

# Spring 2020 MAT303 Recitations

Week of 3/20/20: Sections 3.2 and 3.3

## Section 3.2: General solutions of linear equations

In the first few problems of Homework 6 you're tasked with finding coefficients of a complementary solution to an inhomogeneous equation of type

$$y^{(n)} + p_1(x)y^{(n-1)} + \cdots p_{n-1}(x)y' + p_n(x) = f(x).$$

## Section 3.2: General solutions of linear equations

In the first few problems of Homework 6 you're tasked with finding coefficients of a complementary solution to an inhomogeneous equation of type

$$y^{(n)} + p_1(x)y^{(n-1)} + \cdots p_{n-1}(x)y' + p_n(x) = f(x).$$

Solutions to such equations are obtained by combining a *particular solution*,  $y_p(x)$ , to solutions  $y_c$  of the associated homogeneous equation.

## Section 3.2: General solutions of linear equations

In the first few problems of Homework 6 you're tasked with finding coefficients of a complementary solution to an inhomogeneous equation of type

$$y^{(n)} + p_1(x)y^{(n-1)} + \cdots p_{n-1}(x)y' + p_n(x) = f(x).$$

Solutions to such equations are obtained by combining a *particular solution*,  $y_p(x)$ , to solutions  $y_c$  of the associated homogeneous equation.

$$y(x) = y_c(x) + y_p(x)$$

## Section 3.2: General solutions of linear equations

The example below is extracted from problem 3.2.23 in our textbook.

$$\begin{cases} y'' - 2y' - 3y = 6, \\ y(0) = 3, \\ y'(0) = 11. \end{cases}$$

## Section 3.2: General solutions of linear equations

The example below is extracted from problem 3.2.23 in our textbook.

$$\begin{cases} y'' - 2y' - 3y = 6, \\ y(0) = 3, \\ y'(0) = 11. \end{cases}$$

You are given the forms of complementary and particular solutions,

$$y_c = c_1 e^{-x} + c_2 e^{3x}; \quad y_p = -2$$

Combining these with the initial conditions, a solution to the problem is obtained by solving the algebraic system

$$\begin{cases} c_1 + c_2 = 5, \\ -c_1 + 3c_2 = 11. \end{cases}$$

## Section 3.2: General solutions of linear equations

The example below is extracted from problem 3.2.23 in our textbook.

$$\begin{cases} y'' - 2y' - 3y = 6, \\ y(0) = 3, \\ y'(0) = 11. \end{cases}$$

You are given the forms of complementary and particular solutions,

$$y_c = c_1 e^{-x} + c_2 e^{3x}; \quad y_p = -2$$

Combining these with the initial conditions, a solution to the problem is obtained by solving the algebraic system

$$\begin{cases} c_1 + c_2 = 5, \\ -c_1 + 3c_2 = 11. \end{cases}$$

The solutions are  $c_1 = 1$ ,  $c_2 = 4$ , thus  $y(x) = e^{-x} + 4e^{3x} - 2$ .

## Section 3.2: General solutions of linear equation

Problem 3.2.33 concerns the Wronskian criterion, which we recall below:



## Section 3.2: General solutions of linear equation

Problem 3.2.33 concerns the Wronskian criterion, which we recall below:

### Theorem

*Let  $y_1, y_2, \dots, y_n$  be functions defined on an interval  $I$ , assumed to solve an  $n$ -th order homogeneous, linear differential equation, with continuous coefficients*

$$y^{(n)} + p_1(x)y^{(n-1)} + \cdots p_{n-1}(x)y' + p_n(x) = 0.$$

## Section 3.2: General solutions of linear equation

Problem 3.2.33 concerns the Wronskian criterion, which we recall below:

### Theorem

*Let  $y_1, y_2, \dots, y_n$  be functions defined on an interval  $I$ , assumed to solve an  $n$ -th order homogeneous, linear differential equation, with continuous coefficients*

$$y^{(n)} + p_1(x)y^{(n-1)} + \cdots p_{n-1}(x)y' + p_n(x) = 0.$$

Then either

## Section 3.2: General solutions of linear equation

Problem 3.2.33 concerns the Wronskian criterion, which we recall below:

### Theorem

*Let  $y_1, y_2, \dots, y_n$  be functions defined on an interval  $I$ , assumed to solve an  $n$ -th order homogeneous, linear differential equation, with continuous coefficients*

$$y^{(n)} + p_1(x)y^{(n-1)} + \dots + p_{n-1}(x)y' + p_n(x) = 0.$$

Then either

- (a) Their Wronskian is identically zero on  $I$ , in which case the functions are L.D.

## Section 3.2: General solutions of linear equation

Problem 3.2.33 concerns the Wronskian criterion, which we recall below:

### Theorem

*Let  $y_1, y_2, \dots, y_n$  be functions defined on an interval  $I$ , assumed to solve an  $n$ -th order homogeneous, linear differential equation, with continuous coefficients*

$$y^{(n)} + p_1(x)y^{(n-1)} + \dots + p_{n-1}(x)y' + p_n(x) = 0.$$

Then either

- (a) Their Wronskian is identically zero on  $I$ , in which case the functions are L.D.
- (b) Their Wronskian is nowhere zero on  $I$ , and the functions are L.I.

## Section 3.2: General solutions of linear equation

The following example, extracted from problem 3.2.30 illustrates well the subtleties of this criterion.

## Section 3.2: General solutions of linear equation

The following example, extracted from problem 3.2.30 illustrates well the subtleties of this criterion. Consider the equation

$$x^2 y'' - 2xy' + 2y = 0.$$

## Section 3.2: General solutions of linear equation

The following example, extracted from problem 3.2.30 illustrates well the subtleties of this criterion. Consider the equation

$$x^2 y'' - 2xy' + 2y = 0.$$

It is easy to verify that the functions  $y_1(x) = x, y_2(x) = x^2$  are solutions on  $\mathbb{R}$ . These functions are independent, as if the combination  $y(x) = ax + bx^2$  is null both coefficients must vanish.

## Section 3.2: General solutions of linear equation

The following example, extracted from problem 3.2.30 illustrates well the subtleties of this criterion. Consider the equation

$$x^2 y'' - 2xy' + 2y = 0.$$

It is easy to verify that the functions  $y_1(x) = x, y_2(x) = x^2$  are solutions on  $\mathbb{R}$ . These functions are independent, as if the combination  $y(x) = ax + bx^2$  is null both coefficients must vanish. Meanwhile,

$$W(y_1, y_2) = \det \begin{pmatrix} x & x^2 \\ 1 & 2x \end{pmatrix} = 2x^2 - x^2 = x^2,$$

a function which vanishes when  $x = 0$ .



## Section 3.2: General solutions of linear equation

The following example, extracted from problem 3.2.30 illustrates well the subtleties of this criterion. Consider the equation

$$x^2 y'' - 2xy' + 2y = 0.$$

It is easy to verify that the functions  $y_1(x) = x$ ,  $y_2(x) = x^2$  are solutions on  $\mathbb{R}$ . These functions are independent, as if the combination  $y(x) = ax + bx^2$  is null both coefficients must vanish. Meanwhile,

$$W(y_1, y_2) = \det \begin{pmatrix} x & x^2 \\ 1 & 2x \end{pmatrix} = 2x^2 - x^2 = x^2,$$

a function which vanishes when  $x = 0$ . Why is this not a contradiction?

## Section 3.2: General solutions of linear equations

**Warning:** the Wronskian criterion applies only to solutions

## Section 3.2: General solutions of linear equations

**Warning:** the Wronskian criterion applies only to solutions of equations with continuous coefficients,

## Section 3.2: General solutions of linear equations

**Warning:** the Wronskian criterion applies only to solutions of equations with continuous coefficients, when written in standard form.

The standard form of the previous equation is

$$y'' - \frac{2}{x}y' + \frac{2}{x^2}y = 0,$$

## Section 3.2: General solutions of linear equations

**Warning:** the Wronskian criterion applies only to solutions of equations with continuous coefficients, when written in standard form.

The standard form of the previous equation is

$$y'' - \frac{2}{x}y' + \frac{2}{x^2}y = 0,$$

discontinuous at  $x = 0$ ,

## Section 3.2: General solutions of linear equations

**Warning:** the Wronskian criterion applies only to solutions of equations with continuous coefficients, when written in standard form.

The standard form of the previous equation is

$$y'' - \frac{2}{x}y' + \frac{2}{x^2}y = 0,$$

discontinuous at  $x = 0$ , precisely where the criterion fails.

## Section 3.2: General solutions of linear equations

**Warning:** the Wronskian criterion applies only to solutions of equations with continuous coefficients, when written in standard form.

The standard form of the previous equation is

$$y'' - \frac{2}{x}y' + \frac{2}{x^2}y = 0,$$

discontinuous at  $x = 0$ , precisely where the criterion fails.

As a consequence: there is no linear, second-order differential equation defined on  $\mathbb{R}$  whose solutions are  $x$  and  $x^2$ .

## Section 3.2: General solutions of linear equations

**Warning:** the Wronskian criterion applies only to solutions of equations with continuous coefficients, when written in standard form.

The standard form of the previous equation is

$$y'' - \frac{2}{x}y' + \frac{2}{x^2}y = 0,$$

discontinuous at  $x = 0$ , precisely where the criterion fails.

As a consequence: there is no linear, second-order differential equation defined on  $\mathbb{R}$  whose solutions are  $x$  and  $x^2$ . Think about problem 33 in this context. Why is the Wronskian criterion applicable to the three functions involved? Can you think of a differential equation whose solutions are given by such functions?



## Section 3.3: Homogeneous Equations with Constant Coefficients

The method of characteristic equations assigns to an equation of type

$$a_n y^{(n)} + a_{n-1} y^{(n-1)} + \cdots + a_1 y' + a_0 = 0$$

its characteristic polynomial:

$$p(r) = a_n r^n + a_{n-1} r^{n-1} + \cdots + a_1 r + a_0,$$

where each  $a_i$  is a constant,  $a_n \neq 0$ ,

## Section 3.3: Homogeneous Equations with Constant Coefficients

The method of characteristic equations assigns to an equation of type

$$a_n y^{(n)} + a_{n-1} y^{(n-1)} + \cdots + a_1 y' + a_0 = 0$$

its characteristic polynomial:

$$p(r) = a_n r^n + a_{n-1} r^{n-1} + \cdots + a_1 r + a_0,$$

where each  $a_i$  is a constant,  $a_n \neq 0$ ,

To each (possibly complex) root  $r$  of the polynomial, there corresponds a solution of the differential equation of the form  $e^{rx}$ . If the root has a multiplicity, then new solutions may be generated by monomial multiplication:  $xe^{rx}, x^2e^{rx}, \dots, x^{m-1}e^{rx}$ , where  $m$  is the multiplicity.

## Section 3.3: Homogeneous Equations with Constant Coefficients

The following example is extracted from problem 3.3.12 in our textbook,

$$y^{(4)} - 3y^{(3)} + 3y'' - y' = 0.$$

## Section 3.3: Homogeneous Equations with Constant Coefficients

The following example is extracted from problem 3.3.12 in our textbook,

$$y^{(4)} - 3y^{(3)} + 3y'' - y' = 0.$$

Its characteristic polynomial is

$$p(r) = r^4 - 3r^3 + 3r^2 - r = r(r - 1)^3.$$

The roots are  $r_1 = 0$ , with multiplicity one, and  $r_2 = 1$ , with multiplicity three.

## Section 3.3: Homogeneous Equations with Constant Coefficients

The following example is extracted from problem 3.3.12 in our textbook,

$$y^{(4)} - 3y^{(3)} + 3y'' - y' = 0.$$

Its characteristic polynomial is

$$p(r) = r^4 - 3r^3 + 3r^2 - r = r(r - 1)^3.$$

The roots are  $r_1 = 0$ , with multiplicity one, and  $r_2 = 1$ , with multiplicity three. The general solution of the equation takes the form

$$y(x) = A + Be^x + Cxe^x + Dx^2e^x.$$

## Section 3.3: Homogeneous Equations with Constant Coefficients

The next example is a slight modification of problem 3.3.23:

$$y'' - 8y' + 25y = 0,$$

with initial conditions  $y(0) = 1, y'(0) = 2$ .

## Section 3.3: Homogeneous Equations with Constant Coefficients

The next example is a slight modification of problem 3.3.23:

$$y'' - 8y' + 25y = 0,$$

with initial conditions  $y(0) = 1, y'(0) = 2$ .

The characteristic equation of the problem is

$$r^2 - 8r + 25 = 0,$$

and its solutions are the complex numbers  $r_1 = 4 + 3i, r_2 = 4 - 3i$ .

## Section 3.3: Homogeneous Equations with Constant Coefficients

The next example is a slight modification of problem 3.3.23:

$$y'' - 8y' + 25y = 0,$$

with initial conditions  $y(0) = 1, y'(0) = 2$ .

The characteristic equation of the problem is

$$r^2 - 8r + 25 = 0,$$

and its solutions are the complex numbers  $r_1 = 4 + 3i, r_2 = 4 - 3i$ .

The general real solution of the D.E. takes the form

$$y(x) = e^{4x}(A \cos(3x) + B \sin(3x)).$$



## Section 3.3: Homogeneous Equations with Constant Coefficients

To find the specific values of  $A$  and  $B$ , we impose initial conditions.

## Section 3.3: Homogeneous Equations with Constant Coefficients

To find the specific values of  $A$  and  $B$ , we impose initial conditions. Since

$$y'(x) = 4y(x) - 3e^{4x}(A \sin(3x) - B \cos(3x)),$$

we have at  $x = 0$ :

## Section 3.3: Homogeneous Equations with Constant Coefficients

To find the specific values of  $A$  and  $B$ , we impose initial conditions. Since

$$y'(x) = 4y(x) - 3e^{4x}(A \sin(3x) - B \cos(3x)),$$

we have at  $x = 0$ :

$$A = 1$$

$$4A + 3B = 2,$$

## Section 3.3: Homogeneous Equations with Constant Coefficients

To find the specific values of  $A$  and  $B$ , we impose initial conditions. Since

$$y'(x) = 4y(x) - 3e^{4x}(A \sin(3x) - B \cos(3x)),$$

we have at  $x = 0$ :

$$\begin{aligned} A &= 1 \\ 4A + 3B &= 2, \end{aligned}$$

thus the desired solution is

$$y(x) = e^{4x} \left( \cos(3x) - \frac{2}{3} \sin(3x) \right).$$

## Section 3.3: Homogeneous Equations with Constant Coefficients

In problems 3.3.44 and 3.3.46, you are confronted with differential equations with *complex coefficients*. In this case the roots of the characteristic polynomial need not be conjugates to each other, as the following example (problem 3.3.45) shows.

## Section 3.3: Homogeneous Equations with Constant Coefficients

In problems 3.3.44 and 3.3.46, you are confronted with differential equations with *complex coefficients*. In this case the roots of the characteristic polynomial need not be conjugates to each other, as the following example (problem 3.3.45) shows.

The differential equation

$$y'' - 2iy' + 3y = 0$$

has characteristic equation  $r^2 - 2ir + 3 = 0$ , whose solutions are  $-i$  and  $3i$ .

## Section 3.3: Homogeneous Equations with Constant Coefficients

In problems 3.3.44 and 3.3.46, you are confronted with differential equations with *complex coefficients*. In this case the roots of the characteristic polynomial need not be conjugates to each other, as the following example (problem 3.3.45) shows.

The differential equation

$$y'' - 2iy' + 3y = 0$$

has characteristic equation  $r^2 - 2ir + 3 = 0$ , whose solutions are  $-i$  and  $3i$ . The complex-valued solutions to the D.E. are:

$$y_1(x) = e^{-ix} = \cos(x) - i \sin(x),$$

and

$$y_2(x) = e^{3ix} = \cos(3x) + i \sin(3x).$$

## Section 3.3: Homogeneous Equations with Constant Coefficients

In problem 3.3.52 you are tasked with solving an Euler equation by means of a logarithmic substitution.



## Section 3.3: Homogeneous Equations with Constant Coefficients

In problem 3.3.52 you are tasked with solving an Euler equation by means of a logarithmic substitution. Let's see this procedure in practice by solving problem 3.3.53.

## Section 3.3: Homogeneous Equations with Constant Coefficients

In problem 3.3.52 you are tasked with solving an Euler equation by means of a logarithmic substitution. Let's see this procedure in practice by solving problem 3.3.53.

Consider the equation

$$x^2 y'' + 7xy' + 25y = 0.$$

## Section 3.3: Homogeneous Equations with Constant Coefficients

In problem 3.3.52 you are tasked with solving an Euler equation by means of a logarithmic substitution. Let's see this procedure in practice by solving problem 3.3.53.

Consider the equation

$$x^2 y'' + 7xy' + 25y = 0.$$

Using the substitution  $v = \log(x)$ ,  $x > 0$ , we get

## Section 3.3: Homogeneous Equations with Constant Coefficients

In problem 3.3.52 you are tasked with solving an Euler equation by means of a logarithmic substitution. Let's see this procedure in practice by solving problem 3.3.53.

Consider the equation

$$x^2 y'' + 7xy' + 25y = 0.$$

Using the substitution  $v = \log(x)$ ,  $x > 0$ , we get

$$\begin{aligned}\frac{dy}{dx} &= \frac{1}{x} \frac{dy}{dv} \\ \frac{d^2 y}{dx^2} &= -\frac{1}{x^2} \frac{dy}{dv} + \frac{1}{x^2} \frac{d^2 y}{dv^2}\end{aligned}$$

## Section 3.3: Homogeneous Equations with Constant Coefficients

Rephrasing the original equation in terms of  $v$  yields

$$\frac{d^2y}{dv^2} + 6\frac{dy}{dv} + 25y = 0,$$

whose solutions (obtained via characteristic equation) take the form

$$y(v) = e^{-3v}(A \cos(4v) + B \sin(4v)),$$

## Section 3.3: Homogeneous Equations with Constant Coefficients

Rephrasing the original equation in terms of  $v$  yields

$$\frac{d^2y}{dv^2} + 6\frac{dy}{dv} + 25y = 0,$$

whose solutions (obtained via characteristic equation) take the form

$$y(v) = e^{-3v}(A \cos(4v) + B \sin(4v)),$$

or, in terms of  $x$ ,

$$y(x) = x^{-3}(A \cos(4 \log(x)) + B \sin(4 \log(x))).$$