

Spring 2020 MAT303 Recitations

Week of 4/6/20: Sections 3.4 and 3.5

Section 3.4: Mechanical Vibrations

Problems in this section deal with second-order differential equations of type

$$mx''(t) + cx'(t) + kx(t) = F(t),$$

Section 3.4: Mechanical Vibrations

Problems in this section deal with second-order differential equations of type

$$mx''(t) + cx'(t) + kx(t) = F(t),$$

where m, c, k are constants ($m \neq 0$).

Section 3.4: Mechanical Vibrations

Problems in this section deal with second-order differential equations of type

$$mx''(t) + cx'(t) + kx(t) = F(t),$$

where m, c, k are constants ($m \neq 0$). In today's recitation, we will discuss a few examples of such equations.

Section 3.4: Mechanical vibrations

The first few homework problems deal with homogeneous equations.

Section 3.4: Mechanical vibrations

The first few homework problems deal with homogeneous equations. We learned how to express solutions as combinations of sines, cosines, and exponentials in the last section, but now we adopt a different convention regarding trigonometric representations.

Section 3.4: Mechanical vibrations

The first few homework problems deal with homogeneous equations. We learned how to express solutions as combinations of sines, cosines, and exponentials in the last section, but now we adopt a different convention regarding trigonometric representations. We will write

$$A \cos(\omega_1 t) + B \sin(\omega_1 t) = C \cos(\omega_1 t - \alpha),$$

Section 3.4: Mechanical vibrations

The first few homework problems deal with homogeneous equations. We learned how to express solutions as combinations of sines, cosines, and exponentials in the last section, but now we adopt a different convention regarding trigonometric representations. We will write

$$A \cos(\omega_1 t) + B \sin(\omega_1 t) = C \cos(\omega_1 t - \alpha),$$

where

- ▶ $C = \sqrt{A^2 + B^2}$ the amplitude;

Section 3.4: Mechanical vibrations

The first few homework problems deal with homogeneous equations. We learned how to express solutions as combinations of sines, cosines, and exponentials in the last section, but now we adopt a different convention regarding trigonometric representations. We will write

$$A \cos(\omega_1 t) + B \sin(\omega_1 t) = C \cos(\omega_1 t - \alpha),$$

where

- ▶ $C = \sqrt{A^2 + B^2}$ the amplitude;
- ▶ $\omega_1 = \frac{\sqrt{4cm - k^2}}{2m}$ is the relative frequency;

Section 3.4: Mechanical vibrations

The first few homework problems deal with homogeneous equations. We learned how to express solutions as combinations of sines, cosines, and exponentials in the last section, but now we adopt a different convention regarding trigonometric representations. We will write

$$A \cos(\omega_1 t) + B \sin(\omega_1 t) = C \cos(\omega_1 t - \alpha),$$

where

- ▶ $C = \sqrt{A^2 + B^2}$ the amplitude;
- ▶ $\omega_1 = \frac{\sqrt{4cm - k^2}}{2m}$ is the relative frequency;
- ▶ $\alpha \in [0, 2\pi)$ is the phase, so that $\cos(\alpha) = \frac{A}{C}$ and $\sin(\alpha) = \frac{B}{C}$.

Section 3.4: Mechanical Vibrations

Problem 3.4.4 concerns the movement of a mass-spring system, with:

Section 3.4: Mechanical Vibrations

Problem 3.4.4 concerns the movement of a mass-spring system, with:

- ▶ $m = 250g = 0.25kg$
- ▶ $k = 9N/0.25m = 36N/m$.
- ▶ at $t = 0s$, $x(0) = 1m$, $x'(0) = -5m/s$.

The movement of the system is described by the equation

$$0.25x'' + 36x = 0 \Leftrightarrow x'' + 144x = 0,$$

Section 3.4: Mechanical Vibrations

Problem 3.4.4 concerns the movement of a mass-spring system, with:

- ▶ $m = 250g = 0.25kg$
- ▶ $k = 9N/0.25m = 36N/m$.
- ▶ at $t = 0s$, $x(0) = 1m$, $x'(0) = -5m/s$.

The movement of the system is described by the equation

$$0.25x'' + 36x = 0 \Leftrightarrow x'' + 144x = 0,$$

thus the relative frequency is $\omega = 12$.

Section 3.4: Mechanical Vibrations

Problem 3.4.4 concerns the movement of a mass-spring system, with:

- ▶ $m = 250g = 0.25kg$
- ▶ $k = 9N/0.25m = 36N/m$.
- ▶ at $t = 0s$, $x(0) = 1m$, $x'(0) = -5m/s$.

The movement of the system is described by the equation

$$0.25x'' + 36x = 0 \Leftrightarrow x'' + 144x = 0,$$

thus the relative frequency is $\omega = 12$. A solution can be written in form

$$x(t) = C \cos(12t - \alpha),$$

for constants $C > 0$, $\alpha \in [0, 2\pi)$ to be determined.

Section 3.4: Mechanical Vibrations

Finding these constants amounts to using the initial conditions.

Section 3.4: Mechanical Vibrations

Finding these constants amounts to using the initial conditions.
From $x(0) = 1$, $x'(0) = -5$, we obtain

$$C \cos(\alpha) = 1, \quad C \sin(\alpha) = -\frac{5}{12},$$

Section 3.4: Mechanical Vibrations

Finding these constants amounts to using the initial conditions.
From $x(0) = 1$, $x'(0) = -5$, we obtain

$$C \cos(\alpha) = 1, \quad C \sin(\alpha) = -\frac{5}{12},$$

$$\text{thus } C = \sqrt{1 + \left(\frac{5}{12}\right)^2} = \frac{13}{12}.$$

Section 3.4: Mechanical Vibrations

Finding these constants amounts to using the initial conditions.
From $x(0) = 1$, $x'(0) = -5$, we obtain

$$C \cos(\alpha) = 1, \quad C \sin(\alpha) = -\frac{5}{12},$$

thus $C = \sqrt{1 + \left(\frac{5}{12}\right)^2} = \frac{13}{12}$. The value of α (in radians) that satisfies the constraints above is found by calculator:

$$\alpha \approx 5.89.$$

Thus

$$x(t) \approx \frac{13}{12} \cos(12t - 5.89).$$

Section 3.4: Mechanical Vibrations

Finding these constants amounts to using the initial conditions. From $x(0) = 1$, $x'(0) = -5$, we obtain

$$C \cos(\alpha) = 1, \quad C \sin(\alpha) = -\frac{5}{12},$$

thus $C = \sqrt{1 + \left(\frac{5}{12}\right)^2} = \frac{13}{12}$. The value of α (in radians) that satisfies the constraints above is found by calculator:

$$\alpha \approx 5.89.$$

Thus

$$x(t) \approx \frac{13}{12} \cos(12t - 5.89).$$

Amplitude: $A = \frac{13}{12}$ meters. Period: $T = \frac{2\pi}{12} = \frac{\pi}{6}$ seconds.

Section 3.4: Mechanical Vibrations

In problem 3.4.14, we have a mass-spring-dashpot system with constants

$$m = 25, c = 10, k = 226,$$

and initial data $x(0) = 20, x'(0) = 41$.

Section 3.4: Mechanical Vibrations

In problem 3.4.14, we have a mass-spring-dashpot system with constants

$$m = 25, c = 10, k = 226,$$

and initial data $x(0) = 20, x'(0) = 41$. Its characteristic polynomial is

$$25r^2 + 10r + 226 = (5r + 1)^2 + 15^2,$$

whose roots are

$$r = -\frac{1}{5} \pm 3i.$$

Section 3.4: Mechanical Vibrations

In problem 3.4.14, we have a mass-spring-dashpot system with constants

$$m = 25, c = 10, k = 226,$$

and initial data $x(0) = 20, x'(0) = 41$. Its characteristic polynomial is

$$25r^2 + 10r + 226 = (5r + 1)^2 + 15^2,$$

whose roots are

$$r = -\frac{1}{5} \pm 3i.$$

Solutions to the differential equation take the form

$$x(t) = Ce^{-\frac{t}{5}} \cos(3t - \alpha).$$

Section 3.4: Mechanical Vibrations

To use the initial data, we compute the first derivative,

$$x'(t) = -Ce^{-\frac{t}{5}} \left[\frac{\cos(3t - \alpha)}{5} + 3 \sin(3t - \alpha) \right],$$

Section 3.4: Mechanical Vibrations

To use the initial data, we compute the first derivative,

$$x'(t) = -Ce^{-\frac{t}{5}} \left[\frac{\cos(3t - \alpha)}{5} + 3 \sin(3t - \alpha) \right],$$

hence

$$\begin{aligned} C \cos(\alpha) &= 20 \\ C \left[-\frac{\cos(\alpha)}{5} + 3 \sin(\alpha) \right] &= 41, \end{aligned}$$

Section 3.4: Mechanical Vibrations

To use the initial data, we compute the first derivative,

$$x'(t) = -Ce^{-\frac{t}{5}} \left[\frac{\cos(3t - \alpha)}{5} + 3 \sin(3t - \alpha) \right],$$

hence

$$\begin{aligned} C \cos(\alpha) &= 20 \\ C \left[-\frac{\cos(\alpha)}{5} + 3 \sin(\alpha) \right] &= 41, \end{aligned}$$

from which we infer $C = 25$, $\alpha \approx 0.64$, so

$$x(t) \approx 25e^{-\frac{t}{5}} \cos(3t - 0.64).$$

Section 3.4: Mechanical Vibrations

In problem 3.4.17, we are meant to contrast the damped motion of a spring-mass-dashpot system with its undamped counterpart.

Section 3.4: Mechanical Vibrations

In problem 3.4.17, we are meant to contrast the damped motion of a spring-mass-dashpot system with its undamped counterpart. The differential equation modelling this problem is

$$x'' + 8x' + 16x = 0,$$

Section 3.4: Mechanical Vibrations

In problem 3.4.17, we are meant to contrast the damped motion of a spring-mass-dashpot system with its undamped counterpart. The differential equation modelling this problem is

$$x'' + 8x' + 16x = 0,$$

with initial data $x(0) = 5, x'(0) = -10$.

The characteristic polynomial of the problem is

$$r^2 + 8r + 16 = (r + 4)^2.$$

Its root is -4 , with multiplicity 2, thus this motion is critically damped.

Section 3.4: Mechanical Vibrations

A solution takes the form

$$x(t) = e^{-4t}(At + B),$$

Section 3.4: Mechanical Vibrations

A solution takes the form

$$x(t) = e^{-4t}(At + B),$$

and has derivative

$$x'(t) = e^{-4t}(-4At + A - 4B).$$

Section 3.4: Mechanical Vibrations

A solution takes the form

$$x(t) = e^{-4t}(At + B),$$

and has derivative

$$x'(t) = e^{-4t}(-4At + A - 4B).$$

Subject to initial conditions $x(0) = 5, x'(0) = -10$, we find

$$x(t) = e^{-4t}(2t + 1).$$

Section 3.4: Mechanical Vibrations

The equation that models undamped motion is

$$u''(t) + 16u(t) = 0.$$

Section 3.4: Mechanical Vibrations

The equation that models undamped motion is

$$u''(t) + 16u(t) = 0.$$

Its solutions take the form

$$u(t) = C \cos(4t - \alpha).$$

Section 3.4: Mechanical Vibrations

The equation that models undamped motion is

$$u''(t) + 16u(t) = 0.$$

Its solutions take the form

$$u(t) = C \cos(4t - \alpha).$$

Assuming the same initial conditions $u(0) = 5$, $u'(0) = -10$, we have

$$u(t) \approx \frac{5\sqrt{5}}{2} \cos(4t - 5.82)$$

Section 3.4: Mechanical Vibrations

In problem 3.4.31, we have a mass-spring-dashpot system with constants $m = 1$, $c = 10$ and $k = 125$, subject to initial conditions $x(0) = 6$, $x'(0) = 50$. The characteristic polynomial of the system is

$$r^2 + 10r + 125 = (r + 5)^2 + 10^2,$$

whose roots are $r = -5 \pm 10i$, characterizing underdamped motion. The corresponding solution to the differential equation is

$$x(t) = Ce^{-5t} \cos(10t - \alpha),$$

for constants $C > 0$, $\alpha \in [0, 2\pi)$ to be determined in what follows.

Section 3.4: Mechanical Vibrations

The first derivative of the solution takes the form

$$x'(t) = Ce^{-5t}[-5\cos(10t - \alpha) - 10\sin(10t - \alpha)].$$

Section 3.4: Mechanical Vibrations

The first derivative of the solution takes the form

$$x'(t) = Ce^{-5t}[-5\cos(10t - \alpha) - 10\sin(10t - \alpha)].$$

The initial conditions amount to

$$\begin{aligned} C\cos(\alpha) &= 6, \\ C[-10\cos(\alpha) - 5\sin(\alpha)] &= 50, \end{aligned}$$

Section 3.4: Mechanical Vibrations

The first derivative of the solution takes the form

$$x'(t) = Ce^{-5t}[-5\cos(10t - \alpha) - 10\sin(10t - \alpha)].$$

The initial conditions amount to

$$\begin{aligned} C\cos(\alpha) &= 6, \\ C[-10\cos(\alpha) - 5\sin(\alpha)] &= 50, \end{aligned}$$

from which we infer $C = 10$, $\alpha \approx 0.93$. The solution is thus approximated by

$$x(t) \approx 10e^{-5t}\cos(10t - 0.93).$$

Section 3.4: Mechanical Vibrations

We will contrast this to solutions of the undamped motion, modelled by the equation

$$u'' + 125u = 0.$$

Section 3.4: Mechanical Vibrations

We will contrast this to solutions of the undamped motion, modelled by the equation

$$u'' + 125u = 0.$$

The characteristic polynomial is $p(r) = r^2 + 125$, with roots $r = \pm 5\sqrt{5}i$. Solutions to this equation can be expressed as

$$u(t) = C_* \cos(5\sqrt{5}t - \alpha_*),$$

Section 3.4: Mechanical Vibrations

We will contrast this to solutions of the undamped motion, modelled by the equation

$$u'' + 125u = 0.$$

The characteristic polynomial is $p(r) = r^2 + 125$, with roots $r = \pm 5\sqrt{5}i$. Solutions to this equation can be expressed as

$$u(t) = C_* \cos(5\sqrt{5}t - \alpha_*),$$

where the constants $C_* > 0$ and $\alpha_* \in [0, 2\pi)$ are determined by the initial conditions $u(0) = 6$, $u'(0) = 50$.

Section 3.4: Mechanical Vibrations

The derivative of a solution takes the form

$$u'(t) = -5\sqrt{5}C_* \sin(5\sqrt{5}t - \alpha_*),$$

Section 3.4: Mechanical Vibrations

The derivative of a solution takes the form

$$u'(t) = -5\sqrt{5}C_* \sin(5\sqrt{5}t - \alpha_*),$$

so using the initial data we obtain

$$C \cos(\alpha_*) = 6$$

$$5\sqrt{5}C_* \sin(\alpha_*) = 50,$$

Section 3.4: Mechanical Vibrations

The derivative of a solution takes the form

$$u'(t) = -5\sqrt{5}C_* \sin(5\sqrt{5}t - \alpha_*),$$

so using the initial data we obtain

$$C \cos(\alpha_*) = 6$$

$$5\sqrt{5}C_* \sin(\alpha_*) = 50,$$

and infer $C_* = 2\sqrt{14}$, $\alpha_* \approx 0.64$. Finally, the motion of the undamped system is

$$u(t) \approx 2\sqrt{14} \cos(5\sqrt{5}t - 0.64).$$

Section 3.5: Homogeneous Equations with Constant Coefficients

In problem 3.5.3, we consider the equation

$$y'' - y' - 6y = 2 \sin(3x). \quad (1)$$

Section 3.5: Homogeneous Equations with Constant Coefficients

In problem 3.5.3, we consider the equation

$$y'' - y' - 6y = 2 \sin(3x). \quad (1)$$

The associated homogeneous equation has characteristic polynomial

$$r^2 - r - 6 = (r + 2)(r - 3),$$

Section 3.5: Homogeneous Equations with Constant Coefficients

In problem 3.5.3, we consider the equation

$$y'' - y' - 6y = 2 \sin(3x). \quad (1)$$

The associated homogeneous equation has characteristic polynomial

$$r^2 - r - 6 = (r + 2)(r - 3),$$

with roots $r_1 = -2$, $r_2 = 3$. The corresponding solutions to the complementary equation take the form

$$y_c = A_1 e^{-2x} + A_2 e^{3x}.$$

Section 3.5: Homogeneous Equations with Constant Coefficients

As the solutions to the homogeneous problem and the inhomogeneous term of equation (1) are linearly independent (verify!), we can use the method of undetermined coefficients to guess a particular solution of the form

$$y_p = C_1 \sin(3x) + C_2 \cos(3x).$$

Section 3.5: Homogeneous Equations with Constant Coefficients

As the solutions to the homogeneous problem and the inhomogeneous term of equation (1) are linearly independent (verify!), we can use the method of undetermined coefficients to guess a particular solution of the form

$$y_p = C_1 \sin(3x) + C_2 \cos(3x).$$

To find the coefficients C_1, C_2 , we need to compute the derivatives of the particular solution,

$$\begin{aligned} y_p' &= 3C_1 \cos(3x) - 3C_2 \sin(3x), \\ y_p'' &= -9C_1 \sin(3x) - 9C_2 \cos(3x). \end{aligned}$$

Section 3.5: Homogeneous Equations with Constant Coefficients

Combining all of this data into the equation we get

$$(-15C_1 + 3C_2)\sin(3x) + (-3C_1 - 15C_2)\cos(3x) = 2\sin(3x),$$

Section 3.5: Homogeneous Equations with Constant Coefficients

Combining all of this data into the equation we get

$$(-15C_1 + 3C_2)\sin(3x) + (-3C_1 - 15C_2)\cos(3x) = 2\sin(3x),$$

from which we infer $C_1 = -\frac{5}{39}$, $C_2 = \frac{1}{39}$.

Section 3.5: Homogeneous Equations with Constant Coefficients

Combining all of this data into the equation we get

$$(-15C_1 + 3C_2)\sin(3x) + (-3C_1 - 15C_2)\cos(3x) = 2\sin(3x),$$

from which we infer $C_1 = -\frac{5}{39}$, $C_2 = \frac{1}{39}$. A general solution to the differential equation takes the form

$$y(x) = A_1 e^{-2x} + A_2 e^{3x} - \frac{5}{39} \sin(3x) + \frac{1}{39} \cos(3x),$$

Section 3.5: Homogeneous Equations with Constant Coefficients

Combining all of this data into the equation we get

$$(-15C_1 + 3C_2)\sin(3x) + (-3C_1 - 15C_2)\cos(3x) = 2\sin(3x),$$

from which we infer $C_1 = -\frac{5}{39}$, $C_2 = \frac{1}{39}$. A general solution to the differential equation takes the form

$$y(x) = A_1 e^{-2x} + A_2 e^{3x} - \frac{5}{39} \sin(3x) + \frac{1}{39} \cos(3x),$$

for constants A_1 and A_2 to be determined based upon initial conditions.