

Online Resource 1

Supplementary figures and tables

Article: [Article title]

Journal: Cognitive Neurodynamics

Authors: Marlon Valmórbida Cendron, Wilfredo Blanco Figuerola, Flávio Freitas Barbosa

Corresponding author: Marlon Valmórbida Cendron, Laboratory for Memory and Cognition Studies, Department of Psychology, Federal University of Paraíba, João Pessoa, Brazil. E-mail: marlonvcendron@gmail.com

Cell	<i>In Vivo</i> Firing Rate (Hz)	Reference	Model Firing Rate (Hz)	<i>In Vivo</i> Sparsity	Reference	Model Sparsity
● Mature granule	1.2 (0.56-2.38)	McHugh et al. (2022)	1.08	2-5%	Galloni et al. (2022); Hainmueller and Bartos (2018) GoodSmith et al. (2017)	8.0%
	0.49	Diamantaki et al. (2016)		9%		
	0.031 ± 0.096	Zhang et al. (2020)				
	0.4	Wheeler et al. (2023)				
	0.96 ± 0.19	GoodSmith et al. (2019)				
	≈ 0.2	Senzai and Gyorgy Buzsaki (2016)				
	72 ± 8 ^{max}	Lübke et al. (1998)				
● Immature granule	2.3 (1.6-6.7)	McHugh et al. (2022)	2.96			63.8%
◇ Mossy	0.84	Wheeler et al. (2023)	3.45	≈ 100%	Danielson et al. (2017) GoodSmith et al. (2017)	96.7%
	1.05 ± 0.07	GoodSmith et al. (2019)		88%		
	≈ 0.8	Senzai and Gyorgy Buzsaki (2016)				
	50 ± 6 ^{max}	Lübke et al. (1998)				
● HIPP	—		6.38			100%
▲ Basket	13.6 ± 0.8	Li et al. (2022)	17.35			100%
	230 ± 15 ^{max}	Lübke et al. (1998)				
▲ CA3 pyramidal	0.2	Mizuseki and Buzsaki (2013)	1.17	29%	GoodSmith et al. (2017) Galloni et al. (2022)	30.0%
	0.5	Kay et al. (2016)		20-30%		
	0.72 ± 0.51	Lasztóczy et al. (2011)				
	CA3a: 0.4; CA3b: 0.3	Oliva et al. (2016)				
	1.2	McHugh et al. (2022)				
	0.92 ± 0.06	GoodSmith et al. (2019)				
	≈ 0.6	Senzai and Gyorgy Buzsaki (2016)				
● CA3 inhibitory	20 ± 7	Tukker et al. (2013)	6.51			65.4%
	17 ± 7	Lapray et al. (2012)				
	8.2 ± 5.6	Leão et al. (2012)				

Table S1: *In vivo* firing rates and population sparsity for each cell type, alongside model values (mean). *In vivo* values are mean ± SD. ^{max} maximum observed rate. ≈ approximate value read from figure. Model firing rates are the mean over active cells; values are from the control network, except the immature granule cells (absent in the control), which are from the neurogenesis model at 50% EC→iGC excitability.




Figure S1: Pattern separation of the mature granule cells resolved by input similarity. Each curve shows the pattern separation degree $\mathcal{S}_D$ of the mGC population as a function of the similarity between the input patterns, for the control network without neurogenesis (black dashed line) and for neurogenesis networks at representative EC→iGC excitability levels. Each point is the mean over the 30 pattern sets and the shaded band shows the standard error of the mean. The grey dashed line marks $\mathcal{S}_D = 1$, above which the output patterns are more distinct than their inputs. Separation is strongest for the most similar (most overlapping) inputs, and every neurogenesis network separates better than the control across the whole similarity range.




Figure S2: Population sparsity measured with the Hoyer index, as a cross-validation of the Gini results reported in the main text. Each panel shows the Hoyer sparsity of one population — the whole granule-cell population (All GC), the mature granule cells (mGC), the immature granule cells (iGC) and the pyramidal CA3 cells (pCA3) — as a function of the EC→iGC excitability level; the shaded bands show the standard error of the mean and the dashed line marks the control network without neurogenesis (absent for the iGC panel, which has no counterpart in the control). As with the Gini index, the sparsity of the All GC, mGC and pCA3 populations rises with iGC excitability and stays above the control, whereas the iGC population becomes less sparse.




Figure S3: Orthogonalization of the mature granule-cell output as a function of EC→iGC excitability. The curve shows the orthogonalization degree $\mathcal{O}_D$ of the mGC population, taken as the ratio of the output to the input orthogonalization ($O^{(out)}/O^{(in)}$); the shaded band shows the standard error of the mean and the dashed line marks the control network without neurogenesis. The orthogonalization rises with iGC excitability, indicating that the mGC outputs become increasingly decorrelated as neurogenesis increases, beyond what the reduction in the number of active cells alone would produce.




Figure S4: Population sparsity (Gini index) immediately before and after direct iGC activation, for every EC→iGC excitability level. Each panel corresponds to one excitability level (10%–100%, given in the panel title) and compares the Gini index of the mGC, iGC, combined granule-cell (all GC) and pCA3 populations in the 50 ms windows immediately before (pre) and after (post) the depolarizing pulse, pooled over all runs. Bars show the mean across runs. The result of the paired Wilcoxon signed-rank test between the pre and post values is indicated above each pair (ns, not significant; * $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$).

References

Nathan B. Danielson, Gergely F. Turi, Max Ladow, Spyridon Chavlis, Panagiotis C. Petrantakis, Panayiota Poirazi, and Attila Losonczy. In Vivo Imaging of Dentate Gyrus Mossy Cells in Behaving Mice. *Neuron*, 93(3):552–559.e4, February 2017. ISSN 08966273. doi: 10.1016/j.neuron.2016.12.019.

- Maria Diamantaki, Markus Frey, Philipp Berens, Patricia Preston-Ferrer, and Andrea Burgalossi. Sparse activity of identified dentate granule cells during spatial exploration. *eLife*, 5: e20252, October 2016. ISSN 2050-084X. doi: 10.7554/eLife.20252.
- Alessandro R. Galloni, Aya Samadzelkava, Kiran Hiremath, Reuben Oumnov, and Aaron D. Milstein. Recurrent Excitatory Feedback From Mossy Cells Enhances Sparsity and Pattern Separation in the Dentate Gyrus via Indirect Feedback Inhibition. *Frontiers in Computational Neuroscience*, 16:826278, February 2022. ISSN 1662-5188. doi: 10.3389/fncom.2022.826278.
- Douglas GoodSmith, Xiaojing Chen, Cheng Wang, Sang Hoon Kim, Hongjun Song, Andrea Burgalossi, Kimberly M. Christian, and James J. Knierim. Spatial Representations of Granule Cells and Mossy Cells of the Dentate Gyrus. *Neuron*, 93(3):677–690.e5, February 2017. ISSN 08966273. doi: 10.1016/j.neuron.2016.12.026.
- Douglas GoodSmith, Heekyung Lee, Joshua P. Neunuebel, Hongjun Song, and James J. Knierim. Dentate Gyrus Mossy Cells Share a Role in Pattern Separation with Dentate Granule Cells and Proximal CA3 Pyramidal Cells. *The Journal of Neuroscience*, 39(48):9570–9584, November 2019. ISSN 0270-6474, 1529-2401. doi: 10.1523/JNEUROSCI.0940-19.2019.
- Thomas Hainmueller and Marlene Bartos. Parallel emergence of stable and dynamic memory engrams in the hippocampus. *Nature*, 558(7709):292–296, June 2018. ISSN 0028-0836, 1476-4687. doi: 10.1038/s41586-018-0191-2.
- Kenneth Kay, Marielena Sosa, Jason E. Chung, Mattias P. Karlsson, Margaret C. Larkin, and Loren M. Frank. A hippocampal network for spatial coding during immobility and sleep. *Nature*, 531(7593):185–190, March 2016. ISSN 0028-0836, 1476-4687. doi: 10.1038/nature17144.
- Damien Lapray, Balint Lasztoczi, Michael Lagler, Tim James Viney, Linda Katona, Ornella Valenti, Katja Hartwich, Zolt Borhegyi, Peter Somogyi, and Thomas Klausberger. Behavior-dependent specialization of identified hippocampal interneurons. *Nature Neuroscience*, 15(9): 1265–1271, September 2012. ISSN 1097-6256, 1546-1726. doi: 10.1038/nn.3176.
- Bálint Lasztóczi, John J. Tukker, Peter Somogyi, and Thomas Klausberger. Terminal Field and Firing Selectivity of Cholecystokinin-Expressing Interneurons in the Hippocampal CA3 Area. *The Journal of Neuroscience*, 31(49):18073–18093, December 2011. ISSN 0270-6474, 1529-2401. doi: 10.1523/JNEUROSCI.3573-11.2011.
- Richardson N Leão, Sanja Mikulovic, Katarina E Leão, Hermany Munguba, Henrik Gezelius, Anders Enjin, Kalicharan Patra, Anders Eriksson, Leslie M Loew, Adriano B L Tort, and Klas Kullander. OLM interneurons differentially modulate CA3 and entorhinal inputs to hippocampal CA1 neurons. *Nature Neuroscience*, 15(11):1524–1530, November 2012. ISSN 1097-6256, 1546-1726. doi: 10.1038/nn.3235.
- Rongrong Li, Weiwei Zhang, Jie Zhang, Haibo Zhang, Hui Chen, Zhian Hu, Zhongxiang Yao, Hao Chen, and Bo Hu. Sustained Activity of Hippocampal Parvalbumin-Expressing Interneurons Supports Trace Eyeblick Conditioning in Mice. *The Journal of Neuroscience*, 42(44): 8343–8360, November 2022. ISSN 0270-6474, 1529-2401. doi: 10.1523/JNEUROSCI.0834-22.2022.
- Joachim Lübke, Michael Frotscher, and Nelson Spruston. Specialized Electrophysiological Properties of Anatomically Identified Neurons in the Hilar Region of the Rat Fascia Dentata. *Journal of Neurophysiology*, 79(3):1518–1534, March 1998. ISSN 0022-3077, 1522-1598. doi: 10.1152/jn.1998.79.3.1518.

- Stephen B. McHugh, Vítor Lopes-dos-Santos, Giuseppe P. Gava, Katja Hartwich, Shu K. E. Tam, David M. Bannerman, and David Dupret. Adult-born dentate granule cells promote hippocampal population sparsity. *Nature Neuroscience*, 25(11):1481–1491, November 2022. ISSN 1097-6256, 1546-1726. doi: 10.1038/s41593-022-01176-5.
- Kenji Mizuseki and György Buzsáki. Preconfigured, Skewed Distribution of Firing Rates in the Hippocampus and Entorhinal Cortex. *Cell Reports*, 4(5):1010–1021, September 2013. ISSN 22111247. doi: 10.1016/j.celrep.2013.07.039.
- Azahara Oliva, Antonio Fernández-Ruiz, György Buzsáki, and Antal Berényi. Role of Hippocampal CA2 Region in Triggering Sharp-Wave Ripples. *Neuron*, 91(6):1342–1355, September 2016. ISSN 08966273. doi: 10.1016/j.neuron.2016.08.008.
- Yuta Senzai and Gyorgy Buzsaki. Extracellular recordings of dentate granule cells, mossy cells and CA3 neurons during spatial navigation and sleep in mice, 2016.
- John J. Tukker, Bálint Lasztóczy, Linda Katona, J. David B. Roberts, Eleftheria K. Pissadaki, Yannis Dalezios, László Márton, Limei Zhang, Thomas Klausberger, and Peter Somogyi. Distinct Dendritic Arborization and *In Vivo* Firing Patterns of Parvalbumin-Expressing Basket Cells in the Hippocampal Area CA3. *The Journal of Neuroscience*, 33(16):6809–6825, April 2013. ISSN 0270-6474, 1529-2401. doi: 10.1523/JNEUROSCI.5052-12.2013.
- Diek W. Wheeler, Jeffrey D. Kopsick, Nate Sutton, Carolina Tecuatl, Alexander O. Komedantov, Kasturi Nadella, and Giorgio A. Ascoli. Hippocampome.org v2.0: A knowledge base enabling data-driven spiking neural network simulations of rodent hippocampal circuits, September 2023.
- Xiaomin Zhang, Alois Schlögl, and Peter Jonas. Selective Routing of Spatial Information Flow from Input to Output in Hippocampal Granule Cells. *Neuron*, 107(6):1212–1225.e7, September 2020. ISSN 08966273. doi: 10.1016/j.neuron.2020.07.006.