

M 361K: Real Analysis

Ishan Shah

Fall 2022

Contents

August 25	Page 3
Algebraic Axioms — 3 • Order Axioms — 4	

August 30	Page 6
Upper and Lower Bounds — 6 • Completeness Axiom — 7	

September 6	Page 8
Cardinality — 8 • Countability — 8	

September 8	Page 10
Limits of Sequences — 10	

September 13	Page 12
Monotone Sequences — 12	

September 15	Page 13
Cauchy Sequences — 13	

September 20	Page 14
Empty Set — 14 • Topology — 14 • Closure — 15	

September 22	Page 17
Set Covers — 17 • Cauchy Convergence — 18	

September 27	Page 19
Limits of Functions — 19	

September 29 _____ **Page 21** _____

Sums of Limits — 21 • Continuity of Functions — 22

October 6 _____ **Page 23** _____

Derivatives — 23

October 13 _____ **Page 26** _____

Differentiability and Continuity — 26

October 20 _____ **Page 28** _____

Mean Value Theorem — 28 • Intermediate Value Theorem — 29

October 25 _____ **Page 30** _____

Cauchy Mean Value Theorem — 30 • L'Hospital's Rule — 30 • Taylor's Theorem — 31

October 27 _____ **Page 32** _____

Applications of Taylor's Theorem — 32 • Riemann Integrability — 33

November 3 _____ **Page 35** _____

Partitions — 35

November 8 _____ **Page 38** _____

Tagged Partitions — 38

November 10 _____ **Page 40** _____

Fundamental Theorem of Calculus — 40

Chapter 1

August 25

1.1 Algebraic Axioms

$\forall a, b, c \in \mathbb{R}$

- (A1) $a + b = b + a$.
- (A2) $(a + b) + c = a + (b + c)$.
- (A3) \exists an element $o \in \mathbb{R}$ such that $a + o = o + a = a$.
- (A4) For each element $a \in \mathbb{R}$, \exists an element $(-a) \in \mathbb{R}$ such that $a + (-a) = 0$.
- (M1) $ab = ba$.
- (M2) $(ab)c = a(bc)$.
- (M3) \exists an element $1 \in \mathbb{R}$ such that $a * 1 = 1 * a = a$.
- (M4) For each element $a \in \mathbb{R} \setminus 0$, \exists an element $\frac{1}{a} \in \mathbb{R}$ such that $a * \frac{1}{a} = \frac{1}{a} * a = 1$.
- (D) $a * (b + c) = a * b + a * c$.

Note

If $a = b$ and $c = d$, then $a + c = b + d$ and $a * c = b * d$.

$\forall x, y, z \in \mathbb{R}$:

Theorem 1

If $x + z = y + z$ then $x = y$.

Proof.

$$\begin{aligned}x + z &= y + z \quad (A4) \\(x + z) + (-z) &= (y + z) + (-z) \quad (A2) \\x + (z + (-z)) &= y + (z + (-z)) \quad (A4) \\x + 0 &= y + 0 \quad (A3) \\x &= y\end{aligned}$$

□

Theorem 2

For any $x \in \mathbb{R}$, $x * 0 = 0$.

Proof.

$$\begin{aligned}
 x * 0 &= x * (0 + 0) \\
 x * 0 &= x * 0 + x * 0 \\
 x * 0 + (-x * 0) &= (x * 0 + x * 0) + (-x * 0) \\
 0 &= x * 0 + (x * 0 + (-x * 0)) \\
 &= x * 0 + 0 \\
 &= x * 0
 \end{aligned}$$

□

Theorem 3

$-1 * x = -x$ i.e. $x + (-1) * x = 0$.

Proof.

$$\begin{aligned}
 x + (-1) * x &= x + x * (-1) \\
 &= x * 1 + x * (-1) \\
 &= x * (1 + (-1)) \\
 &= x * 0 \\
 &= 0
 \end{aligned}$$

□

Theorem 4 Zero-product property

$\forall x, y \in \mathbb{R}, x * y = 0 \iff x = 0 \vee y = 0$.

Proof. Let $x, y \in \mathbb{R}$, if $x = 0$ or $y = 0$, then $x * y = 0$. Suppose $x \neq 0$, then we must show $y = 0$. Since $x \neq 0$, $\frac{1}{x}$ exists. Thus, if:

$$\begin{aligned}
 xy &= 0 \\
 \frac{1}{x} * (xy) &= \frac{1}{x} * 0 \\
 (\frac{1}{x} * (xy)) * y &= 0 \\
 1 * y &= 0 \\
 y &= 0
 \end{aligned}$$

□

1.2 Order Axioms

$\forall x, y \in \mathbb{R}$:

- (O1) One of $x < y$, $x > y$ or $x = y$ is true.
- (O2) If $x < y$ and $y < z$, then $x < z$.
- (O3) If $x < y$ then $x + z < y + z$.
- (O4) If $x < y$ and $z > 0$ then $xz < yz$.

Theorem 5

If $x < y$ then $-y < -x$.

Proof.

$$\begin{aligned}
 x &< y \\
 x + (-x + -y) &< y + (-x + -y) \\
 (x + -x) + -y &< (y + -y) + -x \\
 0 + -y &< 0 + -x \\
 -y &< -x
 \end{aligned}$$

□

Theorem 6

If $x < y$ and $z > 0$ then $xz > yz$.

Proof. If $x < y$ and $z > 0$ then $-z < 0$. Thus, $x(-z) < y(-z)$. But,

$$\begin{aligned}
 x(-z) &= x(-1 * z) \\
 &= (x * -1) * z \\
 &= (-1 * x) * z \\
 &= -1(x * z) \\
 &= -x * z
 \end{aligned}$$

Similarly, $y(-z) = -y * z$. Thus, $-x * z < -y * z$, so $xz > yz$.

□

Note

\mathbb{R} is an ordered field. \mathbb{R} is complete, while \mathbb{Q} is not complete.

Chapter 2

August 30

Theorem 7

$\sqrt{2}$ is irrational.

Proof. Suppose not. Suppose that $\sqrt{2}$ is rational. Then $\exists m, n \in \mathbb{Z}$ such that $\sqrt{2} = \frac{m}{n}, n \neq 0$ and m and n share no common factors. Then,

$$2 = \frac{m^2}{n^2}$$
$$2n^2 = m^2$$

Thus, m^2 is even and m is even. Then, $m = 2k$ for some $k \in \mathbb{Z}$. But, by substituting $m = 2k$ into the above equation, we get

$$2n^2 = (2k)^2$$
$$2n^2 = 4k^2$$
$$n^2 = 2k^2$$

Thus, n^2 is even, so n is even. So, n is a perfect square, which is a contradiction. Thus, $\sqrt{2}$ is irrational. \square

2.1 Upper and Lower Bounds

Theorem: Let S be a subset of \mathbb{R} . If there exists a real number m such that $m \geq s \forall s \in S$, m is called an **upper bound** for S . If $m \leq s \forall s \in S$, m is called a **lower bound** for S . **Minimums** and **maximums** must exist in the set to be valid.

$$T = \{q \in \mathbb{Q} \mid 0 \leq q \leq \sqrt{2}\}$$

- Lower bound: -420, -1
- Upper bound: 100, 5, 2
- Minimum: 0
- Maximum: No max

Because rationals are not complete, there is no upper bound for T .

Definition 1: Supremum

The least upper bound of a set is called the supremum of the set.

Definition 2: Infimum

The greatest lower bound of a set is called the infimum of the set.

2.2 Completeness Axiom

Definition 3: Completeness axiom

Every nonempty subset of \mathbb{R} that is bounded above has a least upper bound. That is, $\sup S$ exists and is a real number.

Theorem 8

The set of natural numbers \mathbb{N} is unbounded above.

Proof. Suppose not. Suppose that \mathbb{N} is bounded above. If \mathbb{N} were bounded above, it must have a supremum m . Since $\sup \mathbb{N} = m$, $m - 1$ is not an upper bound. Thus, $\exists n_0 \in \mathbb{N}$ such that $n_0 > m - 1$. But then, $n_0 + 1 > m$. This is a contradiction since $n_0 + 1 \in \mathbb{N}$. Thus, \mathbb{N} is unbounded above. \square

Theorem 9

If A and B are nonempty subsets of \mathbb{R} , let $C = \{x + y \mid x \in A, y \in B\}$. If $\sup A$ and $\sup B$ exist, then $\sup C = \sup A + \sup B$.

Proof. Let $\sup A = a$ and $\sup B = b$. Then if $z \in C$, $z = x + y$ for some $x \in A, y \in B$. Then,

$$z = x + y \leq a + b = \sup A + \sup B$$

By the completeness axiom, \exists a least upper bound of C , $c = \sup C$. It must be that $c \leq a + b$, so we must show $c \geq a + b$. Let $\varepsilon > 0$. Since $a = \sup A$, $a - \varepsilon$ is not an upper bound for A . $\exists x \in A$ such that $a - \varepsilon < x$. Likewise, $\exists y \in B$ such that $b - \varepsilon < y$. Then,

$$(a - \varepsilon) + (b - \varepsilon) = a + b - 2\varepsilon < x + y \leq c$$

Thus, $a + b < c + 2\varepsilon \forall \varepsilon > 0$. So, $a + b \leq c \therefore c = a + b$. \square

Chapter 3

September 6

3.1 Cardinality

Definition 4: Cardinality

The cardinality of a set A is the number of elements in A . We denote this as $|A|$. We say that two sets A and B have the same cardinality if and only if \exists a bijection $f : A \rightarrow B$, or $|A| = |B|$.

Note

This bijection holds true because cardinality is reflexive (via the identity function), symmetric (via the inverse function), and transitive (via composition).

Note

The following examples demonstrate how to prove whether two sets have the same cardinality.

- $|\text{even integers}| = |\text{odd integers}|$: $f(2n) = 2n + 1$.
- $|\mathbb{Z}| = |\mathbb{Z}^+|$: $f(0) = 1, f(1) = 2, f(-1) = 3, f(2) = 4, \dots$
- $|\mathbb{Q}^+| = |\mathbb{Z}^+|$: We can create a diagonal mapping by taking $\frac{n}{m}$ for counting numbers on the rows and columns.
- $|\mathbb{Q}| = |\mathbb{Z}^+|$: $\mathbb{Q} = \mathbb{Q}^+ \cup \mathbb{Q}^- \cup \{0\}$, so we can repeat the diagonal mapping for \mathbb{Q}^- . This is because any subset of a countable set is countable.
- $|\mathbb{Q}| \neq |\mathbb{R}|$: For the real numbers, Cantor's Diagonal Argument proves the sets have different cardinality since no possible surjection exists.

In essence, if we show that there exists some one-to-one mapping between the two sets we can claim that $|A| = |B|$.

3.2 Countability

Definition 5: Countable

If a set is finite or has the same cardinality as \mathbb{N} (i.e. \mathbb{Z}^+), we say that the set is countable.

Theorem 10

Any subset of a countable set is countable.

Theorem 11

Any set that contains an uncountable set is uncountable.

Theorem 12

If $[a_n, b_n] \forall n \in \mathbb{N}$ is a nested sequence of closed bounded intervals, $\exists \delta \in \mathbb{R}$ such that $\delta \in I_n \forall n \in \mathbb{N}$.

Proof. $I_n \subseteq I_1 \forall n \in \mathbb{N}$. Thus, $a_n \subseteq b_1 \forall n \in \mathbb{N}$. So, b_1 is an upper bound for $\{a_n \mid n \in \mathbb{N}\}$. Let δ be the supremum of $\{a_n \mid n \in \mathbb{N}\}$. Thus, $a_n \leq \delta \forall n \in \mathbb{N}$.

We have now shown that $a_n \leq \delta \forall n \in \mathbb{N}$, and we need to show that $\delta \leq b_n \forall n \in \mathbb{N}$. This is left as an exercise for the reader. \square

Note

A nested sequence means that successive subsets contain the previous subset. For example, $[0, 1] \subseteq [0, 2] \subseteq [0, 3] \subseteq \dots$ is a nested sequence.

Theorem 13

$[0, 1]$ is uncountable.

Proof. Assume $[0, 1]$ is countable. That is, $[0, 1] = I = \{x_1, x_2, x_3, \dots\}$. Select a closed interval $I_1 \subseteq I$ such that $x_1 \notin I_1$. Next, select a closed interval $I_2 \subseteq I_1$ such that $x_2 \notin I_2$, and so on. Then, we have

$$I_n \subseteq \dots \subseteq I_2 \subseteq I_1 \subseteq I$$

and $x_n \notin I_n \forall n \in \mathbb{N}$. By **Theorem 3.3**, $\exists \delta \in I$ such that $\delta \in I_n \forall n \in \mathbb{N}$. This implies that $\delta \neq x_n \forall n \in \mathbb{N}$. Thus, $\delta \notin I$, which is a contradiction. Therefore, $[0, 1]$ is uncountable. \square

Chapter 4

September 8

4.1 Limits of Sequences

Definition 6: Limit of a sequence

A sequence a_n is said to converge to a real number s , if for any $\varepsilon > 0$, \exists a real number k such that for all $n \geq k$, the terms a_n satisfy $|a_n - s| < \varepsilon$.

Theorem 14

$$\lim_{n \rightarrow \infty} \frac{1}{\sqrt{n}} = 0.$$

Proof. We need to find some N such that $n > N \forall \varepsilon > 0$.

$$\begin{aligned} \left| \frac{1}{\sqrt{n}} - 0 \right| &< \varepsilon \\ \frac{1}{\sqrt{n}} &< \varepsilon \\ \frac{1}{n} &< \varepsilon^2 \\ n &> \frac{1}{\varepsilon^2} \end{aligned}$$

Let $\varepsilon > 0$ and $N = \frac{1}{\varepsilon^2}$. Then, if $n > N$, we have that

$$\begin{aligned} \left| \frac{1}{\sqrt{n}} - 0 \right| &= \frac{1}{\sqrt{n}} \\ &< \frac{1}{\sqrt{\frac{1}{\varepsilon^2}}} \\ &= \varepsilon \end{aligned}$$

Thus, $\lim_{n \rightarrow \infty} \frac{1}{\sqrt{n}} = 0$. □

Theorem 15

$$\lim_{n \rightarrow \infty} 1 + \frac{1}{2^n} = 1.$$

Proof. Let $\varepsilon > 0$ and $N = \frac{1}{\varepsilon}$. Then, we have

$$\begin{aligned} |1 + \frac{1}{2^n} - 1| &< \varepsilon \\ |\frac{1}{2^n}| &= \frac{1}{2^n} < \frac{1}{n} < \frac{1}{\frac{1}{\varepsilon}} < \varepsilon \\ n &> \frac{1}{\varepsilon} \end{aligned}$$

Thus, $\lim_{n \rightarrow \infty} 1 + \frac{1}{2^n} = 1$. □

Theorem 16

Every convergent sequence is bounded.

Proof. Let S_n be a convergent sequence with a limit s and $\varepsilon = 1$. Then, there exists some N such that $|S_n - s| < 1$ when $n > N$. That is, $|S_n| < |s| + 1$.

Let $M = \max\{S_1, S_2, \dots, S_N, |s| + 1\}$. Then, $|S_n| \leq M$, so S_n is bounded. □

Theorem 17

If a sequence converges, its limit is unique.

Proof. Suppose a sequence S_n converges to s and t . Let $\varepsilon > 0$. Then, $\exists N_1$ such that $|S_n - s| < \frac{\varepsilon}{2}$. For $n > N_1$, $\exists N_2$ such that $|S_n - t| < \frac{\varepsilon}{2}$. For $n > N_2$, let $N = \max\{N_1, N_2\}$. Then, for $n > N$, we have

$$\begin{aligned} |s - t| &= |s + S_n - S_n - t| \\ &= |s - S_n + S_n - t| \\ &\leq |s - S_n| + |S_n - t| \\ &< \frac{\varepsilon}{2} + \frac{\varepsilon}{2} \\ |s - t| &= \varepsilon \end{aligned}$$

Thus, the limit is unique. □

Chapter 5

September 13

5.1 Monotone Sequences

Definition 7: Monotone sequence

A sequence S_n of real numbers is said to be increasing $\iff S_n \leq S_{n+1} \forall n \in \mathbb{N}$ and decreasing $\iff S_n \geq S_{n+1} \forall n \in \mathbb{N}$.

Note

The Fibonacci sequence is an example of an increasing sequence.

Definition 8: Monotone convergence theorem

A monotone sequence is convergent if and only if it is bounded.

Theorem 18

An increasing bounded sequence is convergent.

Proof. Suppose S_n is a bounded increasing sequence. Let S be the set $\{S_n \mid n \in \mathbb{N}\}$. By the completeness axiom, $\sup S$ exists. Let $s = \sup S$. We claim $\lim_{n \rightarrow \infty} S_n = s$. Given $\varepsilon > 0$, $s - \varepsilon$ is not an upper bound for S . Thus, $\exists N \in \mathbb{N}$ such that $S_N > s - \varepsilon$. Furthermore, since S_n is increasing and s is an upper bound for S , we have $s - \varepsilon < S_N \leq S_n \leq s \forall n \geq N$. \square

Note

This is an elementary proof because it only uses axioms to make the conclusion.

Ex. $S_{n+1} = \sqrt{1 + S_n}, S_1 = 1$.

Theorem 19

If S_n is an unbounded increasing sequence, then $\lim_{n \rightarrow \infty} S_n = \infty$.

Proof. Let S_n be an increasing unbounded sequence. Then, $\{S_n \mid n \in \mathbb{N}\}$ is not bounded above, but S is bounded below by S_1 . Thus, given $M \in \mathbb{R}, \exists N \in \mathbb{N}$ such that $S_N > M$. But since S_n is increasing, $S_n > M \forall n > N$. Thus, $\lim_{n \rightarrow \infty} S_n = \infty$. \square

Chapter 6

September 15

6.1 Cauchy Sequences

Definition 9: Cauchy sequence

A sequence of real numbers S_n is called a Cauchy sequence if and only if for each $\varepsilon > 0$, $\exists N$ such that $m, n > N \implies |S_m - S_n| < \varepsilon$.

Note

This means the elements of the sequence get closer to each other as N increases.

Theorem 20

Every convergent sequence is Cauchy.

Proof. Let S_n be a convergent sequence. Then $\exists N$ such that $n > N \implies |S_n - s| < \frac{\varepsilon}{2}$ for some $s \in \mathbb{R}$. Then, for $n, m > N$, we have

$$\begin{aligned} |S_n - S_m| &= |S_n - s + s - S_m| \\ &\leq |S_n - s| + |s - S_m| \\ &< \frac{\varepsilon}{2} + \frac{\varepsilon}{2} \\ &= \varepsilon \end{aligned}$$

Thus, S_n is Cauchy. □

Theorem 21

A sequence of real numbers is Cauchy if and only if it is convergent.

Note

We cannot prove this yet.

Chapter 7

September 20

7.1 Empty Set

Theorem 22

The empty set is a subset of any set.

Proof. Suppose not. That is, suppose $\exists A$ such that $\emptyset \not\subseteq A$. Thus, $\exists x \in \emptyset$ such that $x \notin A$. This is a contradiction because the empty set has no elements. Therefore, $\emptyset \subseteq A$. \square

Theorem 23

There is only one set with no elements.

Proof. Suppose not. That is, suppose \exists two empty sets E_1, E_2 . Then $E_1 \subseteq E_2$ and $E_2 \subseteq E_1$. Thus, $E_1 = E_2$. This is a contradiction because E_1 and E_2 are two different sets. Therefore, there is only one empty set. \square

Note

Closedness of \emptyset The empty set is open and closed (vacuously true).

7.2 Topology

Let $S \subseteq \mathbb{R}$ for the following definitions.

Definition 10: Neighborhood

A neighborhood of x in S can be thought of an varepsilon-sized ball around x , i.e. $N(x, \varepsilon) = \{y \in R \mid 0 \leq |x - y| < \varepsilon\}$.

Definition 11: Deleted neighborhood

A deleted neighborhood is the same as a neighborhood except that x is not included, i.e. $N^*(x, \varepsilon) = \{y \in R \mid 0 < |x - y| < \varepsilon\}$.

Definition 12: Accumulation point

$x \in \mathbb{R}$ is an accumulation point of S if and only if every deleted neighborhood of x contains a point of S .

Note

$(0, \infty)$ has accumulation points $[0, \infty)$. $(0, 1)$ does not contain all of its accumulation points since 0 and 1 are both accumulation points of the set.

Theorem 24

$S \subseteq \mathbb{R}$ is closed if and only if S contains all of its accumulation points.

Proof. Suppose S is closed. Let x be an accumulation point of S . If $x \notin S$, then $x \in S^c$. Thus, \exists a neighborhood N of x such that $N \subseteq S^c$. But $N \cap S = \emptyset$, which contradicts x being an accumulation point of S .

Conversely, suppose S contains all of its accumulation points. Let $x \in S^c$, then x is not an accumulation point of S . Thus, $\exists N^*(x, \varepsilon)$ that misses S . Since $x \notin S$, $N(x, \varepsilon)$ misses S . Therefore, S^c is open, which means S is closed. \square

Theorem 25

If S is a nonempty closed bounded subset of \mathbb{R} , then S has a max.

Proof. Let $s = \sup S$. Then, s is an accumulation point of S . Since S is closed, $s \in S$. Thus, s is a max of S . \square

Definition 13: Interior point

$x \in S$ is an interior point of S if and only if $\exists N(x, t)$ such that $N(x, t) \subset S$.

Definition 14: Boundary point

$x \in S$ is a boundary point of S if and only if every neighborhood N of x has $N \cap S \neq \emptyset$ and $N \cap S^c \neq \emptyset$.

7.3 Closure

Definition 15: Open set

S is an open set if and only if every point in S is an interior point of S . $\forall x \in S, \exists$ a neighborhood $N(x, \varepsilon)$ for some $\varepsilon > 0$ such that $N(x, \varepsilon) \subseteq S$.

Definition 16: Closed set

S is a closed set if and only if S contains at least one of its boundary points. Additionally, S^c must be an open set.

Note

\mathbb{R} is open because all of its points are interior points. \mathbb{R} is also closed because \mathbb{R} has no boundary points, therefore implying that it contains at least one of its boundary points (vacuously true).

Theorem 26

The union of two open sets is open.

Proof. Let A and B be open sets. Let $x \in A \cup B$. Then $x \in A$ or $x \in B$. If $x \in A$, then \exists a neighborhood N_1 of x such that $N_1 \subseteq A$. But then, $N_1 \subseteq A \cup B$. If $x \in B$, then \exists a neighborhood N_2 of x such that $N_2 \subseteq B$. But then, $N_2 \subseteq A \cup B$.

Thus, in either case, \exists a neighborhood N of x such that $N \subseteq A \cup B$. Therefore, $A \cup B$ is open. \square

Theorem 27

An arbitrary union of open sets is open.

Proof. Let A_1, A_2, \dots, A_n be open sets. Let $x \in \bigcup_{i=1}^n A_i$. Then $x \in A_i$ for some i . Let N_i be a neighborhood of x such that $N_i \subseteq A_i$. Then $N_i \subseteq A_i \subseteq \bigcup_{i=1}^n A_i$. Therefore, $\bigcup_{i=1}^n N_i \subseteq \bigcup_{i=1}^n A_i$.

Thus, $\bigcup_{i=1}^n N_i$ is a neighborhood of x such that $\bigcup_{i=1}^n N_i \subseteq \bigcup_{i=1}^n A_i$. Therefore, $\bigcup_{i=1}^n A_i$ is open. \square

Theorem 28

The intersection of two open sets is open.

Proof. Let A and B be open sets. Let $x \in A \cap B$. Then $x \in A$ and $x \in B$. Thus, \exists neighborhoods $N_1(x, \varepsilon_1)$ and $N_2(x, \varepsilon_2)$. Let $\varepsilon = \min\{\varepsilon_1, \varepsilon_2\}$. Then $N_1(x, \varepsilon) \subseteq A$ and $N_2(x, \varepsilon) \subseteq B$.

Thus, $N(x, \varepsilon) \subseteq A \cap B$. Therefore, $A \cap B$ is open. \square

Theorem 29

A finite intersection of open sets is open.

Proof. Let A_1, A_2, \dots, A_n be open sets. Let $x \in \bigcap_{i=1}^n A_i$. Then $x \in A_i$ for all i . Let N_i be a neighborhood of x such that $N_i \subseteq A_i$. Then $N_i \subseteq A_i \subseteq \bigcap_{i=1}^n A_i$. Therefore, $\bigcap_{i=1}^n N_i \subseteq \bigcap_{i=1}^n A_i$.

Thus, $\bigcap_{i=1}^n N_i$ is a neighborhood of x such that $\bigcap_{i=1}^n N_i \subseteq \bigcap_{i=1}^n A_i$. Therefore, $\bigcap_{i=1}^n A_i$ is open. \square

Theorem 30

An arbitrary intersection of open sets is open.

Note

$$\bigcap_{i=1}^{\infty} \left(-\frac{1}{n}, \frac{1}{n}\right) = \emptyset.$$

Chapter 8

September 22

8.1 Set Covers

Definition 17: Open cover

An open cover F of some subset $S \subseteq \mathbb{R}$ is a collection of open sets whose union contains S .

Note

If $E \subseteq F$ and E also covers S , we call E a **subcover**.

Definition 18: Compact

A set S is said to be compact if and only if whenever S is contained in the union of a family F of open sets, then it is contained in a finite number of the sets in F (every open cover has a finite subcover).

Note

It is hard to show that a set is compact since we have to consider every open cover.

Theorem 31 Heine-Borel

A subset S of \mathbb{R} is compact if and only if S is closed and bounded.

Proof. Let S be a compact set. Observe the open cover $(-n, n) \forall n \in \mathbb{N}$. Since S is compact, \exists a finite subcover $(-n_1, n_1), (-n_2, n_2), \dots, (-n_k, n_k)$. \exists one of these sets such that $\bigcup_{i=1}^k (-n_i, n_i) = (-n_m, n_m)$ for some $m = 1, 2, \dots, k$. Thus, $S \subseteq (-n_m, n_m)$, so S is bounded.

Let S be a compact set. Suppose S is not closed. Let p be a boundary point of S , and Let $U_n = \mathbb{R} \setminus [p - \frac{1}{n}, p + \frac{1}{n}] \forall n \in \mathbb{N}$. $S \subseteq \bigcup U_n = \mathbb{R} \setminus p$. \exists a finite subcover n_1, n_2, \dots, n_k such that $S \subseteq \bigcup_{i=1}^k U_{n_i}$. $\exists k$ such that $S \subseteq U_{n_k}$. But, this is a contradiction with p being a boundary point. Therefore, S is closed.

The proof in the other direction is similar, yet non-trivial. \square

Theorem 32 Bolzano-Weierstrass

If a bounded subset S of \mathbb{R} contains infinitely many points, then \exists at least one accumulation point of S .

Proof. Let S be a bounded infinite subset of \mathbb{R} . Suppose S has no accumulation points, then S is closed. By Heine-Borel, S must be compact. Define neighborhoods N_x such that $N_x(x) \cap S = \{x\} \forall x \in S$. Clearly, $S \subseteq \bigcup_x N_x$. But, the collection of all N_x must contain a finite subcover. That is,

$$S \subseteq N_{x_1} \cup N_{x_2} \cup \dots \cup N_{x_k}$$

for some $k \in \mathbb{N}$. This contradicts that S is infinite. Therefore, S has an accumulation point. \square

8.2 Cauchy Convergence

Theorem 33

Every Cauchy sequence is convergent.

Proof. S_n is Cauchy, so $S = \{S_n \mid n \in \mathbb{N}\}$. By Bolzano-Weierstrass, \exists an accumulation point s of S . We claim that $S_n \rightarrow s$. Given $\varepsilon > 0$, $\exists N$ such that $m, n > N$. Then $|S_m - S_n| < \frac{\varepsilon}{2}$. $(S - \frac{\varepsilon}{2}, S + \frac{\varepsilon}{2})$ contains an infinite number of points.

$\exists m > N$ such that $S_m \in N(s, \frac{\varepsilon}{2})$. But then, $|S_n - s| = |S_n - S_m + S_m - s| \leq |S_n - S_m| + |S_m - s| < \frac{\varepsilon}{2} + \frac{\varepsilon}{2} = \varepsilon$. Therefore, $S_n \rightarrow s$. \square

Theorem 34

Let x_n be a sequence of non-negative real numbers. $\sum x_n$ converges if S_k , the sequence of partial sums is bounded.

Proof. $\sum_{n=1}^{\infty} x_n = \lim_{k \rightarrow \infty} S_k$. S_k is increasing and bounded, it is convergent by the monotone convergence theorem. \square

Chapter 9

September 27

9.1 Limits of Functions

Definition 19: Limit of a function

Let $f : D \rightarrow \mathbb{R}$ and let c be an accumulation point of the function. Then, $\lim_{x \rightarrow c} f(x) = L$ if and only if given $\varepsilon > 0$, $\exists \delta > 0$ such that if $|x - c| < \delta$, then $|f(x) - L| < \varepsilon$.

Note

Suppose we want to show that $\lim_{x \rightarrow 2} S_x + 1 = 11$. We are looking for some $\delta > 0$ such that $0 \leq |x - 2| < \delta$ and $|S_x + 1 - 11| < \varepsilon$. This is structured similarly to proofs of limits of sequences. Additionally, the limit must go to an accumulation point of the function because we cannot find the limit of a value outside the function's domain.

Theorem 35

$$\lim_{x \rightarrow 5} 10x + 2 = 52.$$

Proof. We need to find some $\delta > 0$ such that whenever $0 < |x - 5| < \delta$, $|10x + 2 - 52| < \varepsilon$.

$$\begin{aligned} |10x - 50| &< \varepsilon \\ 10|x - 5| &< \varepsilon \\ |x - 5| &< \frac{\varepsilon}{10} \end{aligned}$$

Given $\varepsilon > 0$, let $\delta = \frac{\varepsilon}{10}$. Then, whenever $0 < |x - 5| < \delta$, we have $|10x + 2 - 52| = |10x - 50| = 10|x - 5| < 10 \cdot \frac{\varepsilon}{10} = \varepsilon$. \square

Theorem 36

$$\lim_{x \rightarrow 3} x^2 + 2x + 6 = 21.$$

Proof. We need to find some $\delta > 0$ such that whenever $0 < |x - 3| < \delta$, $|(x^2 + 2x + 6) - 21| < \varepsilon$.

$$\begin{aligned} |x^2 + 2x + 6 - 21| &< \varepsilon \\ |x^2 + 2x - 15| &< \varepsilon \\ |x + 5||x - 3| &< \varepsilon \end{aligned}$$

If $\delta < 1 \implies |x + 5||x - 3| < 9|x - 3| < \varepsilon$. Thus $|x - 3| < \frac{\varepsilon}{9}$. We let $\delta = \min\{1, \frac{\varepsilon}{9}\}$.

Given $\varepsilon > 0$, let $\delta = \min\{1, \frac{\varepsilon}{9}\}$. Then, whenever $0 < |x - 3| < \delta$, we have that $|x + 5| < 9$, thus, $|(x^2 + 2x + 6) - 21| = |x^2 + 2x - 15| = |x + 5||x - 3| < \min\{1, \frac{\varepsilon}{9}\} \cdot \frac{\varepsilon}{9} = \varepsilon$. \square

Note

These proofs have two phases. First, we determine some δ as an upper bound. Then, we show how this choice of δ implies the limit is bounded by some ε .

Theorem 37

Let $f : D \rightarrow \mathbb{R}$ and c is an accumulation point of D . Then, $\lim_{x \rightarrow c} f(x) = L$ if and only if for every sequence $S_n \in D$ such that $S_n \rightarrow c$, $S_n \neq c \forall n$, then $f(S_n)$ converges to L .

Proof. $\lim_{x \rightarrow c} f(x) = L$ and $S_n \rightarrow c \implies f(S_n) \rightarrow L$. We need to find N such that $n > N$ and $|f(S_n) - L| < \varepsilon$. We know that $\exists \delta$ such that $0 < |x - c| < \delta \implies |f(x) - L| < \varepsilon$ and $\exists N$ such that $n > N \implies |S_n - c| < \delta$. Thus, for $n > N$ we have $|f(S_n) - L| < \varepsilon$.

Suppose L is not the limit of f as x approaches c . We must find (S_n) that converges to c , but $f(S_n)$ does not converge to L (contrapositive). $\exists \varepsilon > 0$ such that $\forall \delta > 0, 0 < |x - c| < \delta \implies |f(x) - L| \geq \varepsilon$. For each $n \in \mathbb{N}$, $\exists S_n \in D$ such that $0 < |S_n - c| < \frac{1}{n}$ and $|f(S_n) - L| \geq \varepsilon$. Then, $S_n \rightarrow c$, but $f(S_n) \not\rightarrow L$. This is a contradiction. \square

Chapter 10

September 29

10.1 Sums of Limits

Theorem 38

Let $\lim_{x \rightarrow c} f(x) = L$, $\lim_{x \rightarrow c} g(x) = M$. Then, $\lim_{x \rightarrow c} (f + g)(x) = L + M$.

Proof (Definition 9.1). Given $\varepsilon > 0$, let $\delta_1 > 0$ be such that $0 < |x - c| < \delta_1 \implies |f(x) - L| < \frac{\varepsilon}{2}$. Let $\delta_2 > 0$ be such that $0 < |x - c| < \delta_2 \implies |g(x) - M| < \frac{\varepsilon}{2}$.

Let $\delta = \min\{\delta_1, \delta_2\}$. Then, for $0 < |x - c| < \delta$, we have

$$|f(x) + g(x) - (L + M)| = |(f(x) - L) + (g(x) - M)| \leq |f(x) - L| + |g(x) - M| < \frac{\varepsilon}{2} + \frac{\varepsilon}{2} = \varepsilon.$$

□

Proof (Theorem 9.3). Let $\lim_{x \rightarrow c} f(x) = L$, $\lim_{x \rightarrow c} g(x) = M$, and S_n be a sequence of real numbers such that $S_n \rightarrow c$. Then,

$$\lim_{n \rightarrow \infty} (f + g)(S_n) = \lim_{n \rightarrow \infty} f(S_n) + g(S_n) = \lim_{n \rightarrow \infty} f(S_n) + \lim_{n \rightarrow \infty} g(S_n) = L + M$$

Thus, $\lim_{x \rightarrow c} (f + g)(x) = L + M$.

□

Note

This is true for $-$, \times , and \div as well.

Definition 20: Sequential criterion for functional limits

$\lim_{x \rightarrow c} f(x) = L$ if and only if whenever $S_n \rightarrow c$, $\lim_{n \rightarrow \infty} f(S_n) = L$.

Theorem 39

Let $k \in \mathbb{R}$. If $\lim_{x \rightarrow c} f(x) = L$, then $\lim_{x \rightarrow c} kf(x) = kL$.

Proof. Let $\lim_{x \rightarrow c} f(x) = L$, $k \in \mathbb{R}$, and S_n be a sequence of real numbers such that $S_n \rightarrow c$. Then,

$$\lim_{n \rightarrow \infty} kf(S_n) = k \lim_{n \rightarrow \infty} f(S_n) = kL$$

Thus, $\lim_{x \rightarrow c} kf(x) = kL$.

□

10.2 Continuity of Functions

Definition 21: Continuous function

A function f is continuous at $x = c$ if and only if $\lim_{x \rightarrow c} f(x) = f(c)$. Let s be an accumulation point of the domain $f : D \rightarrow \mathbb{R}$. Then, f is continuous at s if and only if for each $\varepsilon > 0$, $\exists \delta > 0$ such that whenever $0 < |x - s| < \delta$, $|f(x) - f(s)| < \varepsilon$.

Note

Let $f(x) = x \sin(\frac{1}{x})$ where $x \neq 0$, $f(0) = 0$. If we want to show that this function is continuous, we need to find some $\delta > 0$ such that $|x| < \delta \implies |f(x) - f(0)| < \varepsilon$. Let $\delta = \varepsilon$, then when $|x| < \delta$, $|f(x) - f(0)| = |x \sin(\frac{1}{x}) - 0| = |x \sin(\frac{1}{x})| \leq |x| < \varepsilon$.

Theorem 40

If f and g are continuous at $x = c$, then $f + g$ is also continuous at $x = c$.

Proof. Let f and g be continuous at c and S_n be a sequence of real numbers such that $S_n \rightarrow c$. Then,

$$\lim_{n \rightarrow \infty} (f + g)(S_n) = \lim_{n \rightarrow \infty} f(S_n) + \lim_{n \rightarrow \infty} g(S_n) = f(c) + g(c)$$

Thus, $\lim_{x \rightarrow c} (f + g)(x) = (f + g)(c)$. □

Theorem 41

Let $f : D \rightarrow E$ be continuous at $x = c$ and let $g : E \rightarrow R$ be continuous at $x = f(c)$. Then, the composition $g \circ f$ is continuous at $x = c$.

Proof. This is left as an exercise for the reader. □

Chapter 11

October 6

11.1 Derivatives

Definition 22: Derivative

Let f be a real-valued function defined on an open interval containing c . We say f is differentiable at c if $\lim_{x \rightarrow c} \frac{f(x) - f(c)}{x - c}$ exists. We call this limit $f'(c)$.

Theorem 42

If f is differentiable at c , then f is continuous at c .

Proof. Let f be defined on some interval I containing c . Then if f is differentiable at c , if and only if for $x \neq c$,

$$f(x) = (x - c) \frac{f(x) - f(c)}{x - c} + f(c)$$

Then, $\lim_{x \rightarrow c} f(x) = \lim_{x \rightarrow c} (x - c) \frac{f(x) - f(c)}{x - c} + f(c) = \lim_{x \rightarrow c} (x - c) f'(c) + f(c) = f(c)$. Therefore, f is continuous at c . \square

Derivative Rules

- $\frac{d}{dx} kf = k \frac{df}{dx}$
- $\frac{d}{dx} f + g = \frac{df}{dx} + \frac{dg}{dx}$
- $\frac{d}{dx} f \cdot g = \frac{df}{dx} g + \frac{dg}{dx} f$
- $\frac{d}{dx} \frac{f}{g} = \frac{\frac{df}{dx} g - \frac{dg}{dx} f}{g^2}$

Theorem 43 Product rule

$$(fg)' = f'g + fg'$$

Proof. Suppose f and g are differentiable at c . Then,

$$\begin{aligned} \lim_{x \rightarrow c} \frac{(fg)(x) - (fg)(c)}{x - c} &= \lim_{x \rightarrow c} \frac{f(x)g(x) - f(c)g(c)}{x - c} \\ &= \lim_{x \rightarrow c} \frac{f(x)g(x) - f(x)g(c) + f(x)g(c) - f(c)g(c)}{x - c} \\ &= \lim_{x \rightarrow c} \frac{f(x)(g(x) - g(c))}{x - c} + \frac{g(c)(f(x) - f(c))}{x - c} \\ &= f(c)g'(c) + g(c)f'(c) \end{aligned}$$

□

Theorem 44 Quotient rule

$$\left(\frac{f}{g}\right)' = \frac{f'g - fg'}{g^2}$$

Proof. Let f and g be differentiable at c . Then,

$$\begin{aligned} \lim_{x \rightarrow c} \frac{\frac{f}{g}(x) - \frac{f}{g}(c)}{x - c} &= \lim_{x \rightarrow c} \frac{\frac{f(x)}{g(x)} - \frac{f(c)}{g(c)}}{x - c} \\ &= \lim_{x \rightarrow c} \frac{\frac{f(x)g(c) - f(c)g(x)}{g(x)g(c)}}{x - c} \\ &= \lim_{x \rightarrow c} \frac{f(x)g(c) - f(c)g(x)}{(x - c)g(x)g(c)} \\ &= \lim_{x \rightarrow c} \frac{g(c)\frac{f(x) - f(c)}{(x - c)} + f(c)\frac{g(x) - g(c)}{(x - c)}}{g(c)g(x)} \\ &= \lim_{x \rightarrow c} \frac{g(c)f'(c) - f(c)g'(c)}{g^2(c)} \quad \text{Aa} \end{aligned}$$

□

Theorem 45 Power rule

$$(x^n)' = nx^{n-1}f' \quad \forall n \in \mathbb{N}$$

Proof by induction. $p(n) = (x^n)' = nx^{n-1}f'$.

$$p(1): f(x) = x. \lim_{x \rightarrow c} \frac{x - c}{x - c} = 1 = 1 \cdot x^0.$$

$$p(k) \rightarrow p(k + 1):$$

$$\begin{aligned} \frac{d}{dx} x^{k+1} &= \frac{d}{dx} x^k \cdot x \\ &= \left(\frac{d}{dx} x^k\right) \cdot x + x^k \left(\frac{d}{dx} x\right) \\ &= kx^{k-1} \cdot x + x^k \cdot 1 \\ &= kx^k + x^k \\ &= (k + 1)x^k \end{aligned}$$

□

Theorem 46 Chain rule

$$g(f(x))' = g'(f(x)) \cdot f'(x)$$

Proof.

$$\begin{aligned} \lim_{x \rightarrow c} \frac{g(f(x)) - g(f(c))}{x - c} &= \lim_{x \rightarrow c} \frac{g(f(x)) - g(f(c))}{f(x) - f(c)} \frac{f(x) - f(c)}{x - c} \\ &= g'(f(x))f'(x) \end{aligned}$$

□

Note

This will not hold if $f(x) = f(c)$. This is not the full proof.

Chapter 12

October 13

12.1 Differentiability and Continuity

Theorem 47

Let f be defined on an interval I containing c . Then, f is differentiable at c if and only if \exists a function φ on I such that φ is continuous at c and

$$f(x) - f(c) = \varphi(x)(x - c) \forall x \neq c$$

In this case, we have $\varphi(c) = f'(c)$.

Note

Let $f(x) = x^3$. Then, $f(x) - f(c) = x^3 - c^3 = (x^2 + xc + c^2)(x - c)$. $\varphi(c) = c^2 + c \cdot c + c^2 = 3c^2 = f'(c)$.

Proof. If $f'(c)$ exists, we can define φ as

$$\varphi(x) = \begin{cases} \frac{f(x)-f(c)}{x-c} & \text{if } x \neq c \\ f'(c) & \text{if } x = c \end{cases}$$

Then, φ is continuous. Since $\lim_{x \rightarrow c} \varphi(x) = f'(c) = \varphi(c)$. Thus, the function is differentiable. If $x = c$, the equation from the theorem holds as $0 = 0$.

Assume φ is continuous at c and satisfies the equation. Then, continuity of φ implies $\varphi(c) = \lim_{x \rightarrow c} \varphi(x) = \lim_{x \rightarrow c} \frac{f(x)-f(c)}{x-c} \implies \varphi(c) = f'(c)$ since f is differentiable. \square

Theorem 48 Chain rule

$$g(f(c))' = g'(f(c)) \cdot f'(c)$$

Proof. Let $c \in I$. f is continuous at c . Define

$$\varphi(x) = \begin{cases} \frac{g(y)-g(f(c))}{y-f(c)} & \text{if } y \neq f(c) \\ g'(f(c)) & \text{if } y = f(c) \end{cases}$$

Thus, φ is continuous at c . Then,

$$\begin{aligned}\lim_{x \rightarrow c} \varphi(f(x)) &= \varphi(f(c)) = g'(f(c)) \\ g(y) - g(f(c)) &= \varphi(y)(y - f(c)) \\ g(f(x)) - g(f(c)) &= \varphi(f(x))(f(x) - f(c)) \\ \lim_{x \rightarrow c} \frac{g(f(x)) - g(f(c))}{x - c} &= \lim_{x \rightarrow c} \frac{\varphi(f(x))(f(x) - f(c))}{x - c} \\ g'(f(c)) &= \lim_{x \rightarrow c} \varphi(f(x)) \cdot \lim_{x \rightarrow c} \frac{f(x) - f(c)}{x - c} \\ g'(f(c)) &= g'(f(c)) \cdot f'(c)\end{aligned}$$

Thus, the chain rule holds. \square

Theorem 49

If S is a nonempty compact subset of \mathbb{R} , S has a max and a min.

Proof. Let $m = \sup S$ exist by the completeness axiom. Given $t > 0$, $\exists x$ such that $m - t < x < m$. Then, m is an accumulation point of S . But S is closed by Heine-Borel. Thus, $m \in S$.

The same proof holds for the min. \square

Theorem 50

If f is continuous and D is compact, then $f(D)$ is compact. (Note: this will be on the final).

Proof. We know that the inverse of a continuous function is continuous (final exam proof) and that if an open set is continuous its inverse is also continuous (exam 2 proof).

Take an open cover $U = \{u_i\}$ of $f(D)$. Then, $f^{-1}(u_i)$ is an open cover for D . But, only a finite number are needed $(\{u_1, u_2, \dots, u_n\})$. Then, $(\{f(u_1), f(u_2), \dots, f(u_n)\})$ is a finite subcover of u_i for $f(D)$. \square

Theorem 51

Let D be compact and suppose $f : D \rightarrow \mathbb{R}$ is continuous, then f assumes a min and a max.

Proof. Since D is compact, $f(D)$ is compact. Thus, $f(D)$ has a min y_1 and a max y_2 . Since $y_1, y_2 \in f(D)$, $\exists x_1, x_2 \in D$ such that $f(x_1) = y_1$ and $f(x_2) = y_2$. Thus, $f(x_1) \leq f(x) \leq f(x_2) \forall x \in D$. \square

Theorem 52

If f is differentiable on an (a, b) and f assumes a max or min for some $c \in (a, b)$, then $f'(c) = 0$.

Proof. Suppose f assumes its max is at c . That is to say $f(x) \leq f(c) \forall x \in (a, b)$. Let x_n be a sequence converging to c such that $a < x_n < c$. Then,

$$\frac{f(x_n) - f(c)}{x_n - c}$$

converges to $f'(c)$. But, each term is nonnegative. Therefore, the derivative is nonnegative $\implies f'(c) \geq 0$. Now, define y_n as a sequence converging to c such that $c < y_n < b$.

If we look at the sequence $\frac{f(y_n) - f(c)}{y_n - c}$, we see that it converges to $f'(c)$. But, each term is nonpositive. Therefore, the derivative is nonpositive, so $f'(c) \leq 0$. $\therefore 0 \leq f'(c) \leq 0$, so we must have that $f'(c) = 0$. \square

Chapter 13

October 20

13.1 Mean Value Theorem

Theorem 53 Rolle's theorem

Let f be continuous on $[a, b]$ and differentiable on (a, b) , and let $f(a) = f(b)$. Then $\exists c \in (a, b)$ such that $f'(c) = 0$.

Proof. Since f is continuous and $[a, b]$ is compact, $\exists x_1, x_2 \in [a, b]$ such that $f(x_1) \leq f(x) \leq f(x_2) \forall x \in [a, b]$. If x_1 and x_2 are the endpoints of the interval, then f is a compact function, thus $f'(c) = 0 \forall c \in (a, b)$. Otherwise, f contains a max at $x_2 \therefore f'(x_2) = 0$. Thus $\exists c \in (a, b)$ such that $f'(c) = 0$. \square

Theorem 54 Mean value theorem

Let f be continuous on $[a, b]$ and differentiable on (a, b) . Then $\exists c \in (a, b)$ such that $f'(c) = \frac{f(b)-f(a)}{b-a}$.

Proof. Let $g(x)$ be defined as $g(x) = \frac{f(b)-f(a)}{b-a}(x-a) + f(a)$. Let $h(x)$ be the distance from the graph of $f \circ g$. That is, $h = f - g$. Then, h is continuous on $[a, b]$ and differentiable on (a, b) . Furthermore, $h(a) = h(b) = 0$.

By Rolle's Theorem, $\exists c \in (a, b)$ such that $h'(c) = 0$. Thus,

$$0 = h'(c) = f'(c) - g'(c) = f'(c) - \frac{f(b) - f(a)}{b - a}$$

Therefore, $f'(c) = \frac{f(b)-f(a)}{b-a}$. \square

Theorem 55

Let f be continuous on $[a, b]$ and differentiable on (a, b) . Then if $f'(x) = 0 \forall x \in (a, b)$, then f is constant on $[a, b]$.

Proof. Suppose f is not constant. Then, $\exists x_1, x_2$ such that $a \leq x_1 < x_2 \leq b$ and $f(x_1) \neq f(x_2)$. By the Mean Value Theorem, $\exists c \in (x_1, x_2)$ such that

$$f'(c) = \frac{f(x_2) - f(x_1)}{x_2 - x_1} \neq 0$$

But, this is a contradiction. Therefore, f is constant on $[a, b]$. \square

Theorem 56

Let f be differentiable on an interval I . If $f'(x) > 0 \forall x \in I$, then f is strictly increasing on I .

Proof. Suppose $f'(x) > 0 \forall x \in I$ and $x_1, x_2 \in I$ such that $x_1 < x_2$. Mean Value Theorem implies that $\exists c \in (x_1, x_2)$ such that $f'(c) = \frac{f(x_2) - f(x_1)}{x_2 - x_1}$. Which is to say that

$$f(x_2) - f(x_1) = f'(c)(x_2 - x_1)$$

Thus, $f(x_2) - f(x_1)$ is positive since $f'(c)$ and $(x_2 - x_1)$ are both positive. Therefore, f is increasing. \square

13.2 Intermediate Value Theorem

Theorem 57 Intermediate value theorem

Let f be continuous on $[a, b]$ and suppose $f(a) < 0 < f(b)$. Then $\exists c \in (a, b)$ such that $f(c) = 0$.

Proof. Let c be the largest value for which $f(x) \leq 0$. Let $S = \{x \in [a, b] \mid f(x) \leq 0\}$. Since $a \in S$, S is nonempty. Thus, $\sup S = c$ exists.

We claim that $f(c) = 0$. Suppose $f(c) < 0$, then \exists a neighborhood U of c such that $f(x) < 0 \forall x \in U \cap [a, b]$. Now, $c \neq b$ since $f(a) < 0 < f(b)$. Thus, U contains a point p such that $c < p < b$ where $f(p) < 0$. But, this is a contradiction since $p \in S$ and $p > c$. Therefore, $f(c) \not< 0$.

Similarly, suppose $f(c) > 0$. We can follow this proof in the other direction to show that $f(c) = 0$. \square

Note

This is the baby version of the intermediate value theorem. The full version will be asked on exam 2.

Chapter 14

October 25

14.1 Cauchy Mean Value Theorem

Theorem 58 Cauchy mean value theorem

Let f and g be continuous on $[a, b]$ and differentiable on (a, b) . Then, \exists at least one $c \in (a, b)$ such that

$$(f(b) - f(a))g'(c) = (g(b) - g(a))f'(c)$$

Proof. Let $h(x) = (f(b) - f(a))g'(x) - (g(b) - g(a))f'(x) \forall x \in [a, b]$.

Note that

$$\begin{aligned} h(a) &= (f(b) - f(a))g'(a) - (g(b) - g(a))f'(a) = 0 \\ &= f(b)g'(a) - f(a)g'(b) \end{aligned}$$

and

$$\begin{aligned} h(b) &= (f(b) - f(a))g'(b) - (g(b) - g(a))f'(b) = 0 \\ &= f(b)g'(a) - f(a)g'(b) \end{aligned}$$

Thus, h is continuous on $[a, b]$, differentiable on (a, b) , and $h(a) = h(b)$. Therefore, by Rolle's theorem, $\exists c \in (a, b)$ such that $h'(c) = 0$. That is to say,

$$h'(c) = (f(b) - f(a))g''(c) - (g(b) - g(a))f''(c) = 0$$

which implies the desired equality. □

14.2 L'Hospital's Rule

Theorem 59 L'Hospital's rule

Let f and g be continuous on $[a, b]$ and differentiable on (a, b) and $f(c) = g(c) = 0$. Also suppose that $g'(c) \neq 0$ in some neighborhood of c .

If

$$\lim_{x \rightarrow c} \frac{f'(x)}{g'(x)} = L$$

then

$$\lim_{x \rightarrow c} \frac{f(x)}{g(x)} = L$$

Proof. Let x_n be a sequence that converges to c . By the Cauchy mean value theorem \exists a sequence c_n such that c_n is between x_n and c for each n and

$$(f(x_n) - f(c))g'(c_n) = (g(x_n) - g(c))f'(c_n)$$

Thus,

$$\frac{f(x_n)}{g(x_n)} = \frac{f(x_n) - f(c)}{g(x_n) - g(c)} = \frac{f'(c_n)}{g'(c_n)}$$

Furthermore, since $x_n \rightarrow c$ and $c_n \rightarrow c$, we have that if $\lim_{n \rightarrow \infty} \frac{f'(c_n)}{g'(c_n)} = L$, then $\lim_{h \rightarrow \infty} \frac{f(x_n)}{g(x_n)} = \lim_{x \rightarrow c} \frac{f(x)}{g(x)} = L$. \square

14.3 Taylor's Theorem

Theorem 60 Taylor's theorem

Let f and its first n derivatives be continuous on $[a, b]$ (implying that they are also differentiable). Let $x_0 \in [a, b]$. Then, for each $x \in [a, b]$ with $x \neq x_0$, \exists a c between x and x_0 such that

$$f(x) = f(x_0) + f'(x_0)(x - x_0) + \frac{f''(x_0)}{2!}(x - x_0)^2 + \cdots + \frac{f^{(n+1)}(c)}{(n+1)!}(x - x_0)^{n+1}$$

Proof. Let x_0 and x be given and let $J = [x_0, x]$ or $[x, x_0]$. We will define F on J as follows:

$$F(t) = f(x) - f(t) - (x - t)f'(t) - \frac{(x - t)^2}{2!}f''(t) - \cdots - \frac{(x - t)^n}{n!}f^{(n)}(t)$$

Note that

$$F'(t) = \frac{-(x - t)^n}{n!}f^{(n+1)}(t)$$

and define G by

$$G(t) = F(t) - \left(\frac{x - t}{x - x_0} \right)^{n+1} F(x_0)$$

Note that $G(x_0) = 0 = G(x)$. Then, by Rolle's Theorem, $\exists c$ between x and x_0 such that $G'(c) = 0$. That is,

$$0 = G'(c) = F'(c) + (n+1) \frac{(x - c)^n}{(x - x_0)^{n+1}} F(x_0)$$

Hence,

$$\begin{aligned} F(x_0) &= - \left(\frac{1}{n+1} \right) \left(\frac{(x - x_0)^{n+1}}{(x - c)^n} \right) F'(c) \\ &= \left(\frac{1}{n+1} \right) \left(\frac{(x - x_0)^{n+1}}{(x - c)^n} \right) \left(\frac{(x - c)^n}{n!} \right) f^{(n+1)}(c) \\ &= \left(\frac{(x - x_0)^{n+1}}{(n+1)!} \right) f^{(n+1)}(c) \end{aligned}$$

which implies the desired equality. \square

Chapter 15

October 27

15.1 Applications of Taylor's Theorem

Definition 23: Taylor polynomial

We denote a Taylor polynomial $\mathcal{P}_n(x)$ as

$$\mathcal{P}_n(x) = f(x_0) + f'(x_0)(x - x_0) + \frac{f''(x_0)}{2}(x - x_0)^2 + \frac{f^{(n)}(x_0)}{n!}(x - x_0)^n$$

and a remainder term $R_n(x)$ with some $c \in \mathbb{R}$ where $x_0 \leq c \leq x$ as

$$R_n(x) = \frac{f^{(n+1)}(c)}{(n+1)!}(x - x_0)^{n+1}$$

Example 15.1 (E)

Approximate e^x on $[-1, 1]$ using a Taylor polynomial. Let $f(x) = e^x$, $x_0 = 0$ and $n = 5$.

$$\begin{aligned} e^x &= f(0) + f'(0)x + \frac{f''(0)}{2}x^2 + \frac{f'''(0)}{3!}x^3 + \frac{f^{(4)}(0)}{4!}x^4 + \frac{f^{(5)}(0)}{5!}x^5 + \frac{f^{(6)}(0)}{6!}x^6 \\ &= 1 + x + \frac{x^2}{2} + \frac{x^3}{3!} + \frac{x^4}{4!} + \frac{x^5}{5!} + \frac{x^6}{6!} \end{aligned}$$

You can place an upper bound on the remainder term on the interval $[-1, 1]$ ($c = 1$ maxes out $f'(c)$ and $x = 1$ maxes out x^6).

$$|R_5(x)| = \left| \frac{f^{(6)}(c)}{6!}x^6 \right| = \frac{|f^{(6)}(c)|}{6!}|x^6| \leq \frac{e \cdot 1}{6!}$$

Example 15.2 (E)

Approximate $\cos(1)$ to within $1/1000$ using a Taylor polynomial.

Take $x_0 = 0$, on $[-1, 1]$. We need

$$|R_n(x)| = \left| \frac{f^{(n+1)}(c)}{(n+1)!}x^{n+1} \right| \leq \frac{1}{1000}$$

If you find the Taylor polynomial of cosine to the 6th degree,

$$\begin{aligned}\cos(0) &\approx 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \frac{x^6}{6!} \\ \therefore \left| -\frac{x^6}{6!} \right| &\leq \frac{1}{1000} \text{ on } [-1, 1]\end{aligned}$$

Hence, this is a good enough approximation that estimates $\cos(x)$ on $[-1, 1]$ within an error of $1/1000$.

Note

You can estimate π with $\tan^{-1}(1)$ because it equals $\pi/4$.

Theorem 61

e is irrational.

Proof. We know that $e < 3$. Then, we have that

$$0 < e - \left(1 + 1 + \frac{1}{2!} + \frac{1}{3!} + \dots + \frac{1}{n!} \right) < \frac{3}{(n+1)!} < \frac{3}{(n+1)!}$$

We can assume $e = \frac{a}{b}$ where $b \neq 0$ and $a, b \in \mathbb{Z}$. Then,

$$0 < \frac{a}{b} - \left(1 + 1 + \frac{1}{2!} + \frac{1}{3!} + \dots + \frac{1}{n!} \right) < \frac{3}{(n+1)!}$$

Let M be the middle term. Then, take $n > \max\{b, 3\}$. Finally, we have

$$0 < M < a - n! \left(1 + 1 + \frac{1}{2!} + \frac{1}{3!} + \dots + \frac{1}{n!} \right) < \frac{3}{n+1} < \frac{3}{4}$$

This is a contradiction because there is no integer between 0 and $\frac{3}{4}$ even though M is an integer. \square

15.2 Riemann Integrability

Definition 24: Partition

Let $[a, b]$ be an interval in \mathbb{R} . A partition \mathcal{P} of $[a, b]$ is a finite set of points $\{x_0, x_1, \dots, x_n\}$ such that $a = x_0 < x_1 < x_2 < \dots < x_n = b$.

Definition 25: Upper and lower sums

Let $\mathcal{P} = \{x_0, x_1, \dots, x_n\}$ be a partition of $[a, b]$, and let

$$\begin{aligned}M_i(f) &= \sup \{f(x) : [x_{i-1}, x_i]\} \\ m_i(f) &= \inf \{f(x) : [x_{i-1}, x_i]\}\end{aligned}$$

For example, $f(x) = x + 3$, $x_0 = 1$ and $x_1 = 2$. Hence,

$$\begin{aligned}M_1(f) &= 5 \\ m_1(f) &= 4\end{aligned}$$

Let $\Delta x_i = x_i - x_{i-1}$. We then define $U(f, \mathcal{P}) = \sum_{i=1}^n M_i \Delta x_i$ (the upper sum of f with respect to \mathcal{P}) and $L(f, \mathcal{P}) = \sum_{i=1}^n m_i \Delta x_i$ (the lower sum of f with respect to \mathcal{P}). Now, define

$$\begin{aligned}U(f) &= \inf \{U(f, \mathcal{P}) : \mathcal{P} \text{ is a partition of } [a, b]\} \\ L(f) &= \sup \{L(f, \mathcal{P}) : \mathcal{P} \text{ is a partition of } [a, b]\}\end{aligned}$$

Definition 26: Riemann integrable

We say that f is Riemann integrable if and only if $U(f) = L(f)$. In this case, we write

$$\int_a^b f(x)dx = U(f) = L(f)$$

To show a function f is Riemann integrable on $[a, b]$ given $\varepsilon > 0$, we only need to find one partition such that

$$U(f, \mathcal{P}) - L(f, \mathcal{P}) < \varepsilon$$

Chapter 16

November 3

16.1 Partitions

Theorem 62

Let f be bounded on $[a, b]$ if \mathcal{P} and \mathcal{Q} are partitions of $[a, b]$ such that \mathcal{Q} is a refinement of \mathcal{P} . Then,

$$L(f, \mathcal{P}) \leq L(f, \mathcal{Q}) \leq U(f, \mathcal{Q}) \leq U(f, \mathcal{P})$$

Proof. We know that \mathcal{Q} will contain more points than \mathcal{P} . \mathcal{P} is described by $m_k \cdot (x_k - x_{k-1})$ while \mathcal{Q} is described by $m_x \cdot (x^* - x_{k-1}) + m_x \cdot (x_k - x^*)$. \square

Theorem 63

Let \mathcal{P} and \mathcal{Q} be partitions of $[a, b]$. Then

$$L(f, \mathcal{P}) \leq U(f, \mathcal{Q})$$

Proof. Let \mathcal{P} and \mathcal{Q} be partitions of f . Then $\mathcal{P} \cup \mathcal{Q}$ is a refinement of \mathcal{P} and \mathcal{Q} . Thus,

$$L(f, \mathcal{P}) \leq L(f, \mathcal{P} \cup \mathcal{Q}) \leq U(f, \mathcal{P} \cup \mathcal{Q}) \leq U(f, \mathcal{Q})$$

\square

Theorem 64

Let f be bounded on $[a, b]$. Then, $L(f) \leq U(f)$.

Proof. Let \mathcal{P} and \mathcal{Q} be partitions of $[a, b]$. Then by the previous theorem, $U(f, \mathcal{Q})$ is an upper bound for

$$S = \{L(f, \mathcal{P}) : \mathcal{P} \text{ is a partition of } [a, b]\}$$

So, $U(f, \mathcal{Q})$ is at least as large as $\sup S = L(f)$. That is, $L(f) \leq U(f, \mathcal{Q})$ for each partition \mathcal{Q} . Then,

$$L(f) \leq \inf\{U(f, \mathcal{Q}) : \mathcal{Q} \text{ is a partition of } [a, b]\} = U(f)$$

Therefore, $L(f) \leq U(f)$. \square

Example 16.1

$f(x) = x^2$ on $[0, 1]$ with partition $\mathcal{P}_n = \{0, \frac{1}{n}, \frac{2}{n}, \dots, \frac{n-1}{n}, 1\}$.

$$\begin{aligned}
M_i &= \sup \left\{ f(x) : x \in \left[\frac{i-1}{n}, \frac{i}{n} \right] \right\} = \left(\frac{i^2}{n^2} \right) \\
m_i &= \inf \left\{ f(x) : x \in \left[\frac{i-1}{n}, \frac{i}{n} \right] \right\} = \left(\frac{i-1}{n} \right)^2 \\
U(f, \mathcal{P}_n) &= \sum_{i=1}^n M_i \cdot \Delta x_i = \sum_{i=1}^n \left(\frac{i}{n} \right)^2 \cdot \frac{1}{n} = \frac{1}{n^3} \sum_{i=1}^n i^2 = \left[\frac{1}{n^3} \cdot \frac{n(n+1)(2n+1)}{6} \right] \\
L(f, \mathcal{P}_n) &= \sum_{i=1}^n m_i \cdot \Delta x_i = \sum_{i=1}^n \left(\frac{i-1}{n} \right)^2 \cdot \frac{1}{n} = \frac{1}{n^3} \sum_{i=1}^n (i-1)^2 = \left[\frac{1}{n^3} \cdot \frac{n(n-1)(2n-1)}{6} \right]
\end{aligned}$$

Then, $\lim_{n \rightarrow \infty} U(f, \mathcal{P}_n) = \frac{1}{3}$ and $\lim_{n \rightarrow \infty} L(f, \mathcal{P}_n) = \frac{1}{3}$. Thus, $U(f) \leq \frac{1}{3}$ and $L(f) \geq \frac{1}{3}$. Because $L(f) \leq U(f)$, we have that $L(f) = U(f) = \frac{1}{3}$.

Since $L(f) = U(f)$, this function is Riemann integrable. Therefore,

$$\int_0^1 x^2 = \int_0^1 x^2 dx = \frac{1}{3}$$

Theorem 65

Let f be a bounded function on $[a, b]$. Then, f is Riemann integrable if and only if given an $\varepsilon > 0$, \exists a partition of $[a, b]$ such that

$$U(f, \mathcal{P}) - L(f, \mathcal{P}) < \varepsilon$$

Proof. If f is Riemann integrable, since $\varepsilon > 0$, \exists a partition \mathcal{P}_1 such that

$$L(f, \mathcal{P}_1) > L(f) - \frac{\varepsilon}{2}$$

Similarly, $\exists \mathcal{P}_2$ such that

$$U(f, \mathcal{P}_2) < U(f) + \frac{\varepsilon}{2}$$

Let $\mathcal{P} = \mathcal{P}_1 \cup \mathcal{P}_2$. Then,

$$\begin{aligned}
U(f, \mathcal{P}) - L(f, \mathcal{P}) &\leq U(f, \mathcal{P}_2) - L(f, \mathcal{P}_1) \\
&< \left(U(f) + \frac{\varepsilon}{2} \right) - \left(L(f) - \frac{\varepsilon}{2} \right) \\
&= U(f) - L(f) + \varepsilon \\
&= \varepsilon
\end{aligned}$$

Therefore, f is Riemann integrable.

Conversely, given $\varepsilon > 0$, suppose $\exists \mathcal{P}$ such that $U(f, \mathcal{P}) < L(f, \mathcal{P}) + \varepsilon$. Then,

$$U(f, \mathcal{P}) \leq U(f, \mathcal{P}) < L(f, \mathcal{P}) + \varepsilon \leq L(f) + \varepsilon$$

Therefore, $U(f) \leq L(f)$. But then $L(f) = U(f)$, so f is Riemann integrable. \square

Note

Generally, we just need to find some partition \mathcal{P} such that $U(f, \mathcal{P})$ and $L(f, \mathcal{P})$ are within ε of each other.

Theorem 66

Show that $f(x) = x$ is Riemann integrable on $[0, 1]$.

Proof. We need to find a partition \mathcal{P} such that for every $\varepsilon > 0$,

$$U(f, \mathcal{P}) - L(f, \mathcal{P}) < \varepsilon$$

Define $\mathcal{P}_n = \{0, \frac{1}{n}, \frac{2}{n}, \dots, \frac{n-1}{n}, 1\}$. Then,

$$\begin{aligned} U(f, \mathcal{P}_n) - L(f, \mathcal{P}_n) &= \sum_{i=1}^n M_i \Delta x_i - \sum_{i=1}^n m_i \Delta x_i \\ &= \sum_{i=1}^n \left(\frac{i}{n} \right) \cdot \frac{1}{n} - \sum_{i=1}^n \left(\frac{i-1}{n} \right) \cdot \frac{1}{n} \\ &= \frac{1}{n^2} \left(\sum_{i=1}^n i - \sum_{i=1}^n (i-1) \right) \\ &= \frac{1}{n^2} \left(\frac{n(n+1)}{2} - \frac{n(n-1)}{2} \right) \\ &= \frac{1}{n^2} \cdot n \\ &= \frac{1}{n} < \varepsilon \implies n > \frac{1}{\varepsilon} \end{aligned}$$

Thus, we should choose some $n > \frac{1}{\varepsilon}$ so $\mathcal{P} = \mathcal{P}_n$. Therefore, f is Riemann integrable. □

Chapter 17

November 8

17.1 Tagged Partitions

Definition 27: Tagged partition

$\dot{\mathcal{P}}$ is a tagged partition of the form $\{([x_{i-1}, x_i], t_i)\}_{i=1}^n$ where $t_i \in [x_{i-1}, x_i]$. Let $\dot{\mathcal{P}}$ be a tagged partition of $[a, b]$. Then, define the Riemann sum of f with respect to $\dot{\mathcal{P}}$ on $[a, b]$ as

$$S(f, \dot{\mathcal{P}}) = \sum_{i=1}^n f(t_i) \cdot (x_i - x_{i-1})$$

Note

$$\|\dot{\mathcal{P}}\| = \|\mathcal{P}\| = \max\{x_1 - x_0, x_2 - x_1, \dots, x_n - x_{n-1}\}$$

Definition 28: Riemann integrable

A function $f : [a, b] \rightarrow \mathbb{R}$ is said to be Riemann integrable on $[a, b]$ if \exists a number L such that $\forall \varepsilon > 0, \exists \delta > 0$ such that if $\dot{\mathcal{P}}$ is any partition with $\|\dot{\mathcal{P}}\| < \delta$, then

$$|S(f, \dot{\mathcal{P}}) - L| < \varepsilon$$

In this case we say that $\int_a^b f = \int_a^b f(x)dx = L$.

Theorem 67

Every constant function is Riemann integrable on $[a, b]$.

Proof. Given $\varepsilon > 0$, we need to find δ such that $\|\dot{\mathcal{P}}\| < \delta \implies |S(f, \dot{\mathcal{P}}) - k(b-a)| < \varepsilon$. We have that

$$\begin{aligned} S(f, \dot{\mathcal{P}}) &= f(t_1) \cdot \Delta x_1 \\ &= k(b-a) \end{aligned}$$

$$|S(f, \dot{\mathcal{P}}) - k(b-a)| = |k(b-a) - k(b-a)| = 0 < \varepsilon$$

So, we can choose some δ to satisfy this condition. Thus, every constant function is Riemann integrable. \square

Lemma 17.1.1

Let $k \in \mathbb{R}$ and $\dot{\mathcal{P}}$ be a tagged partition, then

$$S(kf, \dot{\mathcal{P}}) = kS(f, \dot{\mathcal{P}})$$

Theorem 68

Let $k \in \mathbb{R}$ and $f \in \mathcal{R}[a, b]$, then

$$\int_a^b kf = k \int_a^b f$$

Proof. Given $\varepsilon < 0$, we need to find δ such that $\|\dot{\mathcal{P}}\| < \delta \implies |S(kf, \dot{\mathcal{P}}) - k \int_a^b f| < \varepsilon$. Since $f \in \mathcal{R}[a, b]$, $\exists \delta$ such that $\|\dot{\mathcal{P}}\| < \delta \implies |S(f, \dot{\mathcal{P}}) - \int_a^b f| < \frac{\varepsilon}{|k|}$. Then, $\forall \dot{\mathcal{P}}$ such that $\|\dot{\mathcal{P}}\| < \delta$, we have that

$$\begin{aligned} \left| S(kf, \dot{\mathcal{P}}) - k \int_a^b f \right| &= \left| kS(f, \dot{\mathcal{P}}) - k \int_a^b f \right| \\ &= \left| kS(f, \dot{\mathcal{P}}) - \int_a^b kf \right| \\ &< \frac{\varepsilon}{|k|} \cdot |k| = \varepsilon \end{aligned}$$

□

Note

Note that $\mathcal{R}[a, b]$ is the set of all Riemann integrable functions on $[a, b]$.

Theorem 69

(On exam 2) If $f, g \in \mathcal{R}[a, b]$, then

$$f + g \in \mathcal{R}[a, b]$$

Proof. We can either find p such that $U(f + g, \mathcal{P}) - L(f + g, \mathcal{P}) < \varepsilon$ to show that $\frac{\varepsilon}{2} + \frac{\varepsilon}{2} < \varepsilon$ OR we can find $\|\dot{\mathcal{P}}\| < \delta$ such that $|S(f + g, \dot{\mathcal{P}}) - \int_a^b f + \int_a^b g| < \varepsilon$ to show that $\frac{\varepsilon}{2} + \frac{\varepsilon}{2} < \varepsilon$. □

Theorem 70

(On exam 2)

$$\left| S(f + g, \dot{\mathcal{P}}) - \left(\int_a^b f + \int_a^b g \right) \right| < \varepsilon$$

Chapter 18

November 10

18.1 Fundamental Theorem of Calculus

Definition 29: Fundamental theorem of calculus

$f(x) : D \rightarrow \mathbb{R}$ is uniformly continuous if and only if given $\varepsilon > 0$, $\exists \delta$ such that $|x - y| < \delta \implies |f(x) - f(y)| < \varepsilon$.

Note

We are closer to proving this!

Theorem 71

A continuous function on a closed interval $[a, b]$ is uniformly continuous.

Note

To be proved.

Theorem 72

Let f be continuous on $[a, b]$. Then, f is Riemann integrable on $[a, b]$.

Proof. Since f is continuous on $[a, b]$, $\exists \delta > 0$ such that when $|x - y| < \delta$, $|f(x) - f(y)| < \frac{\varepsilon}{b-a} \forall \varepsilon > 0$. Let \mathcal{P} be a partition of $[a, b]$ such that $\Delta x_i < \delta \forall i$. On each subinterval $[x_i, x_{i+1}]$, f will obtain a maximum and minimum value at s_i and t_i respectively. Furthermore, $|s_i - t_i| < \delta$, so

$$0 \leq M_i - m_i = f(t_i) - f(s_i) < \frac{\varepsilon}{b-a} \forall i$$

Then,

$$U(f, \mathcal{P}) - L(f, \mathcal{P}) = \sum_{i=1}^n (M_i - m_i) \Delta x_i < \sum_{i=1}^n \frac{\varepsilon}{b-a} \Delta x_i = \frac{\varepsilon}{b-a} (b-a) = \varepsilon$$

□

Theorem 73

If $f \in R[a, c]$ and $f \in R[c, b]$, then $f \in R[a, b]$ and

$$\int_a^b f = \int_a^c f + \int_c^b f$$

Proof. Given $\varepsilon > 0$, \exists a partition \mathcal{P}_1 of $[a, c]$ and \mathcal{P}_2 of $[c, b]$ such that $U(f, \mathcal{P}_1) - L(f, \mathcal{P}_1) < \frac{\varepsilon}{2}$ and $U(f, \mathcal{P}_2) - L(f, \mathcal{P}_2) < \frac{\varepsilon}{2}$. Then, define $\mathcal{P} = \mathcal{P}_1 \cup \mathcal{P}_2$. Then, \mathcal{P} is a partition of $[a, b]$ and

$$\begin{aligned} U(f, \mathcal{P}) - L(f, \mathcal{P}) &= U(f, \mathcal{P}_1) + U(f, \mathcal{P}_2) - L(f, \mathcal{P}_1) - L(f, \mathcal{P}_2) \\ &= U(f, \mathcal{P}_1) - L(f, \mathcal{P}_1) + U(f, \mathcal{P}_2) - L(f, \mathcal{P}_2) \\ &< \frac{\varepsilon}{2} + \frac{\varepsilon}{2} \\ &= \varepsilon \end{aligned}$$

So, $f \in R[a, b]$. Furthermore,

$$\begin{aligned} \int_a^b f &\leq U(f, \mathcal{P}) = U(f, \mathcal{P}_1) + U(f, \mathcal{P}_2) \\ &< L(f, \mathcal{P}_1) + L(f, \mathcal{P}_2) + \varepsilon \\ &\leq \int_a^c f + \int_c^b f + \varepsilon \end{aligned}$$

Similarly,

$$\begin{aligned} \int_a^b f &\geq L(f, \mathcal{P}) = L(f, \mathcal{P}_1) + L(f, \mathcal{P}_2) \\ &> U(f, \mathcal{P}_1) + U(f, \mathcal{P}_2) - \varepsilon \\ &\geq \int_a^c f + \int_c^b f - \varepsilon \end{aligned}$$

Therefore, $\int_a^b f = \int_a^c f + \int_c^b f$. □

Theorem 74

If f is Riemann integrable on $[a, b]$ and g is continuous on $[c, d]$ when $f([a, b]) \subseteq [c, d]$, then $g \circ f$ is Riemann integrable on $[a, b]$.

Note

To be proved.

Theorem 75

Let f be Riemann integrable on a closed interval $[a, b]$. Then, $|f|$ is Riemann integrable on $[a, b]$ and

$$\left| \int_a^b f \right| \leq \int_a^b |f|$$

Proof. $|x|$ is continuous so we can apply the previous theorem. Then,

$$\begin{aligned} -|f(x)| &\leq f(x) \leq |f(x)| \\ -\int_a^b |f| &\leq \int_a^b f \leq \int_a^b |f| \text{ because } \frac{M_i}{m_i} \end{aligned}$$

Thus, $\left| \int_a^b f \right| \leq \int_a^b |f|$. □