

Testing numba/jit accelerated dissipation3 functions

John A. Marohn

Table of contents

1 Preliminaries	1
2 Sample object	1
3 Example friction calculation	3
4 Example BLDS calculation	4
5 Create list of modulation frequencies and charge densities	4
6 Save all figures	7

- Name: dissipation-theory--Study-28.ipynb
- Author: John A. Marohn
- Date: 2024-10-13
- Continued from: dissipation-theory--Study-26.ipynb
- Continued to: dissipation-theory--Study-29.ipynb (probably)
- Summary: Redo Study 26 using numba/jit-accelerated functions coded in `dissipationtheory.dissipation3`. The numba/jit functions run below are 100 to 200 times faster than their pure-Python counterparts run in Study 26, enabling us to plot here a more detailed BLDS spectrum and explore friction and the zero-frequency limit of the BLDS spectrum as a function of many more conductivity points.

1 Preliminaries

2 Sample object

Set up pure Python objects.

```
cantilever = CantileverModel(  
    f_c = ureg.Quantity(75, 'kHz'),  
    k_c = ureg.Quantity(2.8, 'N/m'),  
    V_ts = ureg.Quantity(1, 'V'),  
    R = ureg.Quantity(35, 'nm'),
```

```

d = ureg.Quantity(38, 'nm')
)

```

cantilever

```

    resonance freq = 75.000 kHz
                    = 4.712e+05 rad/s
    spring constant = 2.800 N/m
    tip-sample voltage = 1.000 V
                    radius = 35.000 nm
                    height = 38.000 nm

```

semiconductor

```

    epsilon (real) = 3.000
    epsilon (imag) = -0.200
        thickness = 500.0 nm
        conductivity = 1.000e-05 S/m
        charge density = 1.000e+21 m-3
        reference height = 3.000e+02 nm

    roll-off frequency = 1.129e+06 Hz
        mobility = 6.242e-08 m2/(V s)
    diffusion constant = 1.614e-09 m2/s
        Debye length = 3.780e+01 nm
        diffusion length = 5.852e+01 nm
    effective epsilon (real) = 3.000
    effective epsilon (imag) = -2.597

```

dielectric

```

    epsilon (real) = 1000000.000
    epsilon (imag) = 0.000
        thickness = infinite

```

Magically, set up the jit sample model from the pure Python sample model, as described in Study 27.

cantilever

=====

```

    cantilever freq = 75000.0 Hz
                    = 471238.89803846896 rad/s
    spring constant = 2.8 N/m
    tip-sample voltage = 1.0 V
                    radius = 3.5e-08 m
                    height = 3.8e-08 m

```

```
semiconductor
=====
    epsilon (real) = 3.0
    epsilon (imag) = -0.2
        thickness = 5.000000000000001e-07 m
    conductivity = 1e-05 S/m
    charge density = 1e+21 m{-3}
    reference height = 3.0000000000000004e-07 m

    roll-off frequency = 1129409.0673730192 Hz
        mobility = 6.241509090043337e-08 m2/(V s)
    diffusion constant = 1.6135549206567651e-09 m2/s
        Debye length = 3.7797775481848936e-08 m
    diffusion length = 5.851555252782804e-08 m
effective epsilon (real) = 3.0
effective epsilon (imag) = -2.5966804779363124

dielectric
=====
    epsilon (real) = 1000000.0
    epsilon (imag) = 0.0
        thickness = infinite
```

3 Example friction calculation

Spot-check the new friction-calculation code by comparing a numba/jit result to a pure-Python result.

First, the numba/jit result.

```
CPU times: user 6.99 ms, sys: 231 µs, total: 7.22 ms
Wall time: 7.14 ms
```

```
( )  $\frac{\text{piconewton-second}}{\text{meter}}$ 
```

Now the pure Python result.

```
CPU times: user 2.42 s, sys: 29.5 ms, total: 2.45 s
Wall time: 2.57 s
```

```
( )  $\frac{\text{piconewton-second}}{\text{meter}}$ 
```

By inspection we get the same result for the three terms' contribution to the friction. Success!

The numba/jit calculation is 100 to 200 times faster.

4 Example BLDS calculation

Compare the execution time for pure-Python and numba/jit calculations.

```
CPU times: user 28 ms, sys: 2.12 ms, total: 30.1 ms
Wall time: 31.3 ms
```

```
( ) hertz
```

```
CPU times: user 6.23 s, sys: 55.4 ms, total: 6.28 s
Wall time: 6.5 s
```

```
( ) hertz
```

By inspection we get the same result for the two terms' contribution to the BDLS frequency shift at the . Success!

The numba/jit calculation is 150 to 200 times faster.

5 Create list of modulation frequencies and charge densities

Set up arrays of modulation frequency ω_m and conductivity σ .

Assume a fixed mobility of $\mu = 10^{-8} \text{ m}^2/\text{Vs}$, an ionic mobility.

As we vary σ , we should vary the charge density ρ to keep the mobility constant.

Get ready to run the simulations.

Loop over conductivities, computing the BLDS spectrum and the friction. This takes about

```
CPU times: user 11.9 s, sys: 109 ms, total: 12 s
Wall time: 12.1 s
```

Explore the `data['01']` object. The keys are the conductivity in S/m, running from 10^{-9} to 1.

```
('1e-09', '1.0')
```

The BLDS frequencies are given as a 100×3 array, with 100 being the number of modulation frequencies and 3 being the number of terms in Loring's sum.

```
(100, 3)
```




Figure 1: Broadband local dielectric spectrum of a semi-infinite semiconductor versus charge density.

To get the BLDS frequency, sum up the three components. We do this by summing over the second dimension.

−3.6261448167750916 hertz

From **Study 25**, a helper function to plot the BLDS spectra. Make sure to add up the three terms to get the total BLDS frequency shift.

Plot every 3rd BLDS spectrum.

I have rewritten the **BLDSlimits** function to return low-density and high-density limits for all three terms; see **BLDSzerohigh** and **BLDSzerolow** in **dissipation3**. I want to rewrite the plotting function below, **plotme**, to use these new function.

I think I can consense the code considerably by using **pandas** dataframes. Convert **data['01']** to a **pandas** dataframe and explore.

	1e-09	1.5264179671752366e-09	2
omega_m	[10.0 hertz, 11.768119524349984 hertz, 13.8488...	[10.0 hertz, 11.768119524349984 hertz, 13.8488...	[
f_BLDS	[[-2.6979093482955756 hertz, -0.39730981074967...	[[-2.698090920476028 hertz, -0.397398186774877...	[
gamma	[42.51061503924813 piconewton * second / meter...	[42.519745468636685 piconewton * second / mete...	[
sigma	1e-09 siemens / meter	1.5264179671752366e-09 siemens / meter	2
d	3.8e-08 meter	3.8e-08 meter	3
rho	6.241509090043338e+17 / meter ** 3	9.527151617329711e+17 / meter ** 3	1
LD	1.5129390669059824e-06 meter	1.2245730647396677e-06 meter	9
omega_c	471238.89803846896 hertz	471238.89803846896 hertz	4
omega_0	112.94090673730192 hertz	172.39502927288035 hertz	2

We want to extract the data in the rows, as numbers without units, for plotting. Read the **rho** row and convert it to a numpy array. Loop over the elements of the array, specify the element's units, and get the magnitude of the resulting element. Convert the result list to a numpy array. Print out every 10th element of the resulting array.

```
array([6.24150909e+17, 4.28572501e+19, 2.94278813e+21, 2.02066207e+23,
       1.38748528e+25])
```

Some of the rows return an array of arrays. Run one of these, `omega_m`, through the same procedure. Now we get an array of arrays, stripped of their units.

```
array([1.00000000e+01, 5.09413801e+01, 2.59502421e+02, 1.32194115e+03,
       6.73415066e+03, 3.43046929e+04, 1.74752840e+05, 8.90215085e+05,
       4.53487851e+06, 2.31012970e+07])
```

Generalize this procedure, simultaneously looping of all the rows and specifying each of their units.

Check that the above does what we expect. Look at the shape of the resulting arrays. We have 50 columns, one for each charge density. For the spectra, we have 100 frequencies. Then, for the calculated signals, `f_BLDS` and `gamma`, we have three components, one from each of Loring's terms.

```
(50,)
```

```
(50, 100)
```

```
(50, 100, 3)
```

```
(50, 3)
```

Check out the new code computing the limiting cases.

```
(5, 3)
```

```
(5, 3)
```

Check out the new code for computing the linear approximation to the K_2 term.

```
((60,), (60,))
```


```
(0.001, 3.7649358067924674)
```

Check out the new code for computing the γ_\perp (a) $\rho \rightarrow 0$ limit and (b) linear ρ expansion.

```
((50, 3), (50, 3))
```

Now rewrite the `Study 26` code.

Pass sample parameters via `sample1`, where the parameters have units, not via `sample1_jit`, there the parameters don't have units.



6 Save all figures