

1 - Preliminaries on convex sets and convex functions

Mauro Passacantando

Department of Computer Science, University of Pisa
mauro.passacantando@unipi.it

Optimization Methods and Game Theory
Master of Science in Artificial Intelligence and Data Engineering
University of Pisa – A.Y. 2020/21

Subspaces

Given $x, y \in \mathbb{R}^n$.

A **linear combination** of x and y is a point $\alpha x + \beta y$, where $\alpha, \beta \in \mathbb{R}$.

A set $C \subseteq \mathbb{R}^n$ is a **subspace** if it contains all the linear combinations of any two points in C .

Examples:

- ▶ $\{0\}$
- ▶ any line which passes through zero
- ▶ the solution set of a homogeneous system of linear equations

$$C = \{x \in \mathbb{R}^n : Ax = 0\},$$

where A is a $m \times n$ matrix.

Affine sets

An **affine combination** of x and y is a point $\alpha x + \beta y$, where $\alpha + \beta = 1$.

A set $C \subseteq \mathbb{R}^n$ is an **affine** set if it contains all the affine combinations of any two points in C .

Examples:

- ▶ any single point $\{x\}$
- ▶ any line
- ▶ the solution set of a system of linear equations

$$C = \{x \in \mathbb{R}^n : Ax = b\},$$


where A is a $m \times n$ matrix and $b \in \mathbb{R}^m$

- ▶ any subspace


Convex sets

A **convex combination** of two given points x and y is a point $\alpha x + \beta y$, where $\alpha + \beta = 1$, $\alpha \geq 0$, $\beta \geq 0$.

A set $C \subseteq \mathbb{R}^n$ is **convex** if it contains all the convex combinations of any two points in C .



convex set




non-convex set

Exercise 1.1. Prove that if C is convex, then for any $x^1, \dots, x^k \in C$ and $\alpha_1, \dots, \alpha_k \in (0, 1)$ s.t. $\sum_{i=1}^k \alpha_i = 1$, one has $\sum_{i=1}^k \alpha_i x^i \in C$.

Convex hull

The **convex hull** $\text{conv}(C)$ of a set C is the smallest convex set containing C .



Exercise 1.2. Prove that $\text{conv}(C) = \{\text{all convex combinations of points in } C\}$.

Exercise 1.3. Prove that C is convex if and only if $C = \text{conv}(C)$.

Convex sets - Examples

Examples:

- ▶ subspace
- ▶ affine set
- ▶ line segment
- ▶ halfspace $\{x \in \mathbb{R}^n : a^T x \leq b\}$
- ▶ polyhedron $P = \{x \in \mathbb{R}^n : Ax \leq b\}$ solution set of a system of linear inequalities

Convex sets - Examples

- ball $B(x, r) = \{y \in \mathbb{R}^n : \|y - x\| \leq r\}$, where $\|\cdot\|$ is any norm, e.g.

$$\|x\|_2 = \sqrt{\sum_{i=1}^n x_i^2} \text{ (Euclidean norm)}$$

$$\|x\|_1 = \sum_{i=1}^n |x_i| \text{ (Manhattan distance)}$$

$$\|x\|_\infty = \max_{i=1, \dots, n} |x_i| \text{ (Chebyshev norm)}$$

$$\|x\|_p = \sqrt[p]{\sum_{i=1}^n |x_i|^p}, \text{ with } 1 \leq p < +\infty$$

$$\|x\|_A = \sqrt{x^T A x}, \text{ where } A \text{ is a symmetric and positive definite matrix, i.e.,}$$

$$x^T A x > 0 \quad \forall x \neq 0.$$

Exercise 1.4. Find the unit ball $B(0, 1)$ w.r.t. $\|\cdot\|_1$, $\|\cdot\|_\infty$ and $\|\cdot\|_A$, where

$$A = \begin{pmatrix} 2 & 0 \\ 0 & 1 \end{pmatrix}.$$

Operations that preserve convexity

Sum and difference

If C_1 and C_2 are convex, then $C_1 + C_2 := \{x + y : x \in C_1, y \in C_2\}$ is convex.

If C_1 and C_2 are convex, then $C_1 - C_2 := \{x - y : x \in C_1, y \in C_2\}$ is convex.

Intersection

If C_1 and C_2 are convex, then $C_1 \cap C_2$ is convex.

Exercise 1.5. If $\{C_i\}_{i \in I}$ is any family of convex sets, then $\bigcap_{i \in I} C_i$ is convex.

Union

If C_1 and C_2 are convex, then $C_1 \cup C_2$ is convex?

Closure and interior

If C is convex, then $\text{cl}(C)$ is convex.

If C is convex, then $\text{int}(C)$ is convex.

Operations that preserve convexity

Affine functions

Let $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$ be affine, i.e. $f(x) = Ax + b$, with $A \in \mathbb{R}^{m \times n}$, $b \in \mathbb{R}^m$.

- ▶ If $C \subseteq \mathbb{R}^n$ is convex, then $f(C) = \{f(x) : x \in C\}$ is convex
- ▶ If $C \subseteq \mathbb{R}^m$ is convex, then $f^{-1}(C) = \{x \in \mathbb{R}^n : f(x) \in C\}$ is convex

Examples:

- ▶ scaling, e.g. $f(x) = \alpha x$, with $\alpha > 0$
- ▶ translation, e.g. $f(x) = x + b$, with $b \in \mathbb{R}^n$
- ▶ rotation, e.g. $f(x) = \begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix} x$, with $\theta \in (0, 2\pi)$

Cones

A set $C \subseteq \mathbb{R}^n$ is a **cone** if $\alpha x \in C$ for any $x \in C$ and $\alpha \geq 0$.

Examples:

- ▶ \mathbb{R}_+^n is a convex cone
- ▶ $\{x \in \mathbb{R}^2 : x_1 x_2 = 0\}$ is a non-convex cone
- ▶ Given a polyhedron $P = \{x : Ax \leq b\}$, the recession cone of P is defined as

$$\text{rec}(P) := \{d : x + \alpha d \in P \text{ for any } x \in P, \alpha \geq 0\}.$$

It is easy to prove $\text{rec}(P) = \{x : Ax \leq 0\}$, thus it is a polyhedral cone.

- ▶ $\{x \in \mathbb{R}^3 : x_3 \geq \sqrt{x_1^2 + x_2^2}\}$ is a non-polyhedral cone.

Exercises

1.6. Write the vector $(1, 1)$ as the convex combination of the vectors $(0, 0), (3, 0), (0, 2), (3, 2)$.

1.7. When does one halfspace contain another? Give conditions under which

$$\{x \in \mathbb{R}^n : a_1^T x \leq b_1\} \subseteq \{x \in \mathbb{R}^n : a_2^T x \leq b_2\},$$

where $\|a_1\|_2 = \|a_2\|_2 = 1$. Also find the conditions under which the two halfspaces are equal.

1.8. Which of the following sets are polyhedra?

a) $\{y_1 a_1 + y_2 a_2 : -1 \leq y_1 \leq 1, -1 \leq y_2 \leq 1\}$, where $a_1, a_2 \in \mathbb{R}^n$.

b) $\left\{x \in \mathbb{R}^n : x \geq 0, \sum_{i=1}^n x_i = 1, \sum_{i=1}^n a_i x_i = b_1, \sum_{i=1}^n a_i^2 x_i = b_2\right\}$, where $b_1, b_2, a_1, \dots, a_n \in \mathbb{R}$.


c) $\{x \in \mathbb{R}^n : x \geq 0, a^T x \leq 1 \text{ for all } a \text{ with } \|a\|_2 = 1\}$.

d) $\{x \in \mathbb{R}^n : x \geq 0, a^T x \leq 1 \text{ for all } a \text{ with } \|a\|_1 = 1\}$.

Convex functions

Given a convex set $C \subseteq \mathbb{R}^n$, a function $f : C \rightarrow \mathbb{R}$ is **convex** if

$$f(\alpha y + (1 - \alpha)x) \leq \alpha f(y) + (1 - \alpha)f(x) \quad \forall x, y \in C, x \neq y, \forall \alpha \in (0, 1)$$



f is said **concave** if $-f$ is convex.

Exercise 1.9. Prove that if f is convex, then for any $x^1, \dots, x^k \in C$ and $\alpha_1, \dots, \alpha_k \in (0, 1)$ s.t. $\sum_{i=1}^k \alpha_i = 1$, one has $f\left(\sum_{i=1}^k \alpha_i x^i\right) \leq \sum_{i=1}^k \alpha_i f(x^i)$.

Strictly convex and strongly convex functions

Given a convex set $C \subseteq \mathbb{R}^n$, a function $f : C \rightarrow \mathbb{R}$ is **strictly convex** if

$$f(\alpha y + (1 - \alpha)x) < \alpha f(y) + (1 - \alpha)f(x) \quad \forall x, y \in C, x \neq y, \forall \alpha \in (0, 1)$$

Given a convex set $C \subseteq \mathbb{R}^n$, a function $f : C \rightarrow \mathbb{R}$ is **strongly convex** if there exists $\tau > 0$ s.t.

$$f(\alpha y + (1 - \alpha)x) \leq \alpha f(y) + (1 - \alpha)f(x) - \frac{\tau}{2}\alpha(1 - \alpha)\|y - x\|_2^2$$

$$\forall x, y \in C, x \neq y, \forall \alpha \in (0, 1)$$

Thm. f is strongly convex if and only if $\exists \tau > 0$ s.t. $f(x) - \frac{\tau}{2}\|x\|_2^2$ is convex

Exercise 1.10.

- ▶ Prove that: strongly convex \implies strictly convex \implies convex
- ▶ convex \implies strictly convex ?
- ▶ strictly convex \implies strongly convex ?


First order conditions

Assume that $C \subseteq \mathbb{R}^n$ is open and convex, $f : C \rightarrow \mathbb{R}$ is continuously differentiable.

Theorem

f is **convex** if and only if

$$f(y) \geq f(x) + (y - x)^T \nabla f(x) \quad \forall x, y \in C.$$



First-order approximation of f is a global **underestimator**

First order conditions

Theorem

- ▶ f is **strictly convex** if and only if

$$f(y) > f(x) + (y - x)^T \nabla f(x) \quad \forall x, y \in C, \text{ with } x \neq y.$$

- ▶ f is **strongly convex** if and only if there exists $\tau > 0$ such that

$$f(y) \geq f(x) + (y - x)^T \nabla f(x) + \frac{\tau}{2} \|y - x\|_2^2 \quad \forall x, y \in C.$$

Second order conditions

Assume that $C \subseteq \mathbb{R}^n$ is open and convex, $f : C \rightarrow \mathbb{R}$ is twice continuously differentiable.

Theorem

- ▶ f is **convex** if and only if for all $x \in C$ the Hessian matrix $\nabla^2 f(x)$ is positive semidefinite, i.e.

$$v^T \nabla^2 f(x) v \geq 0 \quad \forall v \neq 0,$$

or, equivalently, the eigenvalues of $\nabla^2 f(x)$ are ≥ 0 .

- ▶ If $\nabla^2 f(x)$ is positive definite for all $x \in C$, then f is **strictly convex**.
- ▶ f is **strongly convex** if and only if there exists $\tau > 0$ such that $\nabla^2 f(x) - \tau I$ is positive semidefinite for all $x \in C$, i.e.

$$v^T \nabla^2 f(x) v \geq \tau \|v\|_2^2 \quad \forall v \neq 0,$$

or, equivalently, the eigenvalues of $\nabla^2 f(x)$ are $\geq \tau$.

Examples

$f(x) = c^T x$ is both convex and concave

$f(x) = \frac{1}{2}x^T Qx + c^T x$ is

- ▶ convex iff Q is positive semidefinite
- ▶ strongly convex iff Q is positive definite
- ▶ concave iff Q is negative semidefinite
- ▶ strongly concave iff Q is negative definite

$f(x) = e^{ax}$ for any $a \in \mathbb{R}$ is strictly convex, but not strongly convex

$f(x) = \log(x)$ is strictly concave, but not strongly concave

$f(x) = x^a$ with $x > 0$ is strictly convex if $a > 1$ or $a < 0$. Is it strongly convex?

$f(x) = x^a$ with $x > 0$ is strictly concave if $0 < a < 1$

$f(x) = \|x\|$ is convex, but not strictly convex

$f(x) = \max\{x_1, \dots, x_n\}$ is convex, but not strictly convex

Exercises

1.11. Prove that the function

$$f(x_1, x_2) = \frac{x_1 x_2}{x_1 - x_2}$$

is convex on the set $\{x \in \mathbb{R}^2 : x_1 - x_2 > 0\}$.

1.12. Prove that $f(x_1, x_2) = \frac{1}{x_1 x_2}$ is convex on the set $\{x \in \mathbb{R}^2 : x_1, x_2 > 0\}$.

1.13. Given a convex set $C \subseteq \mathbb{R}^n$, the distance function is defined as follows:

$$d_C(x) = \inf_{y \in C} \|y - x\|.$$

Prove that d_C is a convex function.

1.14. Given $C = \{x \in \mathbb{R}^2 : x_1^2 + x_2^2 \leq 1\}$, write the distance function d_C explicitly.

1.15. Prove that the arithmetic mean of n positive numbers x_1, \dots, x_n is greater or equal to their geometric mean, i.e.,

$$\frac{x_1 + x_2 + \dots + x_n}{n} \geq \sqrt[n]{x_1 x_2 \dots x_n}.$$

(Hint: exploit the \log function.)

Operations that preserve convexity

Theorem

- ▶ If f is convex and $\alpha > 0$, then αf is convex
- ▶ If f_1 and f_2 are convex, then $f_1 + f_2$ are convex
- ▶ If f is convex, then $f(Ax + b)$ is convex

Examples

- ▶ Log barrier for linear inequalities:

$$f(x) = - \sum_{i=1}^m \log(b_i - a_i^T x) \quad C = \{x \in \mathbb{R}^n : b_i - a_i^T x > 0 \quad \forall i = 1, \dots, m\}$$

- ▶ Norm of affine function: $f(x) = \|Ax + b\|$

Exercise 1.16. If f_1 and f_2 are convex, then is the product $f_1 f_2$ convex?

Pointwise maximum

Theorem

- ▶ If f_1, \dots, f_m are convex, then $f(x) = \max\{f_1(x), \dots, f_m(x)\}$ is convex.
- ▶ If $\{f_i\}_{i \in I}$ is a family of convex functions, then $f(x) = \sup_{i \in I} f_i(x)$ is convex.

Example. If $L(x, \lambda) : \mathbb{R}^n \times \mathbb{R}^m \rightarrow \mathbb{R}$ is convex in x and concave in λ , then

$$\begin{aligned} p(x) &= \sup_{\lambda} L(x, \lambda) && \text{is convex} \\ d(\lambda) &= \inf_x L(x, \lambda) && \text{is concave} \end{aligned}$$

Composition

$f : \mathbb{R}^n \rightarrow \mathbb{R}$ and $g : \mathbb{R} \rightarrow \mathbb{R}$.

Theorem

- ▶ If f is convex and g is convex and nondecreasing, then $g \circ f$ is convex.
- ▶ If f is concave and g is convex and nonincreasing, then $g \circ f$ is convex.
- ▶ If f is concave and g is concave and nondecreasing, then $g \circ f$ is concave.
- ▶ If f is convex and g is concave and nonincreasing, then $g \circ f$ is concave.

Examples

- ▶ If f is convex, then $e^{f(x)}$ is convex
- ▶ If f is concave and positive, then $\log f(x)$ is concave
- ▶ If f is convex, then $-\log(-f(x))$ is convex on $\{x : f(x) < 0\}$
- ▶ If f is concave and positive, then $\frac{1}{f(x)}$ is convex
- ▶ If f is convex and nonnegative, then $f(x)^p$ is convex for all $p \geq 1$

Sublevel sets

Given $f : \mathbb{R}^n \rightarrow \mathbb{R}$ and $\alpha \in \mathbb{R}$, the set

$$S_\alpha(f) = \{x \in \mathbb{R}^n : f(x) \leq \alpha\}$$

is said the α -sublevel set of f .

Exercise 1.17. Prove that if f is convex, then $S_\alpha(f)$ is a convex set for any $\alpha \in \mathbb{R}$.

Is the converse true?

Quasiconvex functions

Given a convex set $C \subseteq \mathbb{R}^n$, a function $f : C \rightarrow \mathbb{R}$ is **quasiconvex** if the α -sublevel sets are convex for all $\alpha \in \mathbb{R}$.

f is said quasiconcave if $-f$ is quasiconvex.

Examples

- ▶ $f(x) = \sqrt{|x|}$ is quasiconvex on \mathbb{R}
- ▶ $f(x_1, x_2) = x_1 x_2$ is quasiconcave on $\{x \in \mathbb{R}^2 : x_1 > 0, x_2 > 0\}$
- ▶ $f(x) = \log x$ is quasiconvex and quasiconcave
- ▶ $f(x) = \text{ceil}(x) = \inf\{z \in \mathbb{Z} : z \geq x\}$ is quasiconvex and quasiconcave

Exercise 1.18

Express each convex set defined below in the form $\bigcap_{i \in I} \{x : f_i(x) \leq 0\}$, where $f_i : \mathbb{R}^n \rightarrow \mathbb{R}$ are suitable convex functions:

a) $\text{conv}\{(-1, -1), (1, 0), (0, 2)\}$

b) $\text{conv}\{(0, 0), (1, 1)\}$

c) $\text{conv}(\{x \in \mathbb{R}^2 : x_1^2 + (x_2 - 1)^2 = 1\} \cup \{x \in \mathbb{R}^2 : x_1^2 + (x_2 + 1)^2 = 1\})$

d) $\text{conv}\{x \in \mathbb{R}^2 : x_1 x_2 = 1\}$