# EL2 Worked Examples: Graphing Exponential Functions

Recording companion for the EL2 CheckIt support video — Math 172, Santa Ana College

## How to use this document

- Each worked example is on its own page, in the same order as the Try It tabs in the interactive explainer, with the same numbers.
- The right-hand third of every page is intentionally blank. It is space for handwriting or ink notation while you narrate (for example, after inserting this PDF as a printout in OneNote).
- Say it at the start of the video: your CheckIt problem will have different numbers than the examples here. The reasoning is what transfers.

## The one question behind every example

**What value does the graph level off to, and does the curve sit above that level or below it?** The level-off value is the horizontal asymptote. Above it, the range runs from the asymptote up. Below it, the range runs from the asymptote down. The domain is all real numbers.

Adapted from TBIL Precalculus (Clontz & Lewis), CC BY-NC-SA 4.0. Drafted with Claude (Anthropic); reviewed by Dr. Marty Romero.

# Example 1: Asymptote 1

$$f(x)=2^x+3$$

## Question

What is the equation of the horizontal asymptote of this function?

**A.** y = 0
**B.** y = 3
**C.** y = −3
**D.** x = 3

## Answer and reasoning

On its own, $2^x$ levels off at $y=0$ as $x$ gets very negative. Adding 3 outside the exponential lifts the whole graph up 3, so the level-off line moves from $y=0$ to $y=3$. It is a horizontal line, so its equation starts with $y$. **The asymptote is $y=3$.**

# Example 2: Asymptote 2

$$f(x)=-3^x-2$$

## Question

What is the equation of the horizontal asymptote of this function?

**A.** x = −2
**B.** y = 2
**C.** y = 0
**D.** y = −2

## Answer and reasoning

The minus sign in front flips the graph over the x-axis, and the line $y=0$ stays where it is. Then subtracting 2 outside the exponential moves the whole graph down 2, so the line moves to $y=-2$. **The asymptote is $y=-2$.**

Adapted from TBIL Precalculus (Clontz & Lewis), CC BY-NC-SA 4.0. Drafted with Claude (Anthropic); reviewed by Dr. Marty Romero.

# Example 3: Asymptote 3

$$f(x)=-2^{x+3}$$

## Question

What is the equation of the horizontal asymptote of this function?

**A.** y = 3
**B.** y = −3
**C.** y = 0
**D.** x = −3

## Answer and reasoning

The *+3* is inside the exponent, so it slides the graph left 3. It does not lift or lower the graph, so the level-off line stays at *y=0*. Nothing is added or subtracted outside the exponential. **The asymptote is *y=0*.**

# Example 4: Asymptote 4

$$f(x)=(1/e)^x-3$$

## Question

What is the equation of the horizontal asymptote of this function?

**A.** y = −3
**B.** y = 3
**C.** $y=1/e$
**D.** x = −3

## Answer and reasoning

The base $1/e$ is only the base of the exponential. It is not added or subtracted outside, so it does not set the asymptote. The $-3$ outside moves the level-off line from $y=0$ down to $y=-3$. **The asymptote is $y=-3$.**

# Example 5: Range 1

$$f(x)=2^x+3$$

## Question

What is the range of this function? Use interval notation.

**A.** $(-\infty, 3)$
**B.** $(3, \infty)$
**C.** $(0, \infty)$
**D.** $(-\infty, \infty)$

## Answer and reasoning

The asymptote is $y=3$. The outputs of $2^x$ are always positive, and adding 3 makes them always greater than 3. They get as close to 3 as you like but never reach it, and they grow without limit. **The range is $(3, \infty)$.**

Adapted from TBIL Precalculus (Clontz & Lewis), CC BY-NC-SA 4.0. Drafted with Claude (Anthropic); reviewed by Dr. Marty Romero.

# Example 6: Range 2

$$f(x)=-3^x-2$$

## Question

What is the range of this function? Use interval notation.

**A.** $(-\infty, -2)$
**B.** $(-2, \infty)$
**C.** $(-\infty, 0)$
**D.** $(2, \infty)$

## Answer and reasoning

The asymptote is $y=-2$. The minus sign out front flips the graph upside down, so the curve lies below the asymptote. The outputs are always less than $-2$, and they fall without limit. **The range is $(-\infty, -2)$.**

Adapted from TBIL Precalculus (Clontz & Lewis), CC BY-NC-SA 4.0. Drafted with Claude (Anthropic); reviewed by Dr. Marty Romero.

# Example 7: Range 3

$$f(x)=(1/2)^x-1$$

## Question

What is the range of this function? Use interval notation.

**A.** $(0, \infty)$
**B.** $(-\infty, -1)$
**C.** $(-1, \infty)$
**D.** $(-\infty, \infty)$

## Answer and reasoning

The asymptote is $y=-1$. The base $1/2$ is the same as $2^{-x}$, which flips the graph left to right, not upside down. The values of $(1/2)^x$ are still positive, so the curve stays above the asymptote. **The range is $(-1, \infty)$.**

# Example 8: Range 4

$$f(x)=-e^{x+1}$$

## Question

What is the range of this function? Use interval notation.

**A.** $(0, \infty)$
**B.** $(-\infty, -1)$
**C.** $(-1, \infty)$
**D.** $(-\infty, 0)$

## Answer and reasoning

Nothing is added or subtracted outside, so the asymptote is $y=0$. The $+1$ in the exponent only slides the graph left. The minus sign out front flips the curve below the x-axis, so the outputs are always negative. **The range is $(-\infty, 0)$.**

# Example 9: Domain 1

$$f(x)=-4^{x-2}+1$$

## Question


What is the domain of this function? Use interval notation.

**A.** $(0, \infty)$
**B.** $(-\infty, \infty)$
**C.** $(2, \infty)$
**D.** $(-\infty, 1)$

## Answer and reasoning

You can put any real number in an exponent, including negative numbers, fractions, and zero. Shifts and flips move the graph around but do not remove any inputs. The number 1 is the asymptote, and it limits the outputs, not the inputs. **The domain is $(-\infty, \infty)$.**

# Example 10: Graph 1




**Plotted data:** a dashed horizontal line at y = −1. A smooth curve lies above the dashed line. Moving left, it gets closer and closer to the dashed line without touching it. Moving right, it rises faster and faster. It passes through (0, 0) and (1, 1).

## Question


Which function has this graph?

- **A.** $f(x)=2^x+1$
- **B.** $f(x)=2^x-1$
- **C.** $f(x)=2^{x-1}$
- **D.** $f(x)=-2^x-1$

## Answer and reasoning

The dashed line at $y=-1$ is the asymptote, so a 1 is subtracted outside the exponential. Check a point: $2^0-1=0$, which matches $(0,0)$, and $2^1-1=1$, which matches $(1,1)$. The other choices have the wrong asymptote or miss the points. **The function is $f(x)=2^x-1$.**

# Example 11: Graph 2




**Plotted data:** a dashed horizontal line at y = 2. A smooth curve lies below the dashed line. Moving left, it gets closer and closer to the dashed line without touching it. Moving right, it falls faster and faster. It passes through (0, 1) and (1, −1).

## Question


What is the range of the function graphed here? Use interval notation.

**A.** (−∞, 2)
**B.** (2, ∞)
**C.** (−∞, ∞)
**D.** (−∞, 1)

## Answer and reasoning

The dashed line at $y=2$ is the asymptote, and the whole curve lies below it. Every output is less than 2. Moving right, the outputs keep falling with no lowest value. The point $(0,1)$ is just one point on the curve. It is not the highest output. **The range is $(-\infty, 2)$.**

# Example 12: Graph 3




**Plotted data:** a dashed horizontal line at y = −2. A smooth curve lies above the dashed line. Moving left, it rises steeply. Moving right, it falls and gets closer and closer to the dashed line without touching it. It passes through (−1, 0) and (0, −1).

## Question

Which function has this graph?

- **A.** $f(x)=2^x-2$
- **B.** $f(x)=(1/2)^x+2$
- **C.** $f(x)=-(1/2)^x-2$
- **D.** $f(x)=(1/2)^x-2$

## Answer and reasoning

The dashed line at $y=-2$ means 2 is subtracted outside. The curve is above the line, so it is not flipped upside down. It falls as $x$ increases, so the base is between 0 and 1. Check a point: $(1/2)^{-1}-2=2-2=0$, which matches $(-1,0)$. **The function is $f(x)=(1/2)^x-2$.**