PF1 Worked Examples: Graphs of Sine and Cosine

Recording companion for the PF1 CheckIt support video — Math 172, Santa Ana College

How to use this document

- Each worked example is on its own page, in the same order as the Try It tabs in the interactive explainer, with the same numbers.
- The right-hand third of every page is intentionally blank. It is space for handwriting or ink notation while you narrate (for example, after inserting this PDF as a printout in OneNote).
- Say it at the start of the video: your CheckIt problem will have different numbers than the examples here. The reasoning is what transfers.

The one question behind every example

How tall, how long, and where does a cycle start? For a sin(bx + c) or a cos(bx + c): the amplitude is |a|, how far the graph rises above and falls below the midline. The period is 2π/b, the length of one cycle. The phase shift is c/b: factor out b to write b(x + c/b), so a plus sign shifts left and a minus sign shifts right. Start a cycle at the shift and mark every quarter period.

Adapted from TBIL Precalculus (Clontz & Lewis), CC BY-NC-SA 4.0. Drafted with Claude (Anthropic); reviewed by Dr. Marty Romero.

Example 1: Amplitude

$$f(x) = -3\sin(2x+\pi)$$

Question

What is the amplitude of f?

- **A.** -3
- **B.** 3
- **C.** 2
- **D.** 6

Answer and reasoning

The amplitude is $|a| = |-3| = 3$: the graph rises 3 above the midline and falls 3 below it. -3 keeps the sign, but a distance is never negative; the minus only flips the graph. 2 is b, which sets the period; and 6 is the full distance from the highest point to the lowest, twice the amplitude. **3.**

Adapted from TBIL Precalculus (Clontz & Lewis), CC BY-NC-SA 4.0. Drafted with Claude (Anthropic); reviewed by Dr. Marty Romero.

Example 2: Period

$$g(x)=4\cos(3x-2\pi)$$

Question

What is the period of g?

- **A.** $2\pi/3$
- **B.** 6π
- **C.** $\pi/3$
- **D.** 2π

Answer and reasoning

The period is $2\pi/b=2\pi/3$: with $b=3$, the graph fits 3 cycles where plain cosine fits 1. The shift does not change the period. 6π multiplies by 3 instead of dividing, which stretches the graph instead of squeezing it; $\pi/3$ is π/b, half a period; and 2π ignores b. ***2π/3*.**

Adapted from TBIL Precalculus (Clontz & Lewis), CC BY-NC-SA 4.0. Drafted with Claude (Anthropic); reviewed by Dr. Marty Romero.

Example 3: Both at Once

$$h(x) = -2\sin(4x + 2\pi)$$

Question

What are the amplitude and the period of h?

- **A.** Amplitude −2, period π/2
- **B.** Amplitude 2, period 8π
- **C.** Amplitude 2, period π/2
- **D.** Amplitude −2, period 8π

Answer and reasoning

Amplitude $|-2| = 2$, and period $2\pi/4 = \pi/2$. Amplitude -2 keeps the sign of a; period 8π multiplies 2π by 4; and amplitude −2 with period 8π makes both slips. **Amplitude 2, period $\pi/2$.**

Adapted from TBIL Precalculus (Clontz & Lewis), CC BY-NC-SA 4.0. Drafted with Claude (Anthropic); reviewed by Dr. Marty Romero.

Example 4: Period with b = 2

$$f(x)=5\cos(2x-2\pi/3)$$

Question

What is the period of f?

- **A.** 4π
- **B.** $\pi/2$
- **C.** π
- **D.** $2\pi/3$

Answer and reasoning

Period $=2\pi/2=\pi$. 4π multiplies by 2 instead of dividing; $\pi/2$ is π/b, half a period; and $2\pi/3$ uses $|c|$, which moves the graph but has nothing to do with the period. **π.**

Adapted from TBIL Precalculus (Clontz & Lewis), CC BY-NC-SA 4.0. Drafted with Claude (Anthropic); reviewed by Dr. Marty Romero.

Example 5: Shift 1

$$f(x)=3\sin(2x-8\pi)$$

Question

What is the phase shift of f?

A. 4π to the right
B. 8π to the right
C. 4π to the left
D. 8π to the left

Answer and reasoning

Factor out 2: $2x-8\pi=2(x-4\pi)$. So the graph is $3\sin(2x)$ moved 4π to the right. That is 4 full periods (the period is π), so the graph looks the same as $3\sin(2x)$, but CheckIt still asks for the shift itself, 4π. 8π to the right uses c without dividing by b; 4π to the left reads the minus sign the wrong way; and 8π to the left makes both slips. ***4π* to the right.**

Example 6: Shift 2

$$g(x) = -4\cos(3x + 4\pi)$$

Question

What is the phase shift of g?

A. 4π to the left
B. 4π/3 to the right
C. 4π/3 to the left
D. 4π to the right

Answer and reasoning

$3x + 4\pi = 3(x + 4\pi/3)$, so the shift is $4\pi/3$ to the left: two periods of $2\pi/3$. The -4 flips the graph but does not move it sideways. 4π to the left uses c without dividing by 3; 4π/3 to the right reads the plus sign the wrong way; and 4π to the right makes both slips. ***4π/3* to the left.**

Adapted from TBIL Precalculus (Clontz & Lewis), CC BY-NC-SA 4.0. Drafted with Claude (Anthropic); reviewed by Dr. Marty Romero.

Example 7: Shift 3

$$h(x)=2\sin(4x+8\pi/3)$$

Question

What is the phase shift of h?

A. 8π/3 to the left
B. 2π/3 to the left
C. 2π/3 to the right
D. 8π/3 to the right

Answer and reasoning

$8\pi/3 \div 4 = 8\pi/12 = 2\pi/3$, and the plus sign means left: $4(x+2\pi/3)$. The shift is longer than the period, $\pi/2$; that is allowed, and CheckIt still wants the shift itself, $2\pi/3$. 8π/3 to the left forgets to divide by 4; 2π/3 to the right reads the sign the wrong way; and 8π/3 to the right makes both slips. ***2π/3* to the left.**

Adapted from TBIL Precalculus (Clontz & Lewis), CC BY-NC-SA 4.0. Drafted with Claude (Anthropic); reviewed by Dr. Marty Romero.

Example 8: Shift 4

$$f(x)=5\sin(3x)$$

Question

What is the phase shift of f?

A. 0 (no shift)
B. 3 to the left
C. 2π/3 to the right
D. π/3 to the left

Answer and reasoning

There is no number added to $3x$, so $c=0$ and the shift is $0/3=0$: a cycle starts at $x=0$. CheckIt's key words this as "0 to the left", which means the same thing. 3 is b, which squeezes the graph instead of moving it; $2\pi/3$ is the period, the length of a cycle, not a shift; and $\pi/3$ is half a period. **0.**

Adapted from TBIL Precalculus (Clontz & Lewis), CC BY-NC-SA 4.0. Drafted with Claude (Anthropic); reviewed by Dr. Marty Romero.

Example 9: Graph 1




Figure: A wave on a grid with x from 0 to 4π/3, marked every π/6 and labeled at π/3, 2π/3, π and 4π/3, and y from −3 to 3. It starts at the origin going down, reaches −2 at x = π/6, crosses back up through 0 at π/3, reaches 2 at π/2, returns to 0 at 2π/3, and repeats once more, ending at 4π/3.

Question

Which function does this graph show?

- **A.** $y=2\sin(3x)$
- **B.** $y=-2\sin(2x)$
- **C.** $y=2\sin(2x)$
- **D.** $y=-2\sin(3x)$

Answer and reasoning

The graph goes from 2 to -2, so the amplitude is 2. It starts on the midline going down, which is sine flipped, so $a=-2$. One cycle runs from 0 to $2\pi/3$, so $2\pi/b=2\pi/3$ and $b=3$. $2\sin(3x)$ would start going up; $-2\sin(2x)$ has period π, so it would fit only one and a third cycles here; and $2\sin(2x)$ makes both slips. ***y=−2 sin(3x).***

Example 10: Graph 2




Figure: A wave on a grid with x from −π/2 to 3π/2, marked every π/4 and labeled at −π/2, π/2, π and 3π/2, and y from −5 to 5. It is at 4 at x = −π/2, falls to −4 at x = 0, rises to 4 at π/2, falls to −4 at π, and rises to 4 at 3π/2.

$$f(x)=4\cos(2x+\pi)$$

Question

This is the graph CheckIt would draw for f. What amplitude and period does it show?

- **A.** Amplitude 4, period π
- **B.** Amplitude 8, period π
- **C.** Amplitude 4, period π/2
- **D.** Amplitude 8, period π/2

Answer and reasoning

The graph goes up to 4 and down to -4: the amplitude is half that 8-unit spread, 4, which matches $|a|$. From one peak at $-\pi/2$ to the next at $\pi/2$ is π, matching $2\pi/2$. Amplitude 8 is the full distance from peak to valley; period $\pi/2$ measures from a peak to the next valley, half a cycle; and amplitude 8 with period $\pi/2$ makes both slips. **Amplitude 4, period π.**

Example 11: Where a Cycle Starts

$$f(x)=3\sin(2x-\pi)$$

Question

Over which interval does one full cycle of f run, starting at the phase shift?

- **A.** From π/2 to 3π/2
- **B.** From π to 3π
- **C.** From −π/2 to π/2
- **D.** From π/2 to 5π/2

Answer and reasoning

$2x-\pi=2(x-\pi/2)$: the shift is $\pi/2$ to the right. The period is $2\pi/2=\pi$. So a cycle starts at $\pi/2$ and ends one period later, at $\pi/2+\pi=3\pi/2$. From π to 3π makes two slips: it shifts by $c=\pi$ without dividing by 2, and uses a period of 2π; from $-\pi/2$ shifts left instead of right; and ending at $5\pi/2$ uses a period of 2π, ignoring b. **From $\pi/2$ to $3\pi/2$.**

Adapted from TBIL Precalculus (Clontz & Lewis), CC BY-NC-SA 4.0. Drafted with Claude (Anthropic); reviewed by Dr. Marty Romero.

Example 12: Graph 3




Figure: A wave on a grid with x from 0 to 4π/3, marked every π/12 and labeled every π/4, and y from −3 to 3. It is at −1 when x = 0, falls to −2 at π/12, rises to a peak of 2 at π/3, falls to −2 at 7π/12, rises to 2 at 5π/6, falls to −2 at 13π/12, and rises to 2 at 4π/3.

Question

The first peak to the right of 0 is at (π/3, 2), and the graph repeats every π/2. Which function does it show?

- **A.** $y=2\cos(4x+4\pi/3)$
- **B.** $y=2\cos(3x-\pi)$
- **C.** $y=2\cos(4x-4\pi/3)$
- **D.** $y=2\sin(4x-4\pi/3)$

Answer and reasoning

The period is $\pi/2$, so $b=4$. Cosine starts a cycle at a peak, so shift a cosine to put its peak at $\pi/3$: $2\cos(4(x-\pi/3))=2\cos(4x-4\pi/3)$. Check: at $x=\pi/3$ the inside is 0, and $2\cos 0=2$. $4x+4\pi/3$ shifts left instead, so its first peak is at $\pi/6$; $2\cos(3x-\pi)$ also peaks at $\pi/3$, but its period is $2\pi/3$, not $\pi/2$; and the sine version is 0 at $\pi/3$, with its first peak at $11\pi/24$. **$y=2\cos(4x-4\pi/3)$.**

Example 13: Slip 1

$$f(x)=2\sin(3x-9\pi/2)$$

phase shift: 9π/2 to the right

Question

A student found the phase shift this way. What is the mistake, if any?

A. There is no mistake.
B. Divide by b: 3π/2 to the right.
C. A minus sign means left: 9π/2 to the left.
D. The shift is the period, 2π/3 to the right.

Answer and reasoning

Factor out 3: $3x-9\pi/2=3(x-3\pi/2)$. The shift is $c/b=9\pi/2\div3=3\pi/2$ to the right; the student used c itself. A minus sign means right, so the direction was correct; and the period, $2\pi/3$, is the length of a cycle, not where it starts. ***3π/2* to the right.**

Adapted from TBIL Precalculus (Clontz & Lewis), CC BY-NC-SA 4.0. Drafted with Claude (Anthropic); reviewed by Dr. Marty Romero.

Example 14: Slip 2

$$f(x)=4\cos(2x+2\pi)$$
$$\text{period}=2\pi\cdot 2=4\pi$$
$$\text{phase shift: } \pi \text{ to the left}$$

Question

A student found the period and the phase shift. What is the mistake, if any?

A. There is no mistake.
B. The shift should be 2π to the left, not π.
C. Divide, not multiply: the period is π.
D. The period is 2π, since cosine repeats every 2π.

Answer and reasoning

The period is $2\pi/b=2\pi/2=\pi$. Multiplying by b would stretch the graph, but a larger b squeezes it. The shift is correct: $2x+2\pi=2(x+\pi)$, π to the left. 2π to the left forgets to divide c by 2; and plain cosine repeats every 2π, but this one is squeezed by $b=2$. **The period is π.**

Adapted from TBIL Precalculus (Clontz & Lewis), CC BY-NC-SA 4.0. Drafted with Claude (Anthropic); reviewed by Dr. Marty Romero.

Example 15: Slip 3

$$f(x)=-5\sin(4x+4\pi/3)$$

amplitude 5, period π/2, shift π/3 left

Question

A student found all three features. What is the mistake, if any?

A. There is no mistake.
B. The amplitude should be −5, since a = −5.
C. The shift should be 4π/3 to the left.
D. The period should be 8π, 2π times 4.

Answer and reasoning

Amplitude $|-5|=5$; period $2\pi/4=\pi/2$; and $4x+4\pi/3=4(x+\pi/3)$, so the shift is $\pi/3$ to the left. The student is right. An amplitude is a distance, so it is positive even when a is negative; $4\pi/3$ is c before dividing by 4; and the period divides by b. **There is no mistake.**

Adapted from TBIL Precalculus (Clontz & Lewis), CC BY-NC-SA 4.0. Drafted with Claude (Anthropic); reviewed by Dr. Marty Romero.

Example 16: Slip 4

$$f(x)=3\cos(2x+2\pi/3)$$

phase shift: π/3 to the right

Question

A student found the phase shift this way. What is the mistake, if any?

A. There is no mistake.
B. Do not divide by b: the shift is 2π/3 to the right.
C. The shift is the period: π to the right.
D. A plus sign means left: π/3 to the left.

Answer and reasoning

$2x+2\pi/3=2(x+\pi/3)=2(x-(-\pi/3))$: the graph moves $\pi/3$ to the left. The size was right, but the direction was backward. A check: the peak of cosine, where the inside is 0, is at $x=-\pi/3$, left of 0. The shift is c/b, so dividing is right; and the period, π, is not the shift. **$\pi/3$ to the left.**

Example 17: Slip 5

$$f(x) = -4\sin(3x - 2\pi)$$

amplitude −4, period 2π/3, shift 2π/3 right

Question

A student found all three features. What is the mistake, if any?

A. There is no mistake.
B. The amplitude is a distance: it is 4.
C. The shift should be 2π to the right.
D. The period should be 6π, 2π times 3.

Answer and reasoning

The amplitude is $|a| = |-4| = 4$; the minus flips the graph upside down but does not make the height negative. The period, $2\pi/3$, and the shift, $2\pi/3$ to the right from $3(x - 2\pi/3)$, are both correct. Here the shift happens to equal the period, so the graph lines up with an unshifted one. 2π to the right forgets to divide by 3; and the period divides by b. **Amplitude 4.**