

TABLE 3 Incidence of the nine most frequent adverse drug reactions (ADRs) for each antiepileptic drug (AED), expressed as absolute number of trials with ADR, and as percentage of all trials

AED	Patients with ADR (%)	Weight change	Lethargy	Tremor	Cognitive impairment	Behavioral disorder	Depression	Gastrointestinal ADRs	Adverse cutaneous reaction	Speech disorder
VPA	93 (37.5)	58 (20.8)	19 (6.8)	30 (10.8)	12 (4.3)	6 (2.2)	2 (0.7)	3 (1.1)	0 (0.0)	1 (0.4)
LTG	25 (16.3)	2 (1.2)	4 (2.5)	4 (2.5)	3 (1.9)	0 (0.0)	2 (1.2)	2 (1.2)	4 (2.5)	0 (0.0)
LEV	30 (24.6)	0 (0.0)	11 (8.9)	1 (0.8)	3 (2.4)	16 (12.9)	4 (3.2)	0 (0.0)	0 (0.0)	1 (0.8)
CBZ	9 (14.5)	0 (0.0)	1 (1.6)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	4 (6.2)	4 (6.2)	0 (0.0)
TPM	25 (45.5)	4 (6.7)	5 (8.3)	2 (3.3)	10 (16.7)	3 (5.0)	3 (5.0)	0 (0.0)	0 (0.0)	3 (5.0)