

Table 1 Agents administered to the mouse. The table summarizes all agents injected including company, injection site, volume, concentration, vehicle, administration time, sacrifice time, and number of retinas used

Agent	Company	Injection route	Volume	Concentration (Vehicle)	Time of administration	Time of sacrifice	n of mice
Tissue plasminogen activator (tPA)	Actilyse; INN alteplase, Boehringer Ingelheim, Germany	Tail vein	50 μ l/min (1 min) + 15 μ l/min (30–60 min)	10 mg/kg	1 h after ischemia	30–60 min after tPA injection	17
Dextran (FITC–70S or/and TRITC–500S)	Sigma	Tail vein	0.2 ml	5 mg/ml (saline)	1 h before ischemia and 1 h after tPA injection	1 h after ischemia/recanalisation	34
Adenosine	Molecular Probes	Intravitreal	3 μ l	1 mg/ml (water)	Just before ischemia	1 h after ischemia	4
					1 h after ischemia	2 h after ischemia	3
Amlodipine	Pfizer	Intravitreal	3 μ l	1 mg/ml (water)	1 h before ischemia	1 h after ischemia	3
Carbenoxolone (CBX)	Sigma	Intravitreal	3 μ l	30 μ M (water)	Just before ischemia	1 h after ischemia	4
Fluorescent phalloidin	Biotium	Intravitreal	3 μ l	200 U/ml (water)	1–2 h before ischemia	1 h after ischemia	5
Propidium iodide (PI)	Molecular Probes, Invitrogen	Intravitreal	1 μ l	1 mg/ml (water)	1 h after ischemia	4–6 h after ischemia	5
Tissue plasminogen activator (tPA) + Fluorescent phalloidin	Actilyse; INN alteplase, Boehringer Ingelheim, Germany; Biotium	Tail vein Intravitreal	50 μ l/min (1 min) + 15 μ l/min (30–60 min) 3 μ l	10 mg/kg 200 U/ml (water)	1 h after ischemia 30 min after ischemia	1 h after tPA injection	6
Fluo-4	Molecular Probes, Invitrogen	Intravitreal	3 μ l	5 μ M (0.005% pluronic acid, 0.00025% cremophor and 0.05% DMSO in artificial CSF)	Just before ischemia	2 h after ischemia	18
Scrambled peptide	Smart-Bioscience	Intravitreal	2 μ l	20 μ M (nuclease free water)	1 h before ischemia	1 h after ischemia	6
Peptide	Smart-Bioscience	Intravitreal	2 μ l	20 μ M (nuclease free water)	1 h before ischemia	1 h after ischemia	6
1,4-Dideoxy-1,4-imino-D-arabinitol hydrochloride (DAB)	Sigma-Aldrich	Intravitreal	3 μ l	2.5 M (saline)	2 h before ischemia	30 min after ischemia	9
Scrambled siRNA	Ambion® LifeTech,	Intravitreal	3 μ l	0.5 mg/ml (10% Glucose, nuclease free water, in-vivo-jetPEI®, 10% Glucose)	48 h before ischemia	1 h after ischemia	3
α SMA siRNA	Ambion® LifeTech,	Intravitreal	3 μ l	0.5 mg/ml (10% Glucose, nuclease free water, in-vivo-jetPEI®, 10% Glucose)	48 h before ischemia	1 h after ischemia	3