

Table 2. Evaluation of the SPRINT outcomes within each recommendation group for validating the treatment recommendation^a

	SPRINT randomization	Total SPRINT population	Intensive treatment recommendation group	Non-intensive treatment recommendation group
Number of participants, <i>n</i>	Actual intensive	4,677	2,916	1761
	Actual non-intensive	4,683	2,899	1784
Mean systolic BP ^b , mmHg	Actual intensive	121.01	120.68	121.55
	Actual non-intensive	135.28	134.99	135.74
Primary composite cardiovascular outcome, per year	Actual intensive	1.65	1.78	1.45
	Actual non-intensive	2.19	2.60	1.55
	Ratio (95% CI)	0.75 (0.64–0.89)	0.68 (0.56–0.84)	0.93 (0.69–1.26)
	<i>P</i> -value	0.00092	0.00022	0.64769
All-cause mortality, per year	Actual intensive	1.03	1.01	1.05
	Actual non-intensive	1.40	1.56	1.14
	Ratio (95% CI)	0.74 (0.60–0.91)	0.65 (0.50–0.84)	0.92 (0.65–1.31)
	<i>P</i> -value	0.00355	0.00105	0.64574
Serious adverse event composite outcome, per year	Actual intensive	2.89	2.59	3.38
	Actual non-intensive	2.03	1.84	2.32
	Ratio (95% CI)	1.42 (1.22–1.65)	1.41 (1.15–1.72)	1.45 (1.16–1.82)
	<i>P</i> -value	< 0.00001	0.00092	0.00123

SPRINT systolic blood pressure intervention trial, BP blood pressure, CI confidence interval

^aValues of outcomes are presented as yearly incidence^bUsing the last three measurements of each participant during follow-up