

Table 1: Characteristics of the subjects by machine-measured nicotine yield of cigarette

Characteristics	Machine-measured nicotine yield (mg/cigarette)			Total	p
	0.1	0.2–0.8	0.9–2.4		
Number of subjects	87	223	148	458	
Mean and SD of machine-measured nicotine yield (mg/cigarette)	0.1	0.5 ± 0.2	1.1 ± 0.4	0.6 ± 0.4	
Mean and SD of age	53.6 ± 10.4	50.6 ± 9.34	50.8 ± 10.4	51.2 ± 9.9	0.07
Mean and SD of number of cigarettes per day	23.4 ± 12.2	24.5 ± 10.7	24.4 ± 9.5	24.4 ± 10.6	0.93
Mean and SD of FTND	5.1 ± 2.5	5.4 ± 2.3	5.6 ± 2.0	5.4 ± 2.2	0.21
Number of smokers at each category of FTND					
FTND score 0–3	19 (21.8%)	47 (21.1%)	21 (14.2%)	87 (19.0%)	0.41
FTND score 4–6	40 (46.0%)	100 (44.8%)	78 (52.7%)	218 (47.6%)	
FTND score 7–10	28 (32.2%)	76 (34.1%)	49 (33.1%)	153 (33.4%)	
Number of smokers having attempted to quit	50 (61.0%)	132 (59.5%)	84 (56.8%)	269 (58.9%)	0.73
Number of smokers at each stage of behavioral change in quitting process					
Precontemplation-1	16 (18.6%)	47 (21.3%)	58 (39.7%)	121 (26.7%)	0.001
Precontemplation-2	48 (55.8%)	131 (59.3%)	62 (42.5%)	241 (53.2%)	
Contemplation	20 (23.3%)	33 (14.9%)	23 (15.6%)	76 (16.8%)	
Preparation	2 (2.3%)	10 (4.5%)	3 (2.1%)	15 (3.3%)	
Mean and SD of urinary cotinine concentration (mean-SD, mean+SD*)	535 (1782,160)	770 (1981,299)	1010 (2071,492)	784 (484, 1264)	<0.001

FTND:Fagerström Test for Nicotine Dependence P for difference is examined by analysis of variance or chi-square test * Back-transformation of log-transformed data