

TABLE 2. SUMMARY OF SURGICAL PROCEDURES, ANAPLASTIC THYROID CARCINOMA VIABILITY, RESECTED TUMOR HISTOPATHOLOGY, AND PRETREATMENT MUTATION PROFILE

	Patient 1	Patient 2	Patient 3	Patient 4	Patient 5	Patient 6
Procedures	TT, central ND, bilateral ND, limited resection of esophageal muscularis	TT, central ND, right ND, <i>en bloc</i> resection of right RLN	TT, central ND, left ND, <i>en bloc</i> resection of left RLN, shave resection of trachea	TT, central ND, revision right ND, left ND, resection of dermal metastasis	TT, central ND, left level IV ND, and resection of occipital scalp lesion	TT, central ND
Resection of ATC	Complete	Complete	Complete	Complete	Complete	Complete
Residual (R) tumor classification	R1	R0	R1	R0	R0	R0
Resected ATC viability, % ^a	50	5	<5	0	0	0
Additional component in resected tumor	PTC	PTC	PTC	PDTC, PTC	PTC	PTC
Pretreatment tumor mutations	<i>BRAF</i> ^{V600E} , <i>TP53</i> ^{R175H} , <i>EGFR</i> ^{G322S} , <i>BRAF</i> amplification ^b	<i>BRAF</i> ^{V600E} , <i>TP53</i> ^{Q331A}	<i>BRAF</i> ^{V600E} , <i>TP53</i> ^{D208V} , <i>CDKN2A</i> ^{pE88a}	<i>BRAF</i> ^{V600E}	<i>BRAF</i> ^{V600E} , <i>ATM</i> ^{T1986V} , <i>MCL1</i> duplication	<i>BRAF</i> ^{V600E}

^aIndicated viability of the ATC component of the postneoadjuvant treatment resected primary thyroid tumor.

^bDetected in cell-free DNA.

ATC, anaplastic thyroid carcinoma; ND, neck dissection; PDTC, poorly differentiated thyroid carcinoma; PTC, papillary thyroid carcinoma; RLN, recurrent laryngeal nerve; TT, total thyroidectomy.