

Association between the manganese superoxide dismutase polymorphism and breast cancer risk in the Shanghai Breast Cancer Study

Genotype	Cases (n [%])	Controls (n [%])	OR ^a	95% CI
All subjects				
Val/Val	831 (73.9)	884 (73.9)	1.0	Reference
Val/Ala	266 (23.6)	290 (24.2)	1.0	0.8–1.2
Ala/Ala	28 (2.5)	23 (1.9)	1.3	0.7–2.3
Val/Val and Val/Ala	1097 (97.5)	1174 (98.1)	1.0	Reference
Ala/Ala	28 (2.5)	23 (1.9)	1.3	0.7–2.3
Stratified analyses by menopausal status				
Premenopausal women				
Val/Val and Val/Ala	729 (97.2)	751 (98.4)	1.0	Reference
Ala/Ala	21 (2.8)	12 (1.6)	1.8	0.9–3.7
Postmenopausal women				
Val/Val and Val/Ala	363 (98.1)	419 (97.4)	1.0	Reference
Ala/Ala	7 (1.9)	11 (2.6)	0.8	0.3–2.0
<i>P</i> for interaction = 0.147				
Stratified analyses by age				
Age < 45 years				
Val/Val and Val/Ala	442 (97.1)	488 (98.4)	1.0	Reference
Ala/Ala	13 (2.9)	8 (1.6)	1.8	0.7–4.3
Age ≥ 45 years				
Val/Val and Val/Ala	655 (97.8)	686 (97.9)	1.0	Reference
Ala/Ala	15 (2.2)	15 (2.1)	1.1	0.5–2.2
<i>P</i> for interaction = 0.380				

^aAdjusted for age. CI, confidence interval; OR, odds ratio.

controls in a Finnish Caucasian population and reported that women or had higher BMI also had an increased risk for