

Birth order distributions and odds ratios for breast cancer among white and African-American women by race

	Minimally adjusted OR ^a								Fully adjusted OR ^b							
	African-American				White				African-American				White			
	Case	Control	OR	95% CI	Case	Control	OR	95% CI	Case	Control	OR	95% CI	Case	Control	OR	95% CI
Full CBCS ^c	<i>n</i> = 331	<i>n</i> = 329			<i>n</i> = 523	<i>n</i> = 456			<i>n</i> = 331	<i>n</i> = 329			<i>n</i> = 523	<i>n</i> = 456		
1st born (ref.)	106	103	Ref.		191	165	Ref.		106	103	Ref.		191	165	Ref.	
2nd–4th born	142	151	0.8	0.6–1.2	264	226	1.0	0.7–1.3	142	151	0.8	0.6–1.2	264	226	1.0	0.7–1.3
≥5th born	83	75	1.0	0.7–1.5	68	65	1.0	0.7–1.5	83	75	1.0	0.7–1.5	68	65	1.0	0.7–1.5
Mean ± SD	3.6 ± 2.7	3.3 ± 2.7			2.6 ± 1.9	2.6 ± 2.0										
Median	2.0	3.3			2.0	2.0										
Range	1–14	1–14			1–12	1–11										
Born ≥1948 ^{d,e}	<i>n</i> = 129	<i>n</i> = 134			<i>n</i> = 233	<i>n</i> = 181			<i>n</i> = 101	<i>n</i> = 109			<i>n</i> = 164	<i>n</i> = 115		
1st born (ref.)	37	41	Ref.		80	67	Ref.		29	31	Ref.		55	45	Ref.	
2nd–4th born	56	61	1.0	0.5–1.7	138	97	1.2	0.8–1.9	46	50	0.7	0.4–1.5	98	59	1.0	0.6–1.8
≥5th born	36	32	1.2	0.6–2.3	15	17	0.7	0.3–1.6	26	28	0.7	0.3–1.6	11	11	0.5	0.2–1.5
Mean ± SD	3.6 ± 2.7)	3.3 ± 2.6)			2.3 ± 1.5)	2.3 ± 1.6)										
Median	3.0	2.5			2.0	2.0										
Range	1–14	1–14			1–10	1–11										

¹Adjusted for age and sampling fractions. ²Adjusted for age, sampling fractions, and other covariates as specified. ³Additional covariates for fully adjusted OR: none. ⁴Additional covariates for fully adjusted OR: adult body mass index >25 kg/m², household income ≥\$30,000 and maternal age. ⁵Restricted to participants born on or after 1 January 1948. CBCS, Carolina Breast Cancer Study; CI, confidence interval; OR, odds ratio.