

# Association of CHEK2 variants with breast cancer risk

| CHEK2 status <sup>a</sup> | Cases (n = 506) |      | Controls (n = 459) |       | OR <sup>b</sup> | 95% CI     |
|---------------------------|-----------------|------|--------------------|-------|-----------------|------------|
|                           | n               | %    | n                  | %     |                 |            |
| 1100del C                 |                 |      |                    |       |                 |            |
| Noncarrier                | 499             | 98.8 | 456                | 99.6  | 1.0             | Reference  |
| Carrier                   | 6               | 1.2  | 2                  | 0.4   | 2.9             | (0.6–14.6) |
| R145W                     |                 |      |                    |       |                 |            |
| Noncarrier                | 505             | 99.8 | 459                | 100.0 | 1.0             | Reference  |
| Carrier                   | 1               | 0.2  | 0                  | -     | -               | -          |
| I175T                     |                 |      |                    |       |                 |            |
| Noncarrier                | 504             | 99.6 | 455                | 99.1  | 1.0             | Reference  |
| Carrier                   | 2               | 0.4  | 4                  | 0.9   | 0.5             | (0.1–2.6)  |

<sup>a</sup>All carriers are heterozygous. <sup>b</sup>Adjusted for age at reference and reference year. CI, confidence interval; OR, odds ratio.