

**Odds ratios for breast cancer and total PCBs in relation to CYP1A1 M1 genotypes in African American and white women**

| Total PCBs <sup>a</sup> | CYP1A1 M1 genotype | All participants      |                          | Premenopausal         |                          | Postmenopausal        |                          |
|-------------------------|--------------------|-----------------------|--------------------------|-----------------------|--------------------------|-----------------------|--------------------------|
|                         |                    | Patients/<br>controls | OR (95% CI) <sup>b</sup> | Patients/<br>controls | OR (95% CI) <sup>b</sup> | Patients/<br>controls | OR (95% CI) <sup>b</sup> |
| African Americans       |                    |                       |                          |                       |                          |                       |                          |
| <0.430                  | Non-M1             | 66/75                 | Referent                 | 46/51                 | Referent                 | 20/24                 | Referent                 |
| ≥0.430                  | Non-M1             | 75/67                 | 1.5 (0.9–2.5)            | 21/16                 | 1.9 (0.9–4.2)            | 54/51                 | 1.0 (0.5–2.3)            |
| <0.430                  | Any M1             | 42/46                 | 1.0 (0.6–1.7)            | 35/33                 | 1.2 (0.6–2.2)            | 7/13                  | 0.6 (0.2–1.8)            |
| ≥0.430                  | Any M1             | 59/54                 | 1.4 (0.8–2.5)            | 17/14                 | 1.7 (0.8–4.1)            | 42/40                 | 1.0 (0.5–2.3)            |
| ICR (95% CI)            |                    |                       | 0.0 (-0.9,0.9)           |                       | -0.3 (-2.2,1.6)          |                       | 0.4 (-0.5,1.3)           |
| Whites                  |                    |                       |                          |                       |                          |                       |                          |
| <0.349                  | Non-M1             | 174/133               | Referent                 | 118/83                | Referent                 | 56/50                 | Referent                 |
| ≥0.349                  | Non-M1             | 122/148               | 0.7 (0.5–1.0)            | 38/43                 | 0.6 (0.4–1.1)            | 84/105                | 0.8 (0.5–1.3)            |
| <0.349                  | Any M1             | 45/44                 | 0.8 (0.5–1.2)            | 29/31                 | 0.7 (0.4–1.2)            | 16/13                 | 1.1 (0.5–2.5)            |
| ≥0.349                  | Any M1             | 29/32                 | 0.8 (0.4–1.4)            | 11/10                 | 0.7 (0.3–1.9)            | 18/22                 | 0.8 (0.4–1.7)            |
| ICR (95% CI)            |                    |                       | 0.4 (-0.2,0.9)           |                       | 0.5 (-0.3,1.2)           |                       | 0.0 (-1.1,1.0)           |

<sup>a</sup>Lipid-adjusted total imputed PCBs (ng/ml).

<sup>b</sup>Odds ratios adjusted for age and sampling fractions.

Any M1, M1/M1 or M1/non-M1; CI, confidence interval; ICR, interaction contrast ratio; non-M1, non-M1/non-M1.