




Recommendations for broad-scale diversity models from an independent study

Martin Jung
Center for Macroecology, Climate and Evolution

BIODIVERSITY MODELS



Predicted change in species richness from Newbold et al. (2015)

POTENTIAL PROBLEMS

POTENTIAL PROBLEMS

- Many broad-scale models generalize over a wide range of variables

POTENTIAL PROBLEMS

- Often only single factor (land use) models

POTENTIAL PROBLEMS


- Based on available (biased ?) data


QUESTION :

QUESTION :

How well does an African-wide model reflect species diversity and abundance on a local land-use gradient where local conditions are known?

METHODS





PV

SV

PL

CL

UR

- Timed point counts

- Timed point counts
- Total 147 sites: 172 species, 2700 individual counts

AUXILIARY DATA FOR THE COMPARISON


AUXILIARY DATA FOR THE COMPARISON

- Remote-sensing and census data
(INDVI and meanNDVI, Human population density, Forest-Cover year 2000)


AUXILIARY DATA FOR THE COMPARISON


- Remote-sensing and census data
(INDVI and meanNDVI, Human population density, Forest-Cover year 2000)
- Functional traits
(Range size, Threat status, Forest specialization)

RESULTS



Transect ● Kilimanjaro ▲ PREDICTS ■ Taita





A wide-angle photograph of a tropical landscape. In the foreground, there is a dense growth of tall, green grasses and some shorter leafy plants. A small, irregular clearing or path is visible in the center-left. The middle ground is dominated by more dense vegetation. In the background, a thick forest of various trees and palms stretches across the horizon under a clear sky.

SUMMARY

SUMMARY

- Cropland higher number of species than the average more intense cropland site (Agroforestry)
- More forest-specialists in cropland
- Primary forest likely lower diversity due to size and fragmentation

CONCLUSION

CONCLUSION

- Large-scale models succeed at detecting overall impacts of land use change.

CONCLUSION

- Large-scale models succeed at detecting overall impacts of land use change.
- However they might lose accuracy if they are used to predict local impacts on biodiversity, if local conditions do not conform

CONCLUSION

- Large-scale models succeed at detecting overall impacts of land use change.
- However they might lose accuracy if they are used to predict local impacts on biodiversity, if local conditions do not conform
- Policy makers need to look not only at large-scale patterns, but also at local differences, when deciding on conservation actions.

ACKNOWLEDGEMENTS

- Tim Newbold and Neil Burgess for scientific advice
- PREDICTS contributors for making data available
(<http://predicts.org.uk>)
- The CHIESA project partners for support and coordination
- Everyone at CMEC and UNEP-WCMC
- Danida Fellowship Centre for financial support



Danida Fellowship Centre
- sustaining development through research and learning

**THANKS FOR
LISTENING!**

conservationecology.wordpress.com