




Recommendations for broad-scale diversity models from an independent study

Martin Jung
Center for Macroecology, Climate and Evolution

BIODIVERSITY MODELS



Predicted change in species richness from Newbold et al. (2015)

POTENTIAL PROBLEMS

POTENTIAL PROBLEMS

- Many broad-scale models generalize over a wide range of variables in a single factor (land use)

POTENTIAL PROBLEMS

- It is tempting to apply those models in a new spatial context

POTENTIAL PROBLEMS


- Based on available (biased ?) data


QUESTION :

QUESTION :

How well does a broad-scale model reflect species diversity and abundance on a local land-use gradient where local conditions are known?

METHODS





PV

SV

PL

CL

UR

- Timed point counts

- Timed point counts
- Total 147 sites: 172 species, 2700 individual counts

AUXILIARY DATA FOR THE COMPARISON


AUXILIARY DATA FOR THE COMPARISON

- Remote-sensing and census data
(INDVI and meanNDVI, Human population density, Forest-Cover year 2000)


AUXILIARY DATA FOR THE COMPARISON


- Remote-sensing and census data
(INDVI and meanNDVI, Human population density, Forest-Cover year 2000)
- Functional traits
(Range size, Threat status, Forest specialization)

RESULTS



Transect ● Kilimanjaro ▲ PREDICTS ■ Taita







SUMMARY

SUMMARY

- Cropland higher number of species than the average more intense cropland site (Agroforestry)

SUMMARY

- Cropland higher number of species than the average more intense cropland site (Agroforestry)
- More forest-specialists in cropland

SUMMARY

- Cropland higher number of species than the average more intense cropland site (Agroforestry)
- More forest-specialists in cropland
- Community differences in plantation forest maybe driven by type (Eucalyptus)

SUMMARY

- Cropland higher number of species than the average more intense cropland site (Agroforestry)
- More forest-specialists in cropland
- Community differences in plantation forest maybe driven by type (Eucalyptus)
- However: Primary forest likely lower diversity due to size and fragmentation

CONCLUSION

CONCLUSION

- Large-scale models succeed at detecting overall impacts of land use change.

CONCLUSION

- Large-scale models succeed at detecting overall impacts of land use change.
- However they might lose accuracy if they are used to predict local impacts on biodiversity, if local conditions do not conform

CONCLUSION

- Large-scale models succeed at detecting overall impacts of land use change.
- However they might lose accuracy if they are used to predict local impacts on biodiversity, if local conditions do not conform
- Species traits and environmental co-variates could provide a window of opportunity to improve predictions in regions, where coarse categories don't capture all of the variability.

ACKNOWLEDGEMENTS

- Tim Newbold and Neil Burgess for scientific advice
- PREDICTS contributors for making data available
(<http://predicts.org.uk>)
- The CHIESA project partners for support and coordination
- Everyone at CMEC and UNEP-WCMC
- Danida Fellowship Centre for financial support



Danida Fellowship Centre
- sustaining development through research and learning

**THANKS FOR
LISTENING!**