

Certified Algorithms for Combinatorial Optimization Problems

Martin Sokolov

May 3, 2025

Contents

1	Introduction	3
2	Preliminaries	5
2.1	Graph Theory	5
2.2	Combinatorial Optimization	5
2.2.1	NP-Complete Problems: Definitions and Variants	5
2.2.2	Approximation Algorithms and Complexity Classes	7
2.2.3	Baker's approach	7
2.2.4	Treewidth	9
2.2.5	Linear Programming and Randomized Rounding	9
2.3	Beyond the Worst-Case Analysis of Algorithms	10
2.4	r-Dominating Set; Exact algorithm and PTAS	10
3	(E)PTAS for combinatorial optimization problems using Baker's technique	12
3.1	Baker's technique for unweighted FVS-4CL on planar graphs	12
3.2	Baker's technique for the weighted FVS-4CL on planar graphs	17
3.3	Baker's technique for weighted FVS-BCL of arbitrary cycle length	20
4	Exact algorithm for FVS-4CL using MSOL	25
4.1	MSOL - a language for graph problems	25
4.2	Exact algorithm for FVS-4CL using MSOL	25
4.3	Exact algorithm for weighted FVS-BCL using MSOL	26
4.4	Exact algorithm for FVS-4CL using dynamic programming	26
5	m-stitching and Π-m-stitching	29
5.1	FVS-4CL and 2-stitching	29
5.2	FVS-BCL and $(\rho/2)$ -stitching	35
5.3	r-Dominating Set and $(r+1)$ -stitching	35
6	Certified Algorithm	38

6.1	FVS-4CL	38
6.2	FVS-BCL	38
6.3	r-Dominating Set	38

1 Introduction

The FEEDBACK VERTEX SET (FVS) problem is central to the field of computer science. Karp’s paper [28] showed that there is a reduction from SATISFIABILITY to twenty-one problems, one of which is FVS. Because of the hardness of the problem, extensive research has been conducted in order to find a good approximation solution. The problem is **APX-complete** and the best approximation ratio so far is two and was first found by Becker and Geiger [8]. Around the same time, Bafna et al. [4] devised the first local ratio technique in order to arrive at the same approximation ratio. These algorithms were translated in primal-dual terms by Chudak et al. [18] who constructed a third algorithm which yields a 2-approximation both for the unweighted and weighted case of the FVS problem.

For planar graphs and H-MINOR-FREE graphs two approaches have yielded PTASs for many problems. Baker’s technique [5] focused on approximation algorithms for **NP-Complete** problems on planar graphs, while Demaine and Hajiaghayi [23] generalized the shifting technique for the latter class of graphs. One of the known limitations of the shifting technique is that it does not handle non-local problems, such as the ones we are examining (FVS, CYCLE PACKING). In our paper, we impose restrictions on the problems in order to introduce a local property that allows us to devise a PTAS for the FVS-BCL problem. Baker’s technique allows for the translation into the weighted version of the problem. We adapt the algorithm to work for the MWFVS-BCL all the while trying to follow the framework for devising certified algorithms established in [31, 1, 16].

The FVS problem has applications in the field of Bayesian networks. As we said, researchers turned to approximation algorithms and made use of the linear programming (LP) framework in order to get desirable ratios. In this paper, we make use of the existing literature on LP for FVS and analyse how to translate the results for the class of graphs with bounded cycle length. In Section 2.3, we familiarize the reader with two ways of designing certified algorithms. One way is by using convex relations. This motivates us to define the problem as an ILP, solve the LP relaxation and exploit a randomized rounding scheme to get the values for the approximate and co-approximate conditions, as defined in the same section. Those values are then used to calculate the γ for the γ -certified algorithm.

Finally, we consider the notions of Bilu-Linial stability [9] and certified algorithms [31]. In the last two paragraphs, we explained how we plan to translate existing techniques to get certified solutions of instances. In the final work, the last sections will be dedicated to proving that these algorithms are indeed certified.

Related Work

This work is inspired by Baker’s seminal paper on designing PTASs in planar graphs [5]. There have been other techniques that generalize this approach, namely Demaine and Hajiaghayi [23]’s bidimensionality theory. The latter framework explores problems in H-MINOR-FREE graphs.

The notion of Bilu-Linial stability has given rise to much exploration of structured instances in many problems. In their paper, Bilu and Linial [9] offer the first evidence that stable instances are much easier than worst-case instances of problems. They propose an exact algorithm for $O(n)$ -stable instances of MAX CUT. This result was improved by Bilu et al. [10] who designed a $O(\sqrt{n})$ for the same problem. Further research yielded a general approach to analysing stable instances in Makarychev et al. [32]. They applied that approach to graph partitioning problems and designed $O(\sqrt{\log n \log \log n})$ -stable instances of MAX CUT and 4-stable instances of MINIMUM MULTIWAY CUT. The study of perturbation resilience of clustering problems is explored in [3, 6, 7]. On the TRAVELLING SALESMAN PROBLEM, Mihalák et al.

[34] gave an algorithm for 1.8-stable instances. We refer the reader to the survey [30].

On the subject of Beyond the Worst-Case Analysis, we refer to Makarychev and Makarychev [31]. In that paper certified algorithms are first described in an attempt to bridge the gap between the worst-case and structured instances. Since then, researchers have made use of the definitions and results in that paper and applied them to other problems. Angelidakis et al. [1] explore the notion of stability and initiate the study of certified algorithms for the MAXIMUM INDEPENDENT SET problem (MIS). They provide robust and certified algorithms for $(1+\epsilon)$ -stable instances of planar MIS, for any fixed $\epsilon > 0$. Covering and Dominating in planar graphs was further explored by Bumpus et al. [16]. They provide $(1 + \epsilon)$ -certified algorithms for DOMINATING SET and H-SUBGRAPH-FREE-DELETION, which for any ϵ run in time $f(1/\epsilon) \cdot n^{O(1)}$ on minor-closed classes of graphs of bounded local treewidth.

Symbol	Meaning
FVS-ρCL	Feedback Vertex Set problem for breaking every cycle of length exactly ρ
ℓ	Outerplanarity parameter; typically is set to $\frac{1}{\epsilon}$
k	Solution size
r	Radius parameter in r -Dominating Set; each node dominates vertices at distance at most r
n	Number of vertices in the graph
m	Number of edges in the graph
μ	The μ -neighbourhood of a vertex (e.g., all vertices within distance μ)
γ	Perturbation parameter, $\gamma \in \mathbb{N}$

Table 1: Summary of Notation

2 Preliminaries

In this section, we provide the reader with the necessary information in order to follow the paper. We start with a reminder of Graph Theory. After that, we move to more advanced definitions in the field of Combinatorial Optimization. Finally, we get on topic by introducing perturbation resilience and certified algorithms from the field of Beyond the Worst-Case Analysis of Algorithms.

2.1 Graph Theory

In this section, the authors refer to Bondy and Murty [13]. A *graph* G is an ordered pair denoted as $(V(G), E(G))$. The set $V(G)$ is called a set of *vertices* and the set $E(G)$ is a set of *edges*. The incidence function ψ_G associates each edge with an unordered pair of (not necessarily unique) vertices. The ordered pair and the incidence function define the graph G . We are going to use n as the cardinality of the set V and m for the set of edges E . With each edge e of G , let there be associated a real number $w(e)$, called its *weight*. Then G , together with the weighting function $w : E \rightarrow \mathbb{R}$, is called a *weighted graph*, and denoted (G, w) .

A graph can be *embedded in the plane*, or is *planar*, if it can be drawn on the plane in a way that edges intersect each other only at their ends. Such a drawing is called a *planar embedding* of the graph. A graph G is called *outerplanar* if it has a planar embedding $\tilde{G}$ in which all the vertices lie on the boundary of the outer face. This is also called a *1-outerplanar graph*. Two forbidden minors of that class are the K_4 and $K_{2,3}$ graphs. In the next subsection, we familiarize the reader with Baker's approach [5]. Here, we define an ℓ -*outerplanar* to be a graph that has a planar embedding in which the vertices belong to at most ℓ concentric layers. Every planar graph is ℓ -*outerplanar* for some ℓ . Given a graph G , we look for the ℓ -*outerplanar* embedding of G for which ℓ is minimal.

Let $G = (V, E)$ be a graph and let $v \in V$. The *closed r -neighbourhood* of v , denoted by $N^r[v]$, is the set of all vertices in G that are at distance at most r from v :

$$N^r[v] = \{u \in V \mid \text{dist}_G(u, v) \leq r\}.$$

2.2 Combinatorial Optimization

In this section, we refer to the book on Combinatorial Optimization by Korte and Vygen [29] and Williamson and Shmoys [39]. Some definitions can be traced to the end of the book by Cygan et al. [21].

2.2.1 NP-Complete Problems: Definitions and Variants

Two problems are of interest, namely the Feedback Vertex Set and the r -Dominating Set. The latter is a generalization of the Dominating Set problem. All of those problems belong to the class of **NP-complete** problems. The decision variant of the FVS problem can be defined as:

Definition 2.1: Feedback Vertex Set (FVS)

Input: A graph G and an integer k .

Question: Does there exist a set X of at most k vertices of G such that $G - X$ is a forest?

The weighted optimization version of the problem can be defined as:

Definition 2.2: Minimum Weight Feedback Vertex Set (MWFVS)

Instance:

- An undirected weighted graph (G, w) , where $G = (V, E)$ and $w : V \rightarrow \mathbb{R}$.

Task: Find a vertex set $X \subseteq V$ such that:

- $G \setminus X$ is acyclic (i.e., a forest)
- The sum of the weights of the vertices in X is minimized.

The suffix 'BCL' in the definitions of the problems FVS-BCL and MWFVS-BCL indicates that they pertain to the bounded cycle length variant.

Definition 2.3: FVS-BCL

Input: A graph G , integer k and an integer ρ .

Question: Does there exist a set X of at most k vertices such that every cycle of length ρ contains a vertex in X ?

The following is the weighted version of the problem; we define it as an optimization task.

Definition 2.4: MWFVS-BCL

Instance:

- An undirected weighted graph (G, w) , where $G = (V, E)$ and $w : V \rightarrow \mathbb{R}$ and an integer ρ .

Task: Find a vertex set $X \subseteq V$ such that:

- Every cycle in $G \setminus X$ of length ρ has a vertex in X .
- The sum of the weights of the vertices in X is minimized.

r -Dominating Set is a generalization of Dominating Set. Setting the parameter r to one would transform the former problem into the latter.

Definition 2.5: The r -Dominating Set Problem.

Input: A graph $G = (V, E)$, an integer k and an integer r .

Question: Does there exist a subset $D \subseteq V$ of size at most k , such that every vertex $v \in V$ is within distance at most r from some vertex in D ; that is,

$$\forall v \in V, \exists u \in D \text{ such that } \text{dist}_G(u, v) \leq r.$$

The weighted version of the problem is defined as follows:

Definition 2.6: The Weighted r -Dominating Set Problem.

Input: A graph $G = (V, E)$, a weight function $w : V \rightarrow \mathbb{N}$, an integer W , and an integer r .

Question: Does there exist a subset $D \subseteq V$ such that the total weight of D is at most W , and every vertex $v \in V$ is within distance at most r from some vertex in D ; that is,

$$\sum_{u \in D} w(u) \leq W \quad \text{and} \quad \forall v \in V, \exists u \in D \text{ such that } \text{dist}_G(u, v) \leq r.$$

2.2.2 Approximation Algorithms and Complexity Classes

In the previous section all of the problems were NP-complete. In order to cope with the hardness of those problems, one can turn to *approximation algorithms*. Let Π be an optimization problem with nonnegative weights and $q \geq 1$. A q -factor approximation algorithm for Π is a polynomial-time algorithm A for Π , such that

$$\frac{1}{q} \cdot \text{OPT}(I) \leq A(I) \leq q \cdot \text{OPT}(I)$$

for all instances I of Π . The first inequality shown is true for maximization problems, while the second one holds for minimization problems. We say A has a performance ratio of q . If a problem falls into that category, we say that it is APX-COMPLETE.

The previous paragraph introduced the class of problems APX in the context of approximation algorithms. Now, we introduce another important class: PTAS, which consists of algorithms that run in polynomial time and produce solutions arbitrarily close to optimal, depending on a trade-off parameter.

Definition 2.7: Polynomial-Time Approximation Scheme (PTAS)

A polynomial-time approximation scheme (PTAS) is a family of algorithms A_ϵ where there is an algorithm for each $\epsilon > 0$, such that A_ϵ is a $(1 + \epsilon)$ -approximation algorithm (for minimization problems). The running time of the algorithm A_ϵ is polynomial, but can depend arbitrarily on $1/\epsilon$.

The dependence, mentioned in Def. 2.7, on $\frac{1}{\epsilon}$ could be exponential or even worse. Thus an algorithm running in time $O(n^{1/\epsilon})$ counts as a PTAS.

Every problem with a PTAS is also in APX. The relation between these two classes is similar to the one between P and NP. For more on the topic, such as APX-completeness, L-reductions and the PCP-theorem, we refer to Korte and Vygen [29].

An important distinction in the complexity zoo is made in Cesati and Trevisan [17]. To the best of the knowledge of the author, Cesati and Trevisan first describes what an efficient PTAS is.

Definition 2.8: Efficient Polynomial-Time Approximation Scheme (EPTAS)

An efficient PTAS yields a $(1 + \epsilon)$ approximation scheme and has a running time of $f(\epsilon) \cdot n^c$ where f is an arbitrary function and c is a constant.

2.2.3 Baker's approach

This subsection presents the Shifting Technique, originally introduced in Baker's seminal paper [5]. The technique outlines a general algorithm that partitions the graph into a disjoint union of vertex levels.

By analysing slices of these levels and, more specifically, the overlap between these slices, one obtains an approximation ratio for the solution to the optimization problem. The slices make up several levels that are specifically chosen for a problem and it should hold that an exact solution can be found in each one slice. It must also hold that this solution can be found in optimal time.

There is also another way to view this approach. Instead of analysing the overlap between slices, one could partition the graph by removing said overlap. In this way, the resulting disjoint union may not cover the entire optimal solution, but it can be shown that the deviation from optimality is bounded. This approach works for hereditary maximization problems like INDEPENDENT SET, MAX-CUT, MAXIMUM P-MATCHING, etc.

The following section presents one example application. It focuses on the general Shifting Technique, using the VERTEX COVER problem as an illustrative case. This example is explored in great detail, as both the problem and the technique closely resemble the approach required for solving FVS-BCL.

It is important to highlight that the general Shifting Technique is more powerful than the vertex-removal partitioning. For problems like Dominating Set, the latter technique does not yield a PTAS.

However, using the general shifting technique, one can obtain a PTAS for DOMINATING SET in planar graphs. This was done by Marzban and Gu [33]. They designed a PTAS with an approximation ratio of $(1 + 2/\ell)$. The two in the approximation ratio appears because of the overlap on two levels.

What follows is a brief explanation of Baker's shifting technique with VERTEX COVER as an example.

Algorithm 1 Baker's shifting technique

Require: Planar graph $G = (V, E)$, parameter ℓ

Ensure: $(1 + \epsilon)$ -approximation for VERTEX COVER

- 1: Perform BFS from some arbitrary vertex r
 - 2: $S \leftarrow \emptyset$
 $\quad i: \text{shift}; j: \text{slice}$
 - 3: **for** each $i = 0$ to $\ell - 1$ **do**
 - 4: Let $G_{i,j}$ be the subgraph induced on vertices at levels $j \cdot \ell + i$ through $(j + 1) \cdot \ell + i$ for $0 \leq i < \ell$.
 - 5: $\forall j$ compute $S_{i,j}$, the min. VC of $G_{i,j}$.
 - 6: $\forall j$ let $S_i = \bigcup_j S_{i,j}$
 - 7: $S \leftarrow S \cup S_i$
 - 8: **end for**
 - 9: **return** S_{i*} from S with minimum cardinality
-

Analysis of the algorithm:

Let V_i be the set of vertices at levels $i \bmod \ell$:

$$V_i = \{v \in V \mid \text{level}(v) \equiv i \bmod \ell\}$$

Let OPT_i denote $OPT \cap V_i$. We notice that OPT_i are disjoint and $OPT = \bigcup_i OPT_i$. Therefore, we can make the following argument:

1. $\exists q : |OPT_q| \leq \frac{1}{\ell} \cdot OPT = \epsilon |OPT|$
2. $S_q \leq \sum_j S_{q,j}$, for some shift q we have this inequality because in step 5. of the algorithm we define

the set to be a union over all slices and some vertices may appear twice in the boundary layers.

3. $\sum_j S_{qj} \leq \sum_j |OPT \cap V(G_{q,j})|$, we know that $S_{q,j}$ is a min. VC for $G_{q,j}$, while the optimal solution may choose different vertices, especially on the boundary for the global optimality of the solution.
4. $\sum_j |OPT \cap V(G_{q,j})| \leq |OPT| + |OPT_q|$, if we consider the graphs $\{G_{q,j}\}$, then we can argue that each vertex appears once, except those at levels $q \bmod \ell$, which appear twice.

From the first, second and fourth step we can conclude that:

$$S_q \leq |OPT| + |OPT_q| \leq (1 + \epsilon) \cdot |OPT|$$

2.2.4 Treewidth

According to Diestel [25, pp. 354-355] the notions of tree decomposition and treewidth were introduced by Halin [27] under the name of S -functions and Robertson and Seymour [35]. We give brief definitions of these concepts and then discuss bounded local treewidth and linear local treewidth. The last two notions are important for the main theorem we work with, namely Bumpus et al. [16, Theorem 1.7].

From Diestel [25, Section 12.3] we have the following definition of a tree decomposition.

Definition 2.9: Tree Decomposition

Let G be a graph, T a tree, and let $\mathcal{V} = (V_t)_{t \in T}$ be a family of vertex sets $V_t \subseteq V(G)$ indexed by the vertices t of T . The pair $(T, \mathcal{V})$ is called a *tree decomposition* of G if it satisfies the following conditions:

- (T1) $V(G) = \bigcup_{t \in T} V_t$.
- (T2) For every edge $e \in E(G)$, there exists a $t \in T$ such that both ends of e lie in V_t .
- (T3) If $t_1, t_2, t_3 \in T$ satisfy t_2 lies on the unique path between t_1 and t_3 in T , then

$$V_{t_1} \cap V_{t_3} \subseteq V_{t_2}.$$

The *width* of a tree decomposition $(T, \mathcal{V})$ is the size of its largest set V_t minus one. The *treewidth* $\text{tw}(G)$ of a graph G is the minimum width among all possible tree decompositions of G .

Definition 2.10: Bounded local treewidth

A graph G has bounded local treewidth if the subgraph induced by vertices at distance at most d is a function of d ($f : \mathbb{N} \rightarrow \mathbb{R}$) and not n .

Furthermore, if there exists $\lambda \in \mathbb{R}$, such that f can be defined as $f : r \rightarrow \lambda \cdot r$, then G would have linear local treewidth.

Building on the work of Eppstein [26], Demaine and Hajiaghayi [24] prove that apex-minor-free graphs have linear local treewidth. Apex graphs are graphs that can be made planar by the removal of a single vertex. From this follows that planar graphs also have linear local treewidth.

2.2.5 Linear Programming and Randomized Rounding

One way of approximating solutions is done through Linear Programming (LP). For a more detailed review of LP, the reader can consult Korte and Vygen [29, Chapter 3]. One solves a linear relaxation, which

provides fractional values as solutions. Afterwards, the fractional solution is rounded to an integral one, but rather than doing it deterministically, this process is done in a randomized fashion. This technique is called *randomized rounding*. This framework provides a way to obtain certified algorithms, for a more in depth look for randomized rounding the reader should consult Makarychev and Makarychev [31].

2.3 Beyond the Worst-Case Analysis of Algorithms

In this section we explore perturbation-resilience, also known as γ -*stability* or Bilu-Linial stability. The authors refer to the original paper of the two authors Bilu and Linial [9] as well as a survey on the matter by Makarychev and Makarychev [30]. In a later work, Makarychev and Makarychev define γ -*certified algorithms* [31].

The following definitions for γ -perturbation are for edge-optimization and vertex-optimization problems, respectively. Subsequently, the notions of stability and certified algorithms are introduced specifically for vertex-optimization problems.

Definition 2.11: γ -perturbation for Edge-Optimization problems

Consider a weighted graph $G = (V, E, w)$ with a set of edge weights w_e . An instance $G' = (V, E, w')$ is a γ -perturbation ($\gamma \in \mathbb{R}_{\geq 1}$) of G if $w(e) \leq w'(e) \leq \gamma \cdot w(e)$; that is, if we can obtain a perturbed instance from the original by multiplying the weight of each edge by a number between 1 and γ (the number may vary for each edge).

From now on, the definitions will concentrate on Vertex-Optimization problems.

Definition 2.12: γ -perturbation for Vertex-Optimization problems

Consider a weighted graph (G, w) . For any $\gamma \in \mathbb{R}_{\geq 1}$ a γ -perturbation of the weight function $w : V \rightarrow \mathbb{R}$ is a function $w' : V \rightarrow \mathbb{R}$ satisfying $w(v) \leq w'(v) \leq \gamma \cdot w(v) \forall v \in V$.

Definition 2.13: γ -stability

For any $\gamma \in \mathbb{R}_{\geq 1}$, a γ -stable instance (G, w) of a *vertex-minimization* problem Π is an instance admitting a unique optimal solution S which remains optimal even under γ -perturbations of (G, w) .

Definition 2.14: Certified Algorithm

A γ -certified solution to an instance (G, w) of a weighted vertex-optimization problem Π is a pair (S, w') where w' is a γ -perturbation of w and S is an optimal solution on (G, w') . A γ -certified algorithm for Π maps instances of Π to γ -certified solutions.

2.4 r-Dominating Set; Exact algorithm and PTAS

For FVS-BCL we will give an algorithm that solves the problem on graphs of bounded treewidth. For Dominating Set and its generalizations there are already existing solutions one can use. The r-Dominating Set problem is one generalization of Dominating Set. We mention an algorithm which solves the generalized version. In this work the parameter r is considered to be constant.

There is an algorithm that solves r-Dominating Set using dynamic programming which is realized by

Borradaile and Le [15, Theorem 1, Appendix B.]. It is a generalization of the work done in van Rooij et al. [37]. As it is argued in Borradaile and Le [15, Appendix B.], the running time needed to solve r -Dominating Set is $O((2r + 1)^{\text{tw} + 1} \text{tw}^2 n)$.

We showed in subsection 2.2.3 how one can use an exact algorithm for graphs of bounded treewidth to get an approximation in a planar graph. In Borradaile and Le [15] it is mentioned that one can use Baker's technique to obtain a PTAS. In the MIT lecture notes of [22], there is an explanation of how to get to that approximation. Essentially, instead of two layers of overlap between subgraphs, one would have to increase that overlap to $r + 1$. This technique will yield results only for sufficiently small values for the parameter r .

3 (E)PTAS for combinatorial optimization problems using Baker's technique

This section presents an EPTAS for both the unweighted and weighted versions of the FVS-BCL problem. We begin with the unweighted case, focusing on the algorithm's correctness and efficiency.

For the special case of breaking 4-cycles—denoted as FVS-4CL—exact solutions on subgraphs of bounded treewidth are obtained using a dynamic programming approach. In contrast, the general FVS-BCL problem, where cycles of arbitrary length ρ must be broken, is handled using Monadic Second Order Logic (MSOL) and Courcelle's Theorem.

The running time of the overall approach remains polynomial in the size of the input.

Some notation used throughout this section is introduced here. The 4-cycle-breaking problem is denoted as FVS-4CL and is used consistently in subsections 3.1 and 3.2, regardless of whether the input is weighted. In subsection 3.3, where arbitrary cycle lengths are considered, the notation FVS-BCL is used. This distinction is maintained in subsequent chapters. Unless stated otherwise, FVS-4CL refers to the weighted variant of the 4-cycle-breaking problem.

3.1 Baker's technique for unweighted FVS-4CL on planar graphs

In this subsection we provide the necessary definitions, lemmas and theorems to demonstrate how to obtain an EPTAS for the problem we defined in 2.3, namely FVS-4CL. In the preliminaries we gave two examples of how to construct algorithms using Baker's technique and how to further analyse them. In the same fashion we now construct Algorithm 2 and reason for its effectiveness.

In the following algorithm we use the letter ℓ for the outerplanarity of each subgraph, i.e. each $G_{i,j}$ that we define on line 4 in Algorithm 2 is ℓ -outerplanar.

Algorithm 2 Baker's technique for the unweighted FVS-4CL

Require: Planar graph $G = (V, E)$, parameter $\ell \leftarrow \frac{1}{\epsilon}$

Ensure: $(1 + \epsilon)$ -approximation for FVS-4CL

- 1: Perform BFS from some arbitrary vertex r
 - 2: $S \leftarrow \emptyset$
 i : *shift*; j : *slice*
 - 3: **for** each $i = 0$ to $\ell - 1$ **do**
 - 4: Let $G_{i,j}$ be the subgraph induced on vertices at levels $j \cdot \ell + i$ through $(j + 1) \cdot \ell + i + 1$ for $0 \leq i < \ell$.
 - 5: $\forall j$ compute $S_{i,j}$, the min. unweighted FVS-4CL of $G_{i,j}$ using Algorithm 4.4 as a subroutine (the weights are equal to ones).
 - 6: $\forall j$ let $S_i = \bigcup_j S_{i,j}$
 - 7: $S \leftarrow S \cup S_i$
 - 8: **end for**
 - 9: **return** S_{i*} from S with minimum cardinality
-

In the analysis of the algorithm we are going to need three lemmas in order to prove that Algorithm 2

obtains a $(1 + 2/\ell)$ -approximation of the optimum solution.

Lemma 3.1: Bound on the Optimum Solution in Subgraphs (Unweighted Case)

Let $S_{i,j}$ be the minimum unweighted feedback vertex set (FVS-4CL) for subgraph $G_{i,j}$, computed using Algorithm 2, and let $F \equiv \text{OPT}$ be the optimum solution for the full graph G .

Define $F_{i,j} := F \cap V(G_{i,j})$, i.e., the restriction of the global solution F to the subgraph $G_{i,j}$.

Then:

$$|S_{i,j}| \leq |F_{i,j}|$$

Proof. We prove this in three steps by fixing a particular shift i and slice j , and analysing the corresponding subgraph $G_{i,j}$, which is induced on the set of vertices at levels from $j \cdot \ell + i$ to $(j + 1) \cdot \ell + i + 1$.

1. **Locality of 4-cycles:** Any 4-cycle that is fully contained in $G_{i,j}$ must be broken by vertices within $G_{i,j}$. That is, no vertex $v \in V(G) \setminus V(G_{i,j})$ can break 4-cycles entirely within $G_{i,j}$.
2. **Feasibility of restricted solution:** Since F is a valid global solution for the 4-cycle-breaking problem in G , its restriction to $G_{i,j}$, namely $F_{i,j} = F \cap V(G_{i,j})$, must also be a feasible 4-cycle-breaking set for the subgraph $G_{i,j}$.
3. **Optimality of $S_{i,j}$:** By assumption, $S_{i,j}$ is the minimum unweighted solution for $G_{i,j}$. Since $F_{i,j}$ is also feasible (though not necessarily optimal), we have:

$$|S_{i,j}| \leq |F_{i,j}|.$$

This argument holds for arbitrary values of i and j , so we conclude that:

$$\forall i, j : \quad |S_{i,j}| \leq |F_{i,j}|.$$

□

One must confront the fact notation is abused throughout the paper, especially when it comes down to j , the slice variable. When we say that a statement holds for all values i and j , we mean that it holds for the subgraphs that these variables outline. We will sketch an example to deepen our understanding of Lemma 3.1. Moreover, we are going to explore one avenue that gives rise to an inequality in the lemma.




Figure 1: BFS tree, showing the distance from the root for each layer.

Let G look like the graph we sketched on the previous page. The vertical dots help us clarify the levels

while the horizontal signify that there may be other vertices of interest. However, there are vertices we would like to inspect, specifically vertices with labels 14 and 17. Let $v_{14}, v_{17} \in F$. What follows is that $v_{14}, v_{17} \in F_{0,0}$. The cycles that give us reason to claim that $v_{14} \in F$ are all within $G_{0,0}$. Therefore, it holds that Algorithm 2 will have $v_{14} \in S_{0,0}$. However, the same does not hold for v_{17} , the cycles that it is a part of are in $G_{0,1}$ and thus $v_{17} \notin S_{0,0}$, while $v_{17} \in F_{0,0}$. This example should not distract the reader from the fact the real reason that $|S_{0,0}| \leq |F_{0,0}|$ is because $S_{0,0}$ is by definition the minimum 4-cycle-breaking set for $G_{0,0}$, while $F_{0,0}$ is, as we previously argued, a feasible 4-cycle-breaking set.

What follows from Lemma 3.1 is a result that holds for any shift i .

Result 3.2: Optimum solution bound for the whole graph (Unweighted Case)

Let S_i be the union of optimal solutions defined on line 6 of Algorithm 2 for some shift i . Let $F \equiv OPT$ be the optimum solution in G and let $F_{i,j} = G_{i,j} \cap F$. For those sets it holds that:

$$|S_i| \leq \sum_j |S_{i,j}| \leq \sum_j |F_{i,j}|.$$

The first inequality in Result 3.2 stems from the fact that some vertices in the $S_{i,j}$ sum may appear twice. This pertains to vertices that are located on the boundary. For example if the $v_{30} \in S_{0,1} \wedge v_{30} \in S_{0,2}$, then it would be counted once in S_0 , but twice in $\sum_j S_{0,j}$. The second inequality is a result from Lemma 3.1 where we argue that $S_{i,j}$ is an optimum solution for $G_{i,j}$ while $F_{i,j}$ is a 4-cycle-breaking set, but not one of necessarily minimum cardinality.

In Lemma 3.4 we look at the bound of the union of the $F_{i,j}$ sets. Before that, however, we make an argument for the vertices on the boundaries in Lemma 3.3.

Lemma 3.3: Bound for the vertices on the boundaries (Unweighted Case)

Let $F_i = F \cap \{\text{vertices at levels } i \bmod \ell\}$. The sets F_i are disjoint and $\bigcup_i F_i = F$.

We claim that:

$$\exists q \in \{0, 1, \dots, \ell - 1\} : |F_q| + |F_{q+1}| \leq \frac{2}{\ell} \cdot |F|.$$

Proof. Assume for the sake of contradiction that

$$\forall i \in \{0, 1, \dots, \ell - 1\} : |F_i| + |F_{i+1}| > \frac{2}{\ell} |F|. \quad (1)$$

Then we argue in three steps:

1. Summing 1 over all $i = 0, \dots, \ell - 1$ gives

$$\sum_{i=0}^{\ell-1} (|F_i| + |F_{i+1}|) > \sum_{i=0}^{\ell-1} \frac{2}{\ell} |F| = 2|F|.$$

2. On the other hand, since the sets $F_0, \dots, F_{\ell-1}$ are pairwise disjoint and $\bigcup_i F_i = F$, each F_i appears

exactly twice in the left-hand sum. Therefore

$$\sum_{i=0}^{\ell-1} (|F_i| + |F_{i+1}|) = 2 \sum_{i=0}^{\ell-1} |F_i| = 2|F|,$$

yielding a contradiction.

3. Hence the assumption $\forall i : |F_i| + |F_{i+1}| > \frac{2}{\ell}|F|$ must be false. By the pigeonhole principle there is at least one index

$$q \in \{0, 1, \dots, \ell - 1\} \quad \text{such that} \quad |F_q| + |F_{q+1}| \leq \frac{2}{\ell}|F|.$$

□

Now we argue for the size of the sets $F_{i,j}$. It is natural to try and tie their size to the size of the optimal solution F using Lemma 3.3. Whenever we refer to F_i and F_{i+1} , one should keep in mind that the second term may wrap around and become F_0 .

Lemma 3.4: Bound for the cardinality of the intersection sets (Unweighted Case)

Let $F \equiv OPT$ be the optimal solution and $F_{i,j} = F \cap G_{i,j}$ be the intersection sets with the subgraphs. Then we have:

$$\sum_j |F_{i^*,j}| \leq |F| + \frac{2}{\ell} \cdot |F|$$

for some specific integer i^* .

Proof. If we look at Fig. 1, then we notice that any vertex that is located in the overlap layers would be counted twice in the sum $\sum_j |F_{i,j}|$. The interior vertices are counted once towards the optimum solution sum. Therefore, we have that:

1. $\sum_j |F_{i,j}| = |F| + |\{\text{vertices on the boundary}\}| = |F| + |F_i| + |F_{i+1}|$ for $i \in \{0, 1, \dots, \ell - 1\}$.
2. For the RHS of the equation in step 1. we argue that:

$$\exists i^* : |F_{i^*}| + |F_{i^*+1}| \leq \frac{2}{\ell} \cdot |F|$$

using Lemma 3.3.

3. For that i^* it holds that $\sum_j |F_{i^*,j}| \leq |F| + \frac{2}{\ell}|F|$.

□

Finally, we chain the inequalities presented in Lemma 3.1, Result 3.2 and Lemma 3.4, and get that for some slice i^* , it holds that:

$$|S_{i^*}| \leq \sum_j |S_{i^*,j}| \leq \sum_j |F_{i^*,j}| \leq |F| + \frac{2}{\ell} \cdot |F| = (1 + \frac{2}{\ell}) \cdot |F|$$

The inequality

$$|S_{i*}| \leq \left(1 + \frac{2}{\ell}\right) \cdot |F|$$

implies that Algorithm 2 returns a $(1 + \frac{2}{\ell})$ -approximate solution. Since we set $\ell \leftarrow \frac{1}{\epsilon}$, this yields a $(1 + 2\epsilon)$ -approximation for the unweighted planar FVS-4CL problem.

Running time:

Lines 1 and 9 of Algorithm 2 can be computed in linear time $O(n)$. The loop that starts on line 3 has a running time that is dominated by step 5 where we call Algorithm 4.4 (with weights equal to ones) as a subroutine. That algorithm takes as input subgraphs that are $\ell + 2$ -outerplanar. As we argued in the Preliminaries section, such graphs have bounded treewidth of tw , Bodlaender [11, Theorem 83]. Therefore, Algorithm 2 has a running time of $O(2^{\text{tw}^2} n)$.

3.2 Baker's technique for the weighted FVS-4CL on planar graphs

In this subsection we look at the weighted version of the problem. Firstly, we construct Algorithm 3 which differs from Algorithm 2 only on step 5. On line 5 we construct Algorithm 4 and use it as a subroutine to get the solution we want.

Algorithm 3 Baker's technique for the weighted FVS-4CL

Require: Planar graph $G = (V, E)$, parameter $\ell \leftarrow \frac{1}{\epsilon}$, $w : V \rightarrow \mathbb{R}$

Ensure: $(1 + \epsilon)$ -approximation for FVS-4CL

- 1: Perform BFS from some arbitrary vertex r
 - 2: $S \leftarrow \emptyset$
i: shift; j: slice
 - 3: **for** each $i = 0$ to $\ell - 1$ **do**
 - 4: Let $G_{i,j}$ be the subgraph induced on vertices at levels $j \cdot \ell + i$ through $(j + 1) \cdot \ell + i + 1$ for $0 \leq i < \ell$.
 - 5: $\forall j$ compute $S_{i,j}$, the min. weighted FVS-4CL of $G_{i,j}$ using Algorithm 4.4 as a subroutine.
 - 6: $\forall j$ let $S_i = \bigcup_j S_{i,j}$
 - 7: $S \leftarrow S \cup S_i$
 - 8: **end for**
 - 9: **return** S_{i*} from S , such that $w(S_{i*}) \leq w(S_i), \forall i \in \{1, 2, \dots, \ell - 1\}$.
-

Much like the proof we did in subsection 3.1, we start off with finding a bound for the optimum solution for each subgraph.

Lemma 3.5: Optimum solution bound (Weighted Case)

Let $G = (V, E)$ be a planar graph and $w : V \rightarrow \mathbb{R}$ be a weight function. Let $S_{i,j}$ be an optimum solution for $G_{i,j}$ for some shift i and slice j , let $F \equiv \text{OPT}$ be the optimum solution in G and let $F_{i,j} = G_{i,j} \cap F$. In the weighted case we claim that:

$$w(S_{i,j}) \leq w(F_{i,j}).$$

Proof. We prove this in three steps by fixing a particular shift i and slice j , and analysing the corresponding

subgraph $G_{i,j}$, which is induced on the set of vertices at levels from $j \cdot \ell + i$ to $(j + 1) \cdot \ell + i + 1$.

1. **Locality of 4-cycles:** Any 4-cycle that is fully contained in $G_{i,j}$ must be broken by vertices within $G_{i,j}$. That is, no vertex $v \in V(G) \setminus V(G_{i,j})$ can break 4-cycles entirely within $G_{i,j}$.
2. **Feasibility of restricted solution:** Since F is a valid global solution for the 4-cycle-breaking problem in G , its restriction to $G_{i,j}$, namely $F_{i,j} = F \cap G_{i,j}$, must also be a feasible 4-cycle-breaking set for the subgraph $G_{i,j}$.
3. **Optimality of $S_{i,j}$:** By assumption, $S_{i,j}$ is the minimum-weight solution for $G_{i,j}$. Since $F_{i,j}$ is also feasible (though not necessarily optimal), we have:

$$w(S_{i,j}) \leq w(F_{i,j}).$$

This argument holds for arbitrary values of i and j , so we conclude that:

$$\forall i, j : \quad w(S_{i,j}) \leq w(F_{i,j}).$$

□

Result 3.6: Optimum solution bound for the whole graph (Weighted Case)

Let S_i be the union of optimal solutions defined on line 6 of Algorithm 3 for some shift i . Let $F \equiv OPT$ be the optimum solution in G and let $F_{i,j} = G_{i,j} \cap F$. For those sets it holds that:

$$w(S_i) \leq \sum_j w(S_{i,j}) \leq \sum_j w(F_{i,j}).$$

Proof. This result immediately follows from the fact that when we define S_i as the union of the subgraphs some vertices, specifically those on the overlapping layers, will have their weights added twice to the sum $\sum_j w(S_{i,j})$. If we inspect the LHS, however, each vertex and its weight will be counted only once towards $w(S_i)$. Therefore, it holds that $w(S_i) \leq \sum_j w(S_{i,j})$ and from Lemma 3.5 we chain the inequalities to get Result 3.6.

□

Lemma 3.7: Bound on the Weight of Boundary Vertices (Weighted Case)

Let $G = (V, E)$ be a planar graph, and let $w : V \rightarrow \mathbb{R}$ be a weight function on the vertices. Let $F \subseteq V$ be a feedback vertex set, and define the sets F_i as:

$$F_i := F \cap \{\text{vertices at BFS level } i \bmod \ell\}$$

for each $i \in \{0, 1, \dots, \ell - 1\}$. Then, the sets F_i are disjoint, and:

$$F = \bigcup_{i=0}^{\ell-1} F_i$$

We claim that:

$$\exists q \in \{0, 1, \dots, \ell - 1\} \text{ such that } w(F_q) + w(F_{q+1}) \leq \frac{2}{\ell} \cdot w(F)$$

Proof. Assume by way of contradiction that the following inequality holds for all $i \in \{0, 1, \dots, \ell - 1\}$ (with indices taken modulo ℓ):

$$w(F_i) + w(F_{i+1}) > \frac{2}{\ell} \cdot w(F) \quad (2)$$

1. From the definitions of the sets F_i , we know:

$$F_i \cap F_j = \emptyset \quad \text{for all } i \neq j$$

$$F = \bigcup_{i=0}^{\ell-1} F_i$$

Therefore, each vertex in F is in exactly one F_i , and:

$$\sum_{i=0}^{\ell-1} w(F_i) = w(F)$$

When summing all pairs $w(F_i) + w(F_{i+1})$, each term $w(F_i)$ appears exactly twice (once in the pair (F_{i-1}, F_i) and once in (F_i, F_{i+1})). So:

$$\sum_{i=0}^{\ell-1} (w(F_i) + w(F_{i+1})) = 2 \cdot w(F)$$

2. Now, under the assumption of inequality (2), we would have:

$$\sum_{i=0}^{\ell-1} (w(F_i) + w(F_{i+1})) > \sum_{i=0}^{\ell-1} \frac{2}{\ell} \cdot w(F) = 2 \cdot w(F)$$

which contradicts the conclusion from step 1 that this sum equals $2 \cdot w(F)$.

3. Therefore, our assumption must be false. By the contrapositive, there must exist at least one index

$q \in \{0, 1, \dots, \ell - 1\}$ such that:

$$w(F_q) + w(F_{q+1}) \leq \frac{2}{\ell} \cdot w(F)$$

This completes the proof. □

Lemma 3.8: Bound for the weight of the intersection sets (Weighted Case)

Let $F \equiv OPT$ be the optimal solution of the FVS-4CL problem. Let $F_{i,j} = F \cap G_{i,j}$ be the intersection sets with the subgraphs. Then we have:

$$\sum_j w(F_{i^*,j}) \leq w(F) + \frac{2}{\ell} \cdot w(F)$$

for some specific integer i^* .

Proof. The interior vertices of each subgraph have their weights counted only once towards the sum $\sum_j w(F_{i,j})$. The same does not hold for the vertices on the boundary, they are counted once in $F_{i,j}$ and once in $F_{i,j+1}$. To account for that we rewrite the sum on the LHS as:

1. $\sum_j w(F_{i,j}) = w(F) + w(\{\text{vertices on the boundary}\})$.
2. From Lemma 3.7 we have an integer $i^* \in \{1, 2, \dots, \ell - 1\}$, such that:
 $w(F_{i^*}) + w(F_{i^*+1}) \leq \frac{2}{\ell} \cdot w(F)$.
3. From the previous two steps we get:

$$\sum_j w(F_{i,j}) = w(F) + w(\{\text{vertices on the boundary}\}) \leq \frac{2}{\ell} \cdot w(F)$$

□

Finally, we chain the inequalities we obtained in Lemma 3.5, Result 3.6, and Lemma 3.8 to argue that for some $i^* \in \{1, 2, \dots, \ell - 1\}$:

$$w(S_{i^*}) \leq \sum_j w(S_{i^*,j}) \leq \sum_j w(F_{i^*,j}) \leq w(F) + \frac{2}{\ell} \cdot w(F)$$

.

Running time:

The only difference from the unweighted case comes from line 5 of Algorithm 3. However, using Algorithm 4.4 we get that the running time of the algorithm is still $O(2^{\text{tw}^2} \cdot n)$.

3.3 Baker's technique for weighted FVS-BCL of arbitrary cycle length

This subsection extends the concepts discussed in subsection 3.2 by examining the outcome of attempting to break all weighted cycles of length ρ .

Algorithm 4 Baker's technique for the weighted FVS-BCL

Require: Planar graph $G = (V, E)$, parameter $\ell \leftarrow \frac{1}{\epsilon}$, $w : V \rightarrow \mathbb{R}$, ρ - integer specifying cycles of what length to break

Ensure: $(1 + \epsilon)$ -approximation for FVS-BCL

- 1: Perform BFS from some arbitrary vertex r
 - 2: $S \leftarrow \emptyset$
 i : *shift*; j : *slice*
 - 3: **for** each $i = 0$ to $\ell - 1$ **do**
 - 4: Let $G_{i,j}$ be the subgraph induced on vertices at levels $j \cdot \ell + i$ through $(j+1) \cdot \ell + i + \lfloor \frac{\rho}{2} \rfloor$ for $0 \leq i < \ell$.
 - 5: $\forall j$ compute $S_{i,j}$, the min. weighted FVS-BCL of $G_{i,j}$ using Algorithm 5 as a subroutine.
 - 6: $\forall j$ let $S_i = \bigcup_j S_{i,j}$
 - 7: $S \leftarrow S \cup S_i$
 - 8: **end for**
 - 9: **return** S_{i*} from S , such that $w(S_{i*}) \leq w(S_i), \forall i \in \{1, 2, \dots, \ell - 1\}$.
-

The changes between Algorithm 3 and Algorithm 4 pertain to lines 4-7. The main difference is on line 4 where we define the subgraphs we want to inspect. This change enables us to argue that the union of the local solutions, defined on line 6, is indeed a feasible solution for the whole graph.

Lemma 3.9: Optimum solution bound for FVS- ρ CL

Let $G = (V, E)$ be a planar graph, $w : V \rightarrow \mathbb{R}$ be a weight function and ρ an integer. Let $F \equiv OPT$ be the optimum solution in the whole graph for FVS- ρ -BCL and $F_{i,j} = G_{i,j} \cap F$. Let $S_{i,j}$ be an optimum solution for $G_{i,j}$ for some shift i and slice j . We claim that:

$$w(S_i) \leq \sum_j w(S_{i,j}) \leq \sum_j w(F_{i,j}).$$

Proof. We prove this in three steps by fixing a particular shift i and analysing the solution constructed from slices $G_{i,j}$, each induced on vertices at levels from $j \cdot \ell + i$ to $(j+1) \cdot \ell + i + \lfloor \frac{\rho}{2} \rfloor$, where ρ is the maximum cycle length we wish to break.

1. **Locality of ρ -cycles:** Any cycle of length ρ that is entirely contained within $G_{i,j}$ must be broken by vertices from within $G_{i,j}$. That is, no vertex $v \in V(G) \setminus V(G_{i,j})$ can break such cycles in $G_{i,j}$.
2. **Feasibility of restricted solution:** Since F is a valid global solution that breaks all cycles of length ρ , the restriction $F_{i,j} = F \cap V(G_{i,j})$ is a feasible solution for the ρ -cycle-breaking problem in $G_{i,j}$. By definition, $S_{i,j}$ is the optimal solution for $G_{i,j}$, so:

$$w(S_{i,j}) \leq w(F_{i,j})$$

Summing over all j , we get:

$$\sum_j w(S_{i,j}) \leq \sum_j w(F_{i,j})$$

3. **Overlap and accounting for duplicates:** The total weight $w(S_i)$ of the combined solution across slices may be smaller than $\sum_j w(S_{i,j})$, because vertices near slice boundaries can appear in multiple

$G_{i,j}$ but are only counted once in S_i . Therefore:

$$w(S_i) \leq \sum_j w(S_{i,j})$$

Chaining both inequalities, we conclude:

$$w(S_i) \leq \sum_j w(S_{i,j}) \leq \sum_j w(F_{i,j}).$$

□

Lemma 3.9: Bound on the Weight of Boundary Vertices for FVS- ρ CL

Let $G = (V, E)$ be a planar graph, and let $w : V \rightarrow \mathbb{R}$ be a weight function on the vertices. Let $F \subseteq V$ be a solution to the FVS- ρ CL problem (i.e., a set that breaks all cycles of length ρ). For a fixed layering from a BFS traversal, define:

$$F_i := F \cap \{\text{vertices at level } i \bmod \ell\}$$

for each $i \in \{0, 1, \dots, \ell - 1\}$. Then the sets F_i are disjoint and together cover all of F :

$$F = \bigcup_{i=0}^{\ell-1} F_i.$$

We claim that there exists some $q \in \{0, 1, \dots, \ell - 1\}$ such that:

$$w(F_q) + w(F_{q+1}) + \dots + w(F_{q+\lfloor \rho/2 \rfloor - 1}) \leq \frac{\lfloor \rho/2 \rfloor}{\ell} \cdot w(F)$$

where the indices are taken modulo ℓ .

Proof. Assume, for contradiction, that the above inequality does not hold for any $i \in \{0, 1, \dots, \ell - 1\}$. That is, suppose:

$$\forall i : w(F_i) + w(F_{i+1}) + \dots + w(F_{i+\lfloor \rho/2 \rfloor - 1}) > \frac{\lfloor \rho/2 \rfloor}{\ell} \cdot w(F)$$

1. From the definition of the sets F_i , we know that:

$$F_i \cap F_j = \emptyset \quad \text{for all } i \neq j, \quad \text{and} \quad \bigcup_{i=0}^{\ell-1} F_i = F$$

Hence,

$$\sum_{i=0}^{\ell-1} w(F_i) = w(F)$$

2. **Counting each vertex $\lfloor \rho/2 \rfloor$ times:** When we sum the weights of all such consecutive windows of length $\lfloor \rho/2 \rfloor$ across ℓ values of i , each set F_j is included in exactly $\lfloor \rho/2 \rfloor$ of these sums (i.e., it

appears in that many overlapping windows). Therefore:

$$\sum_{i=0}^{\ell-1} (w(F_i) + \dots + w(F_{i+\lfloor \rho/2 \rfloor - 1})) = \lfloor \rho/2 \rfloor \cdot w(F)$$

3. **Contradiction:** On the other hand, if every window sum is strictly greater than $\frac{\lfloor \rho/2 \rfloor}{\ell} \cdot w(F)$, then:

$$\sum_{i=0}^{\ell-1} (w(F_i) + \dots + w(F_{i+\lfloor \rho/2 \rfloor - 1})) > \ell \cdot \frac{\lfloor \rho/2 \rfloor}{\ell} \cdot w(F) = \lfloor \rho/2 \rfloor \cdot w(F)$$

which contradicts the equality from step 2.

4. **Conclusion:** Therefore, our assumption must be false. We conclude that:

$$\exists q \in \{0, 1, \dots, \ell - 1\} \text{ such that } w(F_q) + \dots + w(F_{q+\lfloor \rho/2 \rfloor - 1}) \leq \frac{\lfloor \rho/2 \rfloor}{\ell} \cdot w(F)$$

□

Lemma 3.10: Bound for the weight of the intersection sets (Weighted Case)

Let $F \equiv \text{OPT}$ be the optimal solution to the FVS- ρ CL problem for some $\rho \in \mathbb{N}_{\geq 3}$, and let $F_{i,j} = F \cap G_{i,j}$ be the intersection of F with the subgraphs. Then there exists an integer $i^* \in \{0, 1, \dots, \ell - 1\}$ such that:

$$\sum_j w(F_{i^*,j}) \leq w(F) + \frac{\lfloor \rho/2 \rfloor}{\ell} \cdot w(F)$$

Proof. When observing the construction of the subgraphs $G_{i,j}$, we note that the vertices within the overlap regions (boundary layers) are counted $\lfloor \rho/2 \rfloor$ in the total sum $\sum_j w(F_{i,j})$. The remaining (interior) vertices of each $G_{i,j}$ contribute exactly once to the total weight. This gives us:

1. Vertices on the boundary:

$$\sum_j w(F_{i,j}) = w(F) + w(\{\text{vertices on the boundary}\}) = w(F) + w(F_i) + w(F_{i+1}) + \dots + w(F_{i+\lfloor \rho/2 \rfloor - 1})$$

for some $i \in \{0, 1, \dots, \ell - 1\}$.

2. From Lemma 3.9, we know there exists some $i^* \in \{0, 1, \dots, \ell - 1\}$ such that:

$$w(F_{i^*}) + w(F_{i^*+1}) + \dots + w(F_{i^*+\lfloor \rho/2 \rfloor - 1}) \leq \frac{\lfloor \rho/2 \rfloor}{\ell} \cdot w(F)$$

3. Plugging this into step 1, we obtain:

$$\sum_j w(F_{i^*,j}) \leq w(F) + \frac{\lfloor \rho/2 \rfloor}{\ell} \cdot w(F)$$

□

Running time:

In Algorithm 4 which solves FVS-BCL the computation on line 5 dominates the running time. In planar graphs steps 1 and 9 take linear time to the input. Using Courcelle's theorem [19] we argue for the running time of line 5, which uses Alg. 5 as a subroutine. As we will say in the next section, the subroutine runs in time $f(|\phi|, t) \cdot n$ for some computable function f where a tree decomposition of width t is provided. Using a result from Bodlaender [11, Theorem 83] one can incur that the treewidth of each subgraph is bounded tw . Therefore the running time of Alg. 4 is $O(f(\text{tw}) \cdot n)$ for some computable function f .

4 Exact algorithm for FVS-4CL using MSOL

In this section we are looking for an algorithm that finds exact solutions to the FVS-4CL problem in each subgraph. For each subgraph we assume that the treewidth is bounded.

We take a logical approach that was introduced by Courcelle [19]. He shows that graph properties, that can be stated in monadic second-order logic (MSOL), can be solved in linear time for graphs of bounded treewidth. In the following two subsections we briefly explain MSOL and then apply it to our problem. We follow a survey on the matter in Satzinger [36].

4.1 MSOL - a language for graph problems

We want to introduce a language that translates graph problems to formulas, therefore reducing the problem to a satisfiability problem. Consequently, our goal is to find a formula Φ with the property:

The problem Π is solvable for a graph G *iff* $G \models \Phi$, i. e. the formula Φ is satisfied by the graph G .

Problem Π is solvable in polynomial (in this case linear) time for graphs of treewidth at most k .

Monadic second-order logic is an extension of first-order logic. Additionally to having object variables $x, y \dots$, logical connectives $\neg, \wedge, \vee, \neg, \rightarrow, \longleftrightarrow$ and quantifiers for object variables $\forall x, \exists x$, MSOL considers sets of object variables $X, Y \dots$, and relation to them via $\in$.

Whenever graph problems are involved, typically we express solution through vertices and edges. Additionally, as we mentioned, we can construct sets of vertices and edges. Therefore we have the following machinery:

- Object variables: *vertices*: $v_1, v_2, \dots$ and *edges*: $e_1, e_2, \dots$.
- Set variables: sets of vertices $V_1, V_2, \dots$ and sets of edges $E_1, E_2, \dots$.
- Binary relation $\in$: $\{\text{object variable}\} \times \{\text{set variable}\} \rightarrow \{0, 1\}$. Therefore $v \in V$ *iff* v is an element of the corresponding set V .
- The $Adj(e, v_i, v_j)$ relation. It detects whether edge e is an edge from vertex v_i to vertex v_j where $v_i \neq v_j$.

With this machinery, one obtains a first-order language suitable for expressing relations in a graph. To extend this to a traditional second-order logic language, one must add quantifications over set variables.

- Quantification over set variables: $\forall V_i, \forall E_i$ and $\exists V_i, \exists E_i$.

4.2 Exact algorithm for FVS-4CL using MSOL

Courcelle's work was extended to extended monadic second-order logic [2, 14, 20]. These extensions allow to optimize over the size and weight of a free set variable.

$$\begin{aligned}
& \min_{F \subseteq V} |F| \\
& \quad \wedge \forall v \forall u \forall w \forall z \\
& \quad v \in (V \setminus F) \wedge u \in (V \setminus F) \wedge w \in (V \setminus F) \wedge z \in (V \setminus F) \\
& \quad \wedge v \neq w \wedge u \neq z \\
& \quad \wedge \neg((v, u) \in E \wedge (u, w) \in E \wedge (w, z) \in E \wedge (v, z) \in E)
\end{aligned} \tag{3}$$

The minimum weight FVS-4CL can be expressed as:

$$\begin{aligned}
& \min_{F \subseteq V} w(F) : \\
& \quad \forall v \forall u \forall w \forall z \\
& \quad v \in (V \setminus F) \wedge u \in (V \setminus F) \wedge w \in (V \setminus F) \wedge z \in (V \setminus F) \\
& \quad \wedge v \neq w \wedge u \neq z \\
& \quad \wedge \neg((v, u) \in E \wedge (u, w) \in E \wedge (w, z) \in E \wedge (v, z) \in E)
\end{aligned} \tag{4}$$

4.3 Exact algorithm for weighted FVS-BCL using MSOL

The minimum weight FVS- ρ CL can be expressed as:

$$\begin{aligned}
& \min_{F \subseteq V} w(F) : \\
& \quad \wedge \forall v_1 \forall v_2 \dots \forall v_\rho \\
& \quad v_1 \in (V \setminus F) \wedge v_2 \in (V \setminus F) \dots \wedge v_\rho \in (V \setminus F) \\
& \quad \wedge (v_1 \neq v_2 \wedge v_1 \neq v_3 \wedge \dots \wedge v_1 \neq v_\rho) \\
& \quad \wedge (v_2 \neq v_1 \wedge v_2 \neq v_3 \wedge \dots \wedge v_2 \neq v_\rho) \\
& \quad \dots \\
& \quad \wedge (v_\rho \neq v_1 \wedge v_\rho \neq v_2 \wedge \dots \wedge v_\rho \neq v_{\rho-1}) \\
& \quad \wedge \neg((v_1, v_2) \in E \wedge (v_2, v_3) \in E \wedge \dots \wedge (v_{\rho-1}, v_\rho) \in E \wedge (v_\rho, v_1) \in E)
\end{aligned} \tag{5}$$

Theorem 4.1: Courcelle's theorem [19]

Assume that ϕ is a MSOL formula and G is an n -vertex graph, and there is an evaluation of all of the free variables of ϕ . Suppose, that a tree decomposition of G of width t is provided. Then there exists an algorithm that verifies that ϕ is satisfied in G in time $f(|\phi|, t) \cdot n$, for some computable function f .

4.4 Exact algorithm for FVS-4CL using dynamic programming

In order to find an algorithm that solves FVS-4CL, one has to define the following notions. For a more thorough explanation of those concepts, see Bodlaender [12, Section 4].

1. Define the notion of a *solution*. In our case for the FVS-4CL problem, the solution for a graph G is a set F , such that $G - F$ does not contain any cycles of length four.

2. Define the notion of a *partial solution*. For the subgraph $G_i = (V_i, E_i)$, partial solutions F_i are restrictions of solutions given by the subsets $F_i \subseteq V_i$.
3. Define the notion of *extension of a partial solution*. A solution F is an extension of a partial solution F_i for G_i if $F \cap V_i = F_i$.
4. Define the notion of a *characteristic* of a partial solution. For the set X_i the vertices are partitioned twofold.
 - (a) The class $I \subseteq X_i$ corresponds to all vertices that are in the partial solution.
 - (b) The class $Flag \subseteq X_i \times X_i$ is comprised of all of the pairs of vertices $v, u \in X_i : (v, u) \in Flag$ if they have a common neighbour $w \in V_i \setminus X_i$, such that $w \notin I$.

$$ch(G_i, F_i) = (\{v \in X_i : v \in F_i\}, \{(v, u) \in X_i \times X_i : \exists w \in V_i \setminus X_i, (v, w) \in E_i, (u, w) \in E_i, w \notin F_i\})$$

The valuation of a characteristic $ch(G_i, F_i)$ is a table $c[i, I, Flag]$. It represents the minimum weight of a partial solution F_i in G_i and $c[i, I, Flag] \in \mathbb{N} \cup \{\infty\}$. The table keeps the value of infinity if no such set exists.

$$c[i, I, Flag] = \min\{w(W) : W \text{ is a FVS-4CL of } G_i \wedge W \cap X_i = I\}$$

5. Define the notion of a *full set of characteristics*.

The full set for a node X_i is a valuation for each characteristic $I \in \{0, 1\}^{|X_i|}$ and $Flag \in \{0, 1\}^{X_i \times X_i}$. Since there are at most $k+1$ elements, there are at most $2^{(k+1)} \cdot 2^{\binom{k+1}{2}}$ entries. That yields a storage capacity of $O(2^{\binom{k+1}{2} + (k+1)})$.

Leaf node:

For a leaf node i , we have that $X_i = \emptyset$. Hence

$$c[i, \emptyset, \emptyset] = 0$$

Introduce node:

Let i be an introduce node with child node j . We have that $X_i = X_j \cup \{v\}$ for some vertex $v \notin X_j$. There are two possible values.

$$c[i, I \cup \{v\}, Flag] = w(v) + c[j, I, Flag]$$

$$c[i, I, Flag] = \begin{cases} \infty, & \text{if } \exists u, w \in X_i : (u, w) \in Flag, u \notin I, w \notin I, (v, u) \in E_i, (v, w) \in E_i \\ c[j, I, Flag], & \text{otherwise} \end{cases}$$

Forget node:

Let i be a forget node with child node j . We have that $X_i = X_j \setminus \{v\}$ for some vertex $v \in X_j$.

$$c[i, I, Flag] = \min(c[j, I \cup \{v\}, Flag], c[j, I, Flag \cup \{(u, w) : (v, u) \in E_i, (v, w) \in E_i\}])$$

Join node:

Let i be a join node with children j_1 and j_2 . Recall that $X_i = X_{j_1} = X_{j_2}$. Let $Flag_1$ and $Flag_2$ be the $Flag$ set values of X_{j_1} and X_{j_2} respectively.

$$c[i, I, Flag_1 \cup Flag_2] = \begin{cases} \infty, & \text{if } \exists u, w \in X_i : (u, w) \in Flag_1, \\ & (u, w) \in Flag_2, u \notin I, w \notin I \\ c[j_1, I, Flag_1] + c[j_2, I, Flag_2] - w(I), & \text{otherwise} \end{cases}$$

5 m-stitching and Π -m-stitching

This section introduces key concepts that facilitate the construction of certified algorithms. It begins with the definitions of *m-stitching* and Π -*m-stitching*, followed by a presentation of the main theorem from Bumpus et al. [16]. These concepts and the theorem will be used for constructing certified algorithms in the following section. Unless stated otherwise, Π is assumed to be a vertex-minimization problem. We refer to the literature in [16, 38].

5.1 FVS-4CL and 2-stitching

The restriction we imposed on the Feedback Vertex Set problem enables us to look for optimal solutions in a local space. This is what *m-stitching* and Π -*m-stitching* hint at. Here, the variable m is a positive integer that depends on the vertex optimization problem Π . In our case this is tied to the ρ argument, i. e. the length of the cycles one is trying to break.

To introduce Π -*m-stitching*, it is essential to first understand *m-stitching* and *m-stitchable* problems. In the stitching operation, two solutions S_1 and S_2 are given for a vertex optimization problem, along with an induced subgraph J of G . Intuitively, the idea is to take the portion of S_1 that lies outside of J and stitch onto it a suitable part of S_2 within the subgraph J .

The formal definition is as follows:

Def. 5.1: m-stitching

Assume $m \geq 0$ is an integer, J is an induced subgraph of G , and $S_1, S_2 \subseteq V(G)$. Then we define the *m-stitch* of S_2 onto S_1 along J as the set:

$$S_3 := (S_1 \setminus J) \cup (S_2 \cap N_G^m[J]).$$




Figure 2: Vertex set S_1




Figure 3: Vertex set S_2

In the following we show the subgraph J we have chosen and the 2-neighbourhood of the subgraph.

Also, in Fig. 5 we show the first step of the *m-stitch* operation, namely removing the vertices of J from the set S_1 .




Figure 4: Subgraph J of G in light orange and $N_G^2[J]$ in darker orange




Figure 5: Remove J from S_1

What follows is the second step in obtaining set S_3 in Fig. 6 and the stitched set in Fig. 7.




Figure 6: Add $S_2 \cap N_G^2[J]$




Figure 7: Set S_3

A problem Π is *m-stitchable* if the set of feasible solutions are closed under the m-stitching operator. We hope that when we take the union in 5.1 and arrive at the set S_3 , this would too be a feasible solution.

Def. 5.2: m-stitchable

For an integer $m \geq 0$, induced subgraph J and feasible solutions $S_1, S_2 \subseteq V(G)$ of a vertex optimization problem Π we have that the m-stitching of S_2 onto S_1 along J , S_3 is a feasible solution to Π on G .

$$S_3 := (S_1 \setminus J) \cup (S_2 \cap N_G^m[J]).$$

Now that we have a definition for problems Π that are m-stitchable, we introduce Π -m-stitching as defined in [16] and afterwards apply it to the current problem.

Algorithm. 5.3: Π -m-stitching

Input: an instance $(G, w : V(G) \rightarrow \mathbb{N})$ to an m-stitchable vertex optimization problem Π along with a solution S_1 and a vertex set $J \subseteq V(G)$.

Task: find a feasible solution S' to Π on G , such that for all other feasible solutions S^* it holds that $w(S') \leq w(S^* \oplus_{G,J}^2 S_1)$.

Before we proceed with the main theorem that we use to construct a certified algorithm we need one more definition.

Def. 5.4: Guessable

A problem Π is guessable if there is an algorithm that outputs a feasible solution with no requirement for optimality in polynomial time.

The following theorem is the main result in Bumpus et al. [16, Theorem 1.7].

Theorem. 5.5:

Let $\mathbb{G}$ be a minor-closed graph class whose local tree-width is bounded above by a linear function of the form $g : r \rightarrow \lambda \cdot r$ (fixed $\lambda \in \mathbb{R}$ and $r \in \mathbb{N}$). If Π is a vertex-minimization problem and it holds that:

1. Π is guessable.
 2. Π is m-stitchable.
 3. There exists an algorithm A_Π which solves Π -m-stitching in time $f(t) \cdot |V(G)|^{\mathcal{O}(1)}$, where $t = \text{tw}(G[N_G^m(J)])$ and f is a computable function;
- then, for each $\epsilon > 0$ there is a $(1+\epsilon)$ -certified algorithm for Π that runs in time $f(\lambda \cdot m/\epsilon) \cdot |V(G)|^{\mathcal{O}(1)}$.

The proof of theorem 5.5 is beyond the scope of this paper. In this section we only apply the three steps, mentioned to the FVS-BCL problem without generalizing to other vertex-minimization problems.

Firstly, we take a look at the guessable property.

Lemma 5.6: FVS-BCL is guessable

For any planar graph G , the set $V(G)$ is a feasible solution to FVS-BCL.

Proof. The graph $G - X$, where X is the set of all vertices $V(G)$, is the empty graph, therefore there are no 4-cycles left, after selecting that set. $\square$

Lemma 5.7: FVS-BCL is 2-stitchable

Let (G, w) be any instance to the FVS-4CL problem, $J \subseteq V(G)$, some subset of the vertices in the graph and S_1 and S_2 any two feasible solutions to the problem in G . Then the set S_3 :

$$S_3 := (S_1 \setminus J) \cup (S_2 \cap N_G^2[J]).$$

is a feasible solution to the problem.

Proof. Let the vertices $\{v_1, v_2, v_3, v_4\}$ form a 4-cycle. We inspect the possible cases for the 4-cycle $v_1v_2v_3v_4$. In order to prove 5.7 we have to prove that there exists a $v \in \{v_1, v_2, v_3, v_4\}$, such that $v \in S_3$.

- $\exists v' \in \{v_1, v_2, v_3, v_4\}$, such that $v' \in J$. We argue in three steps.
 - (a) Because S_2 is a feasible solution, then $\exists v'' \in \{v_1, v_2, v_3, v_4\}$, such that $v'' \in S_2$.
 - (b) Because there $\exists v' \in \{v_1, v_2, v_3, v_4\} \in J$, then it also holds that $\forall v''' \in \{v_1, v_2, v_3, v_4\}, v''' \in N_G^2[J]$. From that and the previous point it follows that $v'' \in S_2 \cap N_G^2[J]$.
 - (c) Finally, $S_2 \cap N_G^2[J] \subseteq S_3$, therefore $v'' \in S_3$.
- $\forall v' \in \{v_1, v_2, v_3, v_4\}$, it holds that $v' \notin J$. This case follows from the feasibility of S_1 .
 - (a) Because S_1 is a feasible solution, then $\exists v'' \in \{v_1, v_2, v_3, v_4\}$, such that $v'' \in S_1$.
 - (b) From $\forall v' \in \{v_1, v_2, v_3, v_4\} \notin J$ and the previous point, it follows that $v'' \in S_1 \setminus J$.
 - (c) Because $S_1 \setminus J \subseteq S_3$, then $v'' \in S_3$.

□

Now we describe an algorithm to find a feasible solution of minimum weight.

Algorithm 5 Stitch-FVS-4CL

Require: Vertex-weighted planar graph $(G, w : V(G) \rightarrow \mathbb{N})$, a feasible solution S_1 on G and a vertex set $J \subseteq V(G)$.

Ensure: a feasible solution S' on G , such that for all other feasible solutions S^* , we have $w(S') \leq w(S^* \oplus_{G,J}^2 S_1)$.

- 1: Let $H \leftarrow G[N_G^2[J] \setminus (S_1 \setminus J)]$.
 - 2: Let S_2 be the output of algorithm A on input (H, w) .
 - 3: **if** $w(S_2 \oplus_{G,J}^2 S_1) < w(S_1)$ **then**
 - 4: **return** $S_2 \oplus_{G,J}^2 S_1$
 - 5: **else**
 - 6: **return** S_1
 - 7: **end if**
-

The algorithm A on line 5 refers to calling Algorithm 5 that takes as input a graph that has bounded treewidth.

Lemma 5.8: FVS-4CL-2-stitching

Let A be an algorithm that solves FVS-4CL on any vertex-weighted instance $(G, w : V(G) \rightarrow \mathbb{N})$ in time $f(\text{tw}(G)) \cdot |V(G)|^{O(1)}$. Then Algorithm 5 solves FVS-4CL-2-stitching in time $f(\text{tw}(N_G^2[J])) \cdot |V(G)|^{O(1)}$.

Note that it is not obvious that the set $S_2 \oplus_{G,J}^2 S_1$ is a feasible solution in G as S_2 is not necessarily a feasible solution in G . If the last was the case, then the feasibility of the $S_2 \oplus_{G,J}^2 S_1$ would follow from Lemma 5.7. Therefore we first prove feasibility of $S_2 \oplus_{G,J}^2 S_1$. Afterwards, we prove that it is of minimum weight.

Proof. We prove the feasibility of $S_2 \oplus_{G,J}^2 S_1$ by case analysis on any four vertices $v_1v_2v_3v_4$ that form a

4-cycle. Remember that $S_2 \oplus_{G,J}^2 S_1 = (S_1 \setminus J) \cup (S_2 \cap N_G^2[J])$.

1. $\forall v' \in \{v_1, v_2, v_3, v_4\} : v' \in V(G) \setminus J$.

Firstly, because of the feasibility of S_1 , there $\exists v'' \in \{v_1, v_2, v_3, v_4\} : v'' \in S_1$.

Secondly, $\forall v' \in \{v_1, v_2, v_3, v_4\}$, it holds that $v' \in V(G) \setminus J$.

Finally from the first two conditions, using the property $A \setminus B = A \cap B^c$, it holds that $v'' \in S_1 \setminus J$. From this and $S_1 \setminus J \subseteq S_2 \oplus_{G,J}^2 S_1$, it follows that $S_2 \oplus_{G,J}^2 S_1$ is feasible in G .

In Fig. 8 we show (a part of) an example graph G and the feasible solution S_1 for it. In Fig. 9, one can see that 4-cycles that do not have vertices in J , are a part of $S_1 \setminus J \subseteq S_2 \oplus_{G,J}^2 S_1$.




Figure 8: Sets $J \subseteq N_G^1[J] \subseteq N_G^2[J] \subseteq G$; Feasible solution S_1 colored in blue.




Figure 9: $\forall v \in \{v_1, v_2, v_3, v_4\} : v \notin J$

2. $\exists v' \in \{v_1, v_2, v_3, v_4\} : v' \in J$. This also means that in this case $\forall v' \in \{v_1, v_2, v_3, v_4\} : v' \in N_G^2[J]$. There are two cases to consider again. We know there $\exists v'' \in \{v_1, v_2, v_3, v_4\}$, such that $v'' \in S_1$, we still have not considered whether or not it is a part of J .

- (a) $\exists v \in S_1 : v \notin J$:

In this case we can directly apply the previous point. From the properties of the set difference operator we get: $v \in S_1 \setminus J \subseteq S_2 \oplus_{G,J}^2 S_1$. Consult Fig. 10 to see that there exists a vertex $v_{1,4}$ that is a part of the 4-cycle, $v_{1,4} \notin J$, therefore $v_{1,4} \in S_1 \setminus J \subseteq S_2 \oplus_{G,J}^2 S_1$. This means that in the new set $v_{1,4}$ will break that cycle.

- (b) $\forall v \in S_1 : v \in J$:

In line 1 of Algorithm 5, we have the following construction of $H \leftarrow G[N_G^2[J] \setminus (S_1 \setminus J)]$.

$$V(H) = N_G^2[J] \setminus (S_1 \setminus J) \quad (6)$$

$$= N_G^2[J] \cap J \cup (N_G^2[J] \setminus S_1) \quad (7)$$

$$= J \cup (N_G^2[J] \setminus S_1) \quad (8)$$

However, $\forall v \in \{v_1, v_2, v_3, v_4\} : v \in S_1 \wedge v \in J$. Therefore $\{v_1, v_2, v_3, v_4\} \subseteq V(H)$. This means that in line 2 of Algorithm 5 we will add some $v^* \in \{v_1, v_2, v_3, v_4\}$ to S_2 .

Therefore $v \in S_2$, which has the same meaning as $v \in S_2 \cap N_G^2[J]$, and because $S_2 \cap N_G^2[J] \subseteq S_2 \oplus_{G,J}^2 S_1$, we get $v \in S_2 \oplus_{G,J}^2 S_1$.

Consult Fig. 11.




Figure 10: $\exists v \in S_1 : v \notin J$ (Remember we are in the case where $\exists v \in J$)




Figure 11: $\forall v \in S_1 : v \in J$

The proof of optimality of the set $S_2 \oplus_{G,J}^2 S_1$ is similar to the one in Bumpus et al.[16, Lemma 3.2]. It is important to note that we can use the same proof, because of a property of the FVS-4CL problem, namely that a 4-cycle breaking set is closed under taking induced subgraphs.

Remember, we are still trying to prove what we mentioned in Algorithm 5.3, namely that some feasible solution S' (which in our case is the $S_2 \oplus_{G,J}^2 S_1$), is of minimum weight, that is, for all other feasible solutions S^* , it holds that $w(S') \leq w(S^* \oplus_{G,J}^2 S_1)$.

Assume by way of contradiction that there is a feasible solution S_3 for G , such that $w(S_3 \oplus_{G,J}^2 S_1) < w(S_2 \oplus_{G,J}^2 S_1)$. Then it would follow that:

$$\begin{aligned}
w|_{V(F)}(S_2) &= w(S_2) && \text{since } S_2 \subseteq V(F) \\
&= w(S_2 \cap N_G^2[J]) && \text{since } V(F) \subseteq N_G^2[J] \\
&= w(S_1 \setminus J) + w(S_2 \cap N_G^2[J]) - w(S_1 \setminus J) \\
&= w((S_1 \setminus J) \cup (S_2 \cap N_G^2[J])) - w(S_1 \setminus J) && \text{since } V(F) \cap (S_1 \setminus J) = \emptyset \\
&= w(S_2 \oplus_{G,J}^2 S_1) - w(S_1 \setminus J) && \text{by definition of stitch} \\
&> w(S_3 \oplus_{G,J}^2 S_1) - w(S_1 \setminus J) && \text{by assumption on } S_3 \\
&= w((S_1 \setminus J) \cup (S_3 \cap N_G^2[J])) - w(S_1 \setminus J) && \text{by definition of stitch} \\
&= w((S_3 \cap N_G^2[J]) \setminus (S_1 \setminus J)) && \text{since } w(A \cup B) - w(A) = w(B \setminus A) \\
&= w(S_3 \cap (N_G^2[J] \setminus (S_1 \setminus J))) && \text{since } (A \cap B) \setminus C = A \cap (B \setminus C) \\
&= w(S_3 \cap V(F)) && \text{by definition of } F \\
&= w|_{V(F)}(S_3 \cap V(F))
\end{aligned}$$

□

Running time:

The running time of the algorithm is dominated by the call of algorithm A . Therefore, for an integer m and some computable function f , the running time is $O(\text{tw}(N_G^m[J]) \cdot |V(G)|^{O(1)})$.

5.2 FVS-BCL and $(\rho/2)$ -stitching

In this subsection

5.3 r -Dominating Set and $(r+1)$ -stitching

In this subsection we will use the definitions and terminology we have established in the previous section and look at another problem, namely the r -Dominating Set problem. We will argue that in the case of that problem the parameter m has to be set to $r + 1$ in order to maintain feasibility when constructing new sets. Afterwards we will prove that when constructing new sets, one can find a feasible solution which is also optimal. We follow the steps we established in the previous section.

Firstly, prove that the problem is guessable and m -stitchable. Secondly, devise an algorithm $A_{r\text{-DS}}$ that solves the problem in FPT. Finally, using Theorem 5.5, prove that the problem is certified.

Lemma 5.9: r -DS is guessable

For any planar graph G , the set $V(G)$ is a feasible solution to r -DS.

Proof. Let X be the set of all vertices in G , $X \leftarrow V(G)$. Therefore, it is trivially true that $N_G^r[X] = V(G)$. □

Lemma 5.10: r -DS is $(r+1)$ -stitchable

Let (G, w) be any instance to the r -DS problem, $J \subseteq V(G)$, some subset of the vertices in the graph and S_1 and S_2 any two feasible solutions to the problem in G . Then the set S_3 :

$$S_3 := (S_1 \setminus J) \cup (S_2 \cap N_G^{r+1}[J]).$$

is a feasible solution to the problem.

Proof. For a vertex $v \in V(G)$, we have the following cases:

1. $v \in V(G) \setminus N_G^{r+1}[J]$. Because S_1 is a feasible solution, it is clear that v will be dominated by some vertex u in $S_1 \setminus J$. Since $S_1 \setminus J \subseteq S_3$, then $u \in S_3$.
2. $v \in N_G^{r+1}[J]$. Because S_2 is a feasible solution, it is clear that $S_2 \cap N_G^{r+1}[J]$ contains a vertex u which dominates v . Since $S_2 \cap N_G^{r+1}[J] \subseteq S_3$, then $u \in S_3$.

In both cases we have argued that there $\exists u \in S_3$, such that for any vertex $v \in V(G)$, u dominates v .

□

Now we describe an algorithm to find a feasible solution of minimum weight.

Algorithm 6 Stitch- r -DS

Require: Vertex-weighted planar graph $(G, w : V(G) \rightarrow \mathbb{N})$, a feasible solution of r -DS S_1 on G and a vertex set $J \subseteq V(G)$.

Ensure: a feasible solution S' on G , such that for all other feasible solutions S^* , we have $w(S') \leq w(S^* \oplus_{G,J}^{r+1} S_1)$.

- 1: Let $H \leftarrow G[N_G^{r+1}[J] \cup \{q\}]$, where for vertex q it holds that $N_H(q) := N_G(N_G^r[J])$ (Vertex q is adjacent to the vertices at distance exactly $r + 1$ from J).
- 2:

$$w_H(x) = \begin{cases} 0 & \text{if } x \in (S_1 \setminus J) \cup \{q\} \\ w(x) & \text{otherwise} \end{cases}$$

- 3: Let S'_2 be the output of algorithm A on input (H, w) .
 - 4: Let $S_2 = S'_2 \setminus \{q\}$.
 - 5: **if** $w(S_2 \oplus_{G,J}^{r+1} S_1) < w(S_1)$ **then**
 - 6: **return** $S_2 \oplus_{G,J}^{r+1} S_1$
 - 7: **else**
 - 8: **return** S_1
 - 9: **end if**
-

Lemma 5.11: r -DS- $(r+1)$ -stitching

Let A be an algorithm that solves r -stitching on any vertex-weighted instance $(G, w : V(G) \rightarrow \mathbb{N})$ in time $f(\text{tw}(G)) \cdot |V(G)|^{O(1)}$. Then Algorithm 6 solves r -DS in time $f(\text{tw}(N_G^{r+1}[J])) \cdot |V(G)|^{O(1)}$.

The proof of Lemma 5.11 is outside of the scope of this paper as it is similar to the proof of Lemma 5.8. The reader can find a similar proof in Bumpus et al. [16, Lemma 3.3].

Running time:

6 Certified Algorithm

In this section we initiate the study of certified algorithms for FVS-4CL and the more general FVS-BCL. In the final subsection the focus is shifted to the r-Dominating Set problem.

6.1 FVS-4CL

In the second stage, we compute a feasible solution S , set J_κ ¹

6.2 FVS-BCL

6.3 r-Dominating Set

¹A word on the argument κ and the value that we set it to. In our case, for the FVS-4CL problem, m is equal to two. Also, $k = \lceil \frac{2m}{\epsilon} \rceil + 2m \geq 5$. Notice that the set J_κ is of width $k - 2m$, therefore the m -neighbourhood (2-neighbourhood) is of width $k - 2m + 2m = k$. Therefore, in our case, the graphs within we stitch are of width $k = \lceil \frac{2m}{\epsilon} \rceil + 2m \geq 5$. Take for example vertex $v_{2,1}$ in Fig. 12. In order to find every cycle that it is a part of, one would need to consider the layer that it is in, two layers to the left and two layers to the right.




Figure 12: Layer decomposition of graph in Fig. 8

References

- [1] Haris Angelidakis et al. “Bilu-linial stability, certified algorithms and the independent set problem”. In: *arXiv preprint arXiv:1810.08414* (2018) (cit. on pp. 3, 4).
- [2] Stefan Arnborg, Jens Lagergren, and Detlef Seese. “Easy problems for tree-decomposable graphs”. In: *Journal of Algorithms* 12.2 (1991), pp. 308–340 (cit. on p. 25).
- [3] Pranjal Awasthi, Avrim Blum, and Or Sheffet. “Center-based clustering under perturbation stability”. In: *Information Processing Letters* 112.1-2 (2012), pp. 49–54 (cit. on p. 3).
- [4] Vineet Bafna, Piotr Berman, and Toshihiro Fujito. “A 2-Approximation Algorithm for the Undirected Feedback Vertex Set Problem”. In: *SIAM Journal on Discrete Mathematics* 12.3 (1999), pp. 289–297. DOI: 10.1137/S0895480196305124. eprint: <https://doi.org/10.1137/S0895480196305124>. URL: <https://doi.org/10.1137/S0895480196305124> (cit. on p. 3).
- [5] Brenda S. Baker. “Approximation algorithms for NP-complete problems on planar graphs”. In: *J. ACM* 41.1 (Jan. 1994), pp. 153–180. ISSN: 0004-5411. DOI: 10.1145/174644.174650. URL: <https://doi.org/10.1145/174644.174650> (cit. on pp. 3, 5, 7).
- [6] Maria Florina Balcan and Yingyu Liang. “Clustering under perturbation resilience”. In: *SIAM Journal on Computing* 45.1 (2016), pp. 102–155 (cit. on p. 3).
- [7] Maria-Florina Balcan, Nika Haghtalab, and Colin White. “ k -center Clustering under Perturbation Resilience”. In: *arXiv preprint arXiv:1505.03924* (2015) (cit. on p. 3).
- [8] Ann Becker and Dan Geiger. “Optimization of Pearl’s method of conditioning and greedy-like approximation algorithms for the vertex feedback set problem”. In: *Artificial Intelligence* 83.1 (1996), pp. 167–188. ISSN: 0004-3702. DOI: [https://doi.org/10.1016/0004-3702\(95\)00004-6](https://doi.org/10.1016/0004-3702(95)00004-6). URL: <https://www.sciencedirect.com/science/article/pii/0004370295000046> (cit. on p. 3).
- [9] Yonatan Bilu and Nathan Linial. “Are stable instances easy?” In: *Combinatorics, Probability and Computing* 21.5 (2012), pp. 643–660 (cit. on pp. 3, 10).
- [10] Yonatan Bilu et al. “On the practically interesting instances of MAXCUT”. In: *arXiv preprint arXiv:1205.4893* (2012) (cit. on p. 3).
- [11] Hans L Bodlaender. “A partial k -arboretum of graphs with bounded treewidth”. In: *Theoretical computer science* 209.1-2 (1998), pp. 1–45 (cit. on pp. 17, 24).
- [12] Hans L Bodlaender. “Treewidth: Algorithmic techniques and results”. In: *International symposium on mathematical foundations of computer science*. Springer. 1997, pp. 19–36 (cit. on p. 26).
- [13] John Adrian Bondy, Uppaluri Siva Ramachandra Murty, et al. *Graph theory with applications*. Vol. 290. Macmillan London, 1976 (cit. on p. 5).
- [14] Richard B Borie, R Gary Parker, and Craig A Tovey. “Deterministic decomposition of recursive graph classes”. In: *SIAM Journal on Discrete Mathematics* 4.4 (1991), pp. 481–501 (cit. on p. 25).
- [15] Glencora Borradaile and Hung Le. “Optimal dynamic program for r -domination problems over tree decompositions”. In: *arXiv preprint arXiv:1502.00716* (2015) (cit. on p. 11).

- [16] Benjamin Merlin Bumpus, Bart M. P. Jansen, and Jaime Venne. “Fixed-Parameter Tractable Certified Algorithms for Covering and Dominating in Planar Graphs and Beyond”. In: *19th Scandinavian Symposium and Workshops on Algorithm Theory (SWAT 2024)*. Ed. by Hans L. Bodlaender. Vol. 294. Leibniz International Proceedings in Informatics (LIPIcs). Dagstuhl, Germany: Schloss Dagstuhl – Leibniz-Zentrum für Informatik, 2024, 19:1–19:16. ISBN: 978-3-95977-318-8. DOI: 10.4230/LIPIcs.SWAT.2024.19. URL: <https://drops.dagstuhl.de/entities/document/10.4230/LIPIcs.SWAT.2024.19> (cit. on pp. 3, 4, 9, 29–31, 34, 36).
- [17] Marco Cesati and Luca Trevisan. “On the efficiency of polynomial time approximation schemes”. In: *Information Processing Letters* 64.4 (1997), pp. 165–171 (cit. on p. 7).
- [18] Fabián A. Chudak et al. “A primal–dual interpretation of two 2-approximation algorithms for the feedback vertex set problem in undirected graphs”. In: *Operations Research Letters* 22.4 (1998), pp. 111–118. ISSN: 0167-6377. DOI: [https://doi.org/10.1016/S0167-6377\(98\)00021-2](https://doi.org/10.1016/S0167-6377(98)00021-2). URL: <https://www.sciencedirect.com/science/article/pii/S0167637798000212> (cit. on p. 3).
- [19] Bruno Courcelle. “The monadic second-order logic of graphs. I. Recognizable sets of finite graphs”. In: *Information and computation* 85.1 (1990), pp. 12–75 (cit. on pp. 24–26).
- [20] Bruno Courcelle and Mohamed Mosbah. “Monadic second-order evaluations on tree-decomposable graphs”. In: *Theoretical Computer Science* 109.1-2 (1993), pp. 49–82 (cit. on p. 25).
- [21] Marek Cygan et al. *Parameterized algorithms*. Vol. 5. 4. Springer, 2015 (cit. on p. 5).
- [22] Erik Demaine et al. *Lecture 9: Approximation schemes in minor-free graphs I*. Lecture notes. 2011. URL: <https://courses.csail.mit.edu/6.889/fall11/lectures/L09.html> (cit. on p. 11).
- [23] Erik D Demaine and Mohammad Taghi Hajiaghayi. “Bidimensionality: new connections between FPT algorithms and PTASs.” In: *SODA*. Vol. 5. 2005, pp. 590–601 (cit. on p. 3).
- [24] Erik D Demaine and Mohammad Taghi Hajiaghayi. “Equivalence of local treewidth and linear local treewidth and its algorithmic applications”. In: (2003) (cit. on p. 9).
- [25] Reinhard Diestel. “Graph Theory Electronic Edition 2000”. In: (2000) (cit. on p. 9).
- [26] David Eppstein. “Diameter and treewidth in minor-closed graph families”. In: *Algorithmica* 27 (2000), pp. 275–291 (cit. on p. 9).
- [27] Rudolf Halin. “S-functions for graphs”. In: *Journal of geometry* 8 (1976), pp. 171–186 (cit. on p. 9).
- [28] Richard M. Karp. “Reducibility among Combinatorial Problems”. In: *Complexity of Computer Computations: Proceedings of a symposium on the Complexity of Computer Computations, held March 20–22, 1972, at the IBM Thomas J. Watson Research Center, Yorktown Heights, New York, and sponsored by the Office of Naval Research, Mathematics Program, IBM World Trade Corporation, and the IBM Research Mathematical Sciences Department*. Ed. by Raymond E. Miller, James W. Thatcher, and Jean D. Bohlinger. Boston, MA: Springer US, 1972, pp. 85–103. ISBN: 978-1-4684-2001-2. DOI: 10.1007/978-1-4684-2001-2_9. URL: https://doi.org/10.1007/978-1-4684-2001-2_9 (cit. on p. 3).
- [29] Bernhard Korte and Jens Vygen. “Combinatorial Optimization: Theory and Algorithms”. In: 2007. URL: <https://api.semanticscholar.org/CorpusID:265900003> (cit. on pp. 5, 7, 9).
- [30] Konstantin Makarychev. “13 Bilu-Linial Stability”. In: *Perturbations, Optimization, and Statistics* (2017), p. 375 (cit. on pp. 4, 10).

- [31] Konstantin Makarychev and Yury Makarychev. “Perturbation Resilience”. In: *Beyond the Worst-Case Analysis of Algorithms*. Ed. by Tim Roughgarden. Cambridge University Press, 2021, pp. 95–119 (cit. on pp. 3, 4, 10).
- [32] Konstantin Makarychev, Yury Makarychev, and Aravindan Vijayaraghavan. “Bilu–Linial stable instances of max cut and minimum multiway cut”. In: *Proceedings of the twenty-fifth annual ACM-SIAM symposium on Discrete algorithms*. SIAM. 2014, pp. 890–906 (cit. on p. 3).
- [33] Marjan Marzban and Qian-Ping Gu. “Computational Study on a PTAS for Planar Dominating Set Problem”. In: *Algorithms* 6.1 (2013), pp. 43–59. ISSN: 1999-4893. DOI: 10.3390/a6010043. URL: <https://www.mdpi.com/1999-4893/6/1/43> (cit. on p. 8).
- [34] Matúš Mihalák et al. “On the complexity of the metric tsp under stability considerations”. In: *SOFSEM 2011: Theory and Practice of Computer Science: 37th Conference on Current Trends in Theory and Practice of Computer Science, Nový Smokovec, Slovakia, January 22-28, 2011. Proceedings 37*. Springer. 2011, pp. 382–393 (cit. on p. 4).
- [35] Neil Robertson and P.D Seymour. “Graph minors. II. Algorithmic aspects of tree-width”. In: *Journal of Algorithms* 7.3 (1986), pp. 309–322. ISSN: 0196-6774. DOI: [https://doi.org/10.1016/0196-6774\(86\)90023-4](https://doi.org/10.1016/0196-6774(86)90023-4). URL: <https://www.sciencedirect.com/science/article/pii/0196677486900234> (cit. on p. 9).
- [36] Christina Satzinger. “On Tree-Decompositions and their Algorithmic Implications for Bounded-Treewidth Graphs”. PhD thesis (cit. on p. 25).
- [37] Johan MM Van Rooij, Hans L Bodlaender, and Peter Rossmanith. “Dynamic programming on tree decompositions using generalised fast subset convolution”. In: *Algorithms-ESA 2009: 17th Annual European Symposium, Copenhagen, Denmark, September 7-9, 2009. Proceedings 17*. Springer. 2009, pp. 566–577 (cit. on p. 11).
- [38] JC Venne. “Designing (1)-certified algorithms for planar optimization problems with local feasibility properties”. In: (2022) (cit. on p. 29).
- [39] David P Williamson and David B Shmoys. *The design of approximation algorithms*. Cambridge university press, 2011 (cit. on p. 5).