

Communications in Partial Differential Equations

Publication details, including instructions for authors and subscription information:
<http://www.tandfonline.com/loi/lpde20>

Pseudodifferential operators of mellin type

Jeff E. Lewis ^a & Cesare Parenti ^b

^a University of Illinois at Chicago Chicago , Illinois, 60680 , U.S.A.

^b Ist. Matematico S. Pincherle , Bologna, Italy

Published online: 08 May 2007.

To cite this article: Jeff E. Lewis & Cesare Parenti (1983) Pseudodifferential operators of mellin type, Communications in Partial Differential Equations, 8:5, 477-544, DOI: [10.1080/03605308308820276](https://doi.org/10.1080/03605308308820276)
To link to this article: <http://dx.doi.org/10.1080/03605308308820276>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

PSEUDODIFFERENTIAL OPERATORS OF MELLIN TYPE

Jeff E. Lewis
University of Illinois at Chicago
Chicago, Illinois 60680 U.S.A.

Cesare Parenti
Ist. Matematico S. Pincherle
40127 Bologna, Italy

INTRODUCTION

This article studies several algebras of pseudo-differential operators (pdo's) on the half line R^+ . The operators of order 0 include the Hilbert transform on $L^p(R^+)$

$$Hf(t) = \text{p.v.} \frac{1}{\pi} \int_0^\infty \frac{f(s)}{t-s} ds$$

and the point of view is motivated by the representation of H as a Mellin multiplier,

$$Hf(t) = \frac{-1}{2\pi i} \int_{\text{Re } z = 1/p} t^{-z} \cot \pi z \tilde{f}(z) dz$$

where $f(z) = \int_0^\infty t^{z-1} f(t) dt$, $f \in C_0^\infty(R^+)$, is the Mellin transform of f . Since $(-t \frac{d}{dt} f)^\sim(z) = z \tilde{f}(z)$, we consider operators of the form

$$(1) \quad a\left(t, -t \frac{d}{dt}\right) f(t) = \frac{1}{2\pi i} \int_{\operatorname{Re} z = 1/p} t^{-z} a(t, z) \tilde{f}(z) dz$$

where the symbol $a(t, z)$ satisfies suitable smoothness properties in t and holomorphy and growth properties in z . The spectral and continuity properties of the operators (1) on various L^p and Sobolev spaces depend on the domain of holomorphy of the functions $a(0, z)$ and $a(\infty, z)$.

In Chapter 1 we develop a C^∞ theory for the operators (1) which is motivated by the usual theory of pdo's on R^n . The basic operator is

$$\partial = -t \frac{d}{dt}$$

and we study the action of (1) on functions in $C^\infty(R^+)$ which satisfy Holder type continuity conditions at $t = 0$ and $t = \infty$. To include the Hilbert transform, H , we allow the symbols $a(\cdot, z)$ to have a sequence of poles $\{z_k\}$, $\operatorname{Re} z_k \rightarrow -\infty$; this makes the proof of the symbolic calculus theorems for compositions and adjoints more delicate than for the corresponding theorems on R^n . We develop algebras ample enough to include H as well as many Hardy kernel operators of the form

$$Kf(t) = \int_0^\infty k\left(\frac{t}{s}\right) f(s) \frac{ds}{s} = \frac{1}{2\pi i} \int_{\operatorname{Re} z = 1/p} t^{-z} \tilde{k}(z) \tilde{f}(z) dz$$

and multiplication by cutoff functions $\chi(t) \in C_0^\infty(R)$, $\chi(t) = 1$ near $t = 0$.

In Chapter 2 we study the continuity properties of pdo's on various weighted and unweighted L^p and Sobolev spaces on the half line R^+ .

A principal symbol for a class of pdo's on $L^p(R^+)$ is introduced in Chapter 3. The notion of ellipticity on $L^p(R^+)$ of a pdo (i.e., the existence of a parametrix modulo compact operators) depends on p as in [LP1]. We obtain that a pdo is elliptic on $L^p(R^+)$ iff its principal symbol never vanishes. An index theorem for pdo's on $L^2(R^+)$ was proved by Cordes and Herman [CH]. We give an index theorem for an elliptic system on $L^p(R^+)$ which generalizes [CH] and [LP1].

In Chapter 4 we define an algebra of pdo's of order 0 on $L^p(I)$, where I is a finite interval. We work modulo compact operators on $L^p(I)$ to define a principal symbol and prove that a pdo is a Fredholm operator on $L^p(I)$ iff its principal symbol never vanishes. For an elliptic system of pdo's on $L^p(I)$, we prove an index theorem which relates the index of the system to the topological degree of the principal symbol. In this case the principal symbol may be identified with a map from the unit circle S^1 into $GL(N, C)$; of course the degree and hence the index depend on the L^p space under consideration.

In Chapter 5, we apply the theory of pdo's to the study of single and double layer potentials for

Laplace's equation in a plane polygon. We study the Fredholm and index properties of the integral equations which arise for Dirichlet, Neumann, and oblique derivative and mixed problems.

Various classes of pdo's on R^+ , halfspaces, and manifolds with boundary have been considered by many authors. Boutet de Monvel [B] studied pdo's with the transmission property. Melrose [M] defined a class of operators on a half-space via oscillatory integrals. Nourrigat [N1] considered symbols $a(\cdot, z)$ holomorphic in a left half plane and gave applications to Fuchs type equations in [N2]. Shamir [S1, S2] observed how the spectrum of a pdo on R^+ depends on the L^p space on which it acts. Cordes and Herman [CH] and Cordes [C] used Banach algebra techniques to introduce the notion of a principal symbol. Eskin [E] used the Mellin transform representation of the Hilbert transform and Hardy kernels to study mixed boundary problems. Plamenevskii [P1], using the Mellin transform, assigned a pdo on the sphere S^{n-1} (actually a meromorphic function of a complex parameter λ) to an operator on R^n . He then reduced the study of an algebra of operators on R^n to the investigation of an algebra of meromorphic operator-valued functions on S^{n-1} . The authors [LP1] developed an algebra which is properly included in the algebra $Op - \Sigma_{1/p}$ studied in Chapter 3.

For the index properties of boundary value problems in polynomial domains, see the extensive work and bibliographies of Grisvard [G], Kondratiev [K], and Avantaggiati [Av1]. The L^p theory of boundary value problems was studied by Merigot [Mer]. Double layer potentials in sectors and bounded domains were considered by Fabes, Jodeit, and Lewis [JFL]. Applications of the present theory to the biharmonic operator in a polygon have been given by Diomeda and Lisena [DL].

1. The C^∞ theory of pdo's on R^+ .

We shall denote the open half line $(0, \infty)$ as R^+ and $[0, \infty)$ as $\overline{R^+}$. The basic differential operator used will be

$$\partial = -t \frac{d}{dt}$$

and ∂_t will denote $-t \frac{\partial}{\partial t}$. If a is a real number, the integer $[a]$ will be defined as

$$[a] = \text{greatest integer} < a.$$

The following function spaces will be used for the C^∞ theory of pdo's on R^+ .

DEFINITION 1.1. Let $-\infty < a < b < \infty$. By $\mathcal{F}_{a,b}$ we denote the class of functions $f \in C^\infty(R^+)$ such that:

- (1) For $j = 0, \dots, (-a)$, there are scalars f_{j0}

such that for every k and every $\delta > 0$,

$$(1.1) \quad \partial^k(f(t) - \sum_{j=0}^{(-a]} \frac{1}{j!} f_{j0} t^j) = o(t^{-a-\delta}), \quad t \rightarrow 0^+,$$

(2) For $j = 0, \dots, (b]$, there are scalars $f_{j\infty}$ such that for every k and every $\delta > 0$,

$$(1.2) \quad \partial^k(f(t) - \sum_{j=0}^{(b]} \frac{1}{j!} f_{j\infty} t^{-j}) = o(t^{-b+\delta}), \quad t \rightarrow \infty.$$

$\mathfrak{F}_{a,b}$ is a Frechet space with the seminorms

$$(1.3) \quad \begin{aligned} & |f_{j0}|, \quad j = 0, \dots, (-a]; \quad |f_{j\infty}|, \quad j = 0, \dots, (b]; \\ & \sup_{0 < t < 2} \left| t^{a+\delta} \partial^k \left(f(t) - \sum_{j=0}^{(-a]} \frac{1}{j!} f_{j0} t^j \right) \right|; \\ & \sup_{1 < t < \infty} \left| t^{b-\delta} \partial^k \left(f(t) - \sum_{j=0}^{(b]} \frac{1}{j!} f_{j\infty} t^{-j} \right) \right|. \end{aligned}$$

We shall abbreviate (1.1) as $f(t) \sim \sum \frac{1}{j!} f_{j0} t^j$, $t \rightarrow 0^+$, and shall write (1.2) as $f(t) \sim \sum \frac{1}{j!} f_{j\infty} t^{-j}$, $t \rightarrow \infty$. If $b > 0$, we denote by $\mathfrak{F}'_{a,b}$ the space of functions $f \in \mathfrak{F}_{a,b}$ such that $f \sim 0$, $t \rightarrow \infty$; i.e., $\partial^k f(t) = o(t^{-b+\delta})$, $t \rightarrow \infty$, for every $\delta > 0$. Similarly $\mathring{\mathfrak{F}}_{a,b}$ consists of these functions $f \in \mathfrak{F}_{a,b}$ such that $f(t) \sim 0$, $t \rightarrow 0^+$. We define

$$\mathfrak{F}''_{a,b} = \mathring{\mathfrak{F}}_{a,b} \cap \mathfrak{F}'_{a,b}.$$

The space $\check{\mathcal{F}}_{a,b}$ is stable under multiplication by $\log t$, and $C_0^\infty(\mathbb{R}^+)$ is dense in $\check{\mathcal{F}}_{a,b}$. We use the convention that $\check{\mathcal{F}}_{-\infty,b} = \bigcap_{a < b} \check{\mathcal{F}}_{a,b}$ and note that

$\check{\mathcal{F}}_{a,b} = \bigcap_{\delta > 0} \check{\mathcal{F}}_{a+\delta, b-\delta}$. If we define

$$(1.4) \quad \check{f}(t) = f\left(\frac{1}{t}\right),$$

then $f \in \check{\mathcal{F}}_{a,b}$ iff $f \in \check{\mathcal{F}}_{-b,-a}$.

The spaces $\check{\mathcal{M}}_{a,b}$ defined in Definition 1.2 will be the Mellin transforms of function in $\check{\mathcal{F}}_{a,b}$. By $S_{a,b}$ we denote the strip

$$(1.5) \quad S_{a,b} = \{z \in \mathbb{C} : a < \operatorname{Re} z < b\}$$

and let Z_- be the set of nonpositive integers.

DEFINITION 1.2. Let $0 < b$. The space $\check{\mathcal{M}}_{a,b}$ consists of functions $g(z)$ defined and meromorphic on the strip $S_{a,b}$ such that

1. g has poles only at $z = -j \in Z_- \cap S_{a,b}$ which are at most simple; denote the residue at $z = -j$ by $(1/j!)g_j$.
2. $g(z)$ is rapidly decreasing as $|\operatorname{Im} z| \rightarrow \infty$ in $S_{a,b}$; i.e., on every strip with base $(a', b') \subset (a, b)$

$$(1.6) \quad |g(z)| = O(|\operatorname{Im} z|^{-N})$$

for every N as $|\operatorname{Im} z| \rightarrow \infty$, $z \in S_{a', b'}$.

The following Lemma follows from the arguments of Avantaggiati [Av2], Nourrigat [N2], or [LP2].

LEMMA 1.1. Let $0 < b$. For $f \in \mathcal{F}_{a, b}'$ and
 $\max(a, 0) < \operatorname{Re} z < b$, define

$$(1.7) \quad \tilde{f}(z) = \int_0^\infty t^{z-1} f(t) dt.$$

Then $f(z)$ may be extended as a meromorphic function
in $\mathcal{M}_{a, b}'$ with residues at $z = -j \in \mathbb{Z}_- \cap S_{a, b}$ given
by $(1/j!)f_{j0}$.

Conversely, given $g(z) \in \mathcal{M}_{a, b}'$, for any γ such
that $\max(a, 0) < \gamma < b$, define

$$(1.8) \quad f(t) = \frac{1}{2\pi i} \int_{\operatorname{Re} z = \gamma} t^{-z} g(z) dz.$$

Then $f \in \mathcal{F}_{a, b}'$, $f \sim \sum_{j=0}^{(-a]} \frac{g_j}{j!} t^j$, $t \rightarrow 0^+$, and
 $\tilde{f}(z) = g(z)$, where f is defined by (1.7).

DEFINITION 1.3. For $-\infty \leq c < d \leq +\infty$ and m real
the class $\mathcal{S}_{c, d}^m$ consists of those functions $a(z)$
holomorphic in the strip $S_{c, d}$ such that for every
 $(c', d') \subset (c, d)$, for every k

$$(1.9) \quad \frac{d^k}{dz^k} a(z) = O(|\operatorname{Im} z|^{m-k})$$

as $|\operatorname{Im} z| \rightarrow \infty$, $z \in S_{c', d'}$.

$\mathcal{O}_{c, d}^m$ is a Frechet space with the seminorms

$$(1.10) \quad \sup_{z \in S_{c', d'}} |(1+|z|)^{k-m} \frac{d^k}{dz^k} a(z)|, (c', d') \subset (c, d).$$

For $-\infty \leq c < d \leq +\infty$ and $-\infty \leq a < b \leq +\infty$, $\mathcal{F}_{a, b}(\mathcal{O}_{c, d}^m)$ denotes the Frechet space $\mathcal{F}_{a, b}(\mathbb{R}^+; \mathcal{O}_{c, d}^m)$ of functions $a(t, z) \in \mathcal{F}_{a, b}$ as functions of t with values in $\mathcal{O}_{c, d}^m$.

We shall write $\mathcal{O}_{c, d}^{-\infty} = \bigcap_m \mathcal{O}_{c, d}^m$.

DEFINITION 1.4. For $-\infty \leq c < d \leq \infty$ and $m \in \mathbb{R}$, the symbol class $\Sigma_{c, d}^m$ is defined by

$$(1.11) \quad \Sigma_{c, d}^m = \bigcap_{(c', d') \subset (c, d)} \mathcal{F}_{c-c', d-d'}(\mathcal{O}_{c', d'}^m).$$

Also

$$(1.12) \quad \Sigma_{c, d}^m = \bigcap_{(c', d') \subset (c, d)} \mathcal{F}_{c-c', d-d'}(\mathcal{O}_{c', d'}^m).$$

REMARKS.

(1) If $\chi(t)$ is a cutoff function (i.e., $\chi(t) \in C_0^\infty(\mathbb{R})$, $\chi(t) = 1$ near $t = 0$), $\alpha > 0$, and $\operatorname{Re} z_0 = c$, then for any $a(z) \in \mathcal{O}_{c, d}^{-\infty}$, the function

(1.13) $a(t,z) = \chi(t)(\alpha t)^{z-z_0} a(z)$

is in $\Sigma_{c,d}^{0,-\infty}$.

2. If $a(t,z) \in \Sigma_{c,d}^m$ and $c < c' < \operatorname{Re} z < d$, we have the asymptotic expansion

(1.14) $a(t,z) \sim \sum_{j=0}^{(c'-c]} \frac{a_{j0}(z)}{j!} t^j, \quad t \rightarrow 0^+,$

where $a_{j0}(z) \in \mathcal{O}_{c+j,d}^m$.

We are now ready to define a class of pseudo-differential operators.

DEFINITION 1.5. Let $-\infty \leq c < d \leq +\infty$ and suppose that

(1.15) $J = (c,d) \cap (0,1) \neq \emptyset.$

If $a(t,z) \in \Sigma_{c,d}^m$ define the operator

$A = a(t,\partial) \in \operatorname{Op} - \Sigma_{c,d}^m$ as follows:

Choose $\gamma \in J$ and for $f \in \mathcal{S}'_{c,d}$, let

(1.16) $Af(t) = a(t,\partial)f(t) = \frac{1}{2\pi i} \int_{\operatorname{Re} z = \gamma} t^{-z} a(t,z) \tilde{f}(z) dz.$

The definition (1.16) is independent of $\gamma \in J$.

THEOREM 1.1. If $a(t,z) \in \Sigma_{c,d}^m$, then

$$a(t, \partial) : \mathcal{S}'_{c,d} \rightarrow \mathcal{S}'_{c,d}$$

is continuous; moreover, if

$$(1.17) \quad f(t) \sim \sum_{j=0}^{(-c]} \frac{1}{j!} f_{j0} t^j, \quad t \rightarrow 0^+$$

then

$$(1.18) \quad Af(t) \sim \sum_{k=0}^{(-c]} \frac{1}{k!} \left\{ \sum_{j=0}^k \binom{k}{j} a_{k-j,0}(-j) f_{j0} \right\} t^k, \quad t \rightarrow 0^+,$$

where $a_{k-j,0}(z)$ is given by (1.14). Hence

$$(1.19) \quad a(t, \partial) : \mathcal{S}_{c,d} \rightarrow \mathcal{S}_{c,d}$$

is continuous.

PROOF: Let $f \in \mathcal{S}'_{c,d}$ satisfy (1.17). For all sufficiently small $\delta > 0$, we shift the contour of integration in (1.16) to $\operatorname{Re} z = d - \delta$ and rewrite (1.16) as

$$(1.20) \quad Af(t) = \frac{1}{2\pi i} \int_{\operatorname{Re} z = d - \delta} t^{-z} a(t, z) \tilde{f}(z) dz.$$

This shows that $Af(t)$ is in $C^\infty(\mathbb{R}^+)$. Since for $\operatorname{Re} z = d - \delta$, $\partial_t^k (a(t, z) - a(\infty, z)) = O(t^{-\delta + \epsilon} (1 + |z|)^m)$, $t \rightarrow \infty$, we readily obtain from (1.20) that $\partial^k f(t) = O(t^{-d+\delta})$, $t \rightarrow \infty$.

To study the behavior of (1.16) near $t = 0$, we choose any small $\delta > 0$ such that $c + \delta$ is not a nonnegative integer. In (1.16) shift the contour of

integration to $\operatorname{Re} z = c + \delta$, and obtain that

$$Af(t) = \sum_{j=0}^{(-c]} \operatorname{Res}\{t^{-z}a(t,z)\tilde{f}(z); z = -j\} + I(t),$$

where $I(t) = \frac{1}{2\pi i} \int_{\operatorname{Re} z = c+\delta} t^{-z}a(t,z)\tilde{f}(z)dz$. It is easy to show that $\partial^k I(t) = O(t^{-c+\delta})$, $t \rightarrow 0^+$. We now study the residue

$$(1.22) \quad R_{-j} = \operatorname{Res}\{t^{-z}a(t,z)\tilde{f}(z); z = -j\}.$$

Choose a small $\epsilon > 0$ such that $(-c-\epsilon] = (-c]$. Since $a(t,z) \in \mathfrak{F}_{c+j+\epsilon, d+j-\epsilon}(\mathfrak{S}_{-j-\epsilon, -j+\epsilon}^m)$, then for $|z+j| \leq \epsilon/2$,

$$\partial_t^k \{a(t,z) - \sum_{k=0}^{(-c-j]} \frac{1}{k!} a_{k0}(z)t^k\} = O(t^{-c-k-2\epsilon}),$$

with $a_{k0}(z) \in \mathfrak{S}_{-j-\epsilon, -j+\epsilon}^m$.

Using the representation

$$R_{-j} = \frac{1}{2\pi i} \int_{|z+j| = \epsilon/2} t^{-z}a(t,z)\tilde{f}(z)dz$$

for (1.22) we have that $\partial_t^k R_{-j}$ is

$$\partial_t^k \left\{ \sum_{k=0}^{(-c-j]} \frac{1}{k!} a_{k0}(-j) \frac{1}{j!} f_{j0} t^{k+j} \right\} + O(t^{-c-j-2\epsilon}).$$

This proves that $Af \in \mathfrak{F}_{c,d}'$ and formulas (1.18) and (1.19). q.e.d.

We now study the composition of two operators.

THEOREM 1.2. Let $a(t, z) \in \Sigma_{c, d}^m$ and $b(t, z) \in \Sigma_{c, d}^{m'}$.
Then there is a symbol $c(t, z) \in \Sigma_{c, d}^{m+m'}$ such that

1. For every N

$$(1.23) \quad c(t, z) - \sum_{k=0}^N \frac{1}{k!} \frac{\partial^k}{\partial z^k} b(t, z) (\partial_t^k a)(t, z) \in \Sigma_{c, d}^{m+m'-N-1}.$$

2. $b(t, \partial) \circ a(t, \partial) = c(t, \partial)$ (as operators on
 $\mathcal{S}_{c, d}'$).

PROOF: Fixing a cutoff function $\chi(t)$, we split the symbol of $a(t, z)$ as

$$(1.24) \quad a(t, z) = \chi(t)a(t, z) + (1-\chi(t))a(t, z) \\
= a_0(t, z) + a_\infty(t, z).$$

Consider the contribution of $b(t, \partial) \circ a_\infty(t, \partial)$. Since $a_\infty(t, z) \in \Sigma_{c, d}^m$, for $\operatorname{Re} z - d < \operatorname{Re} w < 0$ we define

$$(1.25) \quad m_{a_\infty}(w, z) = \int_0^\infty t^{w-1} a_\infty(t, z) dt.$$

Since $m_{a_\infty}(w, z)$ is the Mellin transform of the function $t \rightarrow a_\infty\left(\frac{1}{t}, z\right)$ evaluated at the point $(-w)$, and $t \rightarrow a_\infty\left(\frac{1}{t}, z\right)$ is a function in $\mathcal{S}_{\operatorname{Re} z - d, \operatorname{Re} z - c}'$, the function $m_{a_\infty}(w, z)$ is a meromorphic function of w in $S_z = S_{c - \operatorname{Re} z, \operatorname{Re} z}$ with simple poles at $\{0, 1, 2, \dots\} \cap S_z$ with residues given by $-a_{j_\infty}(z)/j!$; $j = 0, \dots, (d - \operatorname{Re} z]$; and we have the inversion formula

$$(1.26) \quad a_\infty(t, z) = \frac{1}{2\pi i} \int_{\operatorname{Re} w = \gamma'} t^{-w} m_{a_\infty}(w, z) dw,$$

$$\operatorname{Re} z - d < \gamma' < 0.$$

Fix $f \in \mathcal{F}_{c,d}$ and let $g(t) = a_{\infty}(t, \partial)f(t)$. If $\max(0, c) < \gamma < d$ and $c < \operatorname{Re} w = \gamma' < \gamma$, then

$$\begin{aligned} \tilde{g}(w) &= \int_0^{\infty} t^{w-1} \left\{ \frac{1}{2\pi i} \int_{\operatorname{Re} z = \gamma} t^{-z} a_{\infty}(t, z) \tilde{f}(z) dz \right\} dw \\ (1.27) \quad &= \frac{1}{2\pi i} \int_{\operatorname{Re} z = \gamma} m a_{\infty}(w-z, z) \tilde{f}(z) dz. \end{aligned}$$

The use of Fubini's Theorem in (1.27) is justified since $c - \operatorname{Re} z < \operatorname{Re} w - \operatorname{Re} z < 0$. Since $g(t) \in \mathcal{F}_{c,d}$ $\tilde{g}(w)$ has no pole in $S_{c,d}$; hence for any γ' , $c < \gamma' < d$, we have

$$(1.28) \quad b(t, \partial)g(t) = \frac{1}{2\pi i} \int_{\operatorname{Re} w = \gamma'} t^{-w} b(t, w) \tilde{g}(w) dw.$$

Using (1.27) and Peetre's inequality, $(1+|w-z|)^{-N} \leq C(1+|w|)^{-N}(1+|z|)^N$, we apply Fubini's Theorem to obtain

$$(1.29) \quad b(t, \partial) \circ a_{\infty}(t, \partial)f(t) = \frac{1}{2\pi i} \int_{\operatorname{Re} z = \gamma} t^{-z} c_{\infty}(t, z) \tilde{f}(z) dz,$$

where

$$(1.30) \quad c_{\infty}(t, z) = \frac{1}{2\pi i} \int_{\operatorname{Re} w = \gamma'} t^{z-w} b(t, w) m a_{\infty}(w-z, z) dw.$$

In (1.30) choose any $\gamma' \in (c, \operatorname{Re} z)$. We make the change of variables $w-z = v$, $\operatorname{Re} v = \gamma_1 \in (c-\operatorname{Re} z, 0)$,

and apply Taylor's formula with integral remainder and the inversion formula (1.26), so that

$$(1.31) \quad c_{\infty}(t, z) = \sum_{k=0}^N \frac{1}{k!} \frac{\partial^k}{\partial z^k} b(t, z) (\partial_t^k a_{\infty})(t, z) + R_{\infty}^N(t, z),$$

with

$$(1.32) \quad R_{\infty}^N(t, z)$$

$$= \frac{1}{N!} \int_0^1 (1-s)^N \frac{1}{2\pi i} \int_{\operatorname{Re} v = \gamma_1} t^{-v} \frac{\partial^{N+1}}{\partial z^{N+1}} b(t, z+sv) v^N m_{\infty}(v, z) dv ds$$

Fix s , $0 \leq s \leq 1$, and let

$$(1.33) \quad c_s(t, z) = \frac{1}{2\pi i} \int_{\operatorname{Re} v = \gamma_1} t^{-v} \frac{\partial^{N+1}}{\partial z^{N+1}} b(t, z+sv) v^N m_{\infty}(v, z) dv$$

where $\gamma_1 \in (c - \operatorname{Re} z, 0)$. The function $c_s(t, z)$, originally defined by (1.33) for $\operatorname{Re} z > 0$ may be prolonged as a holomorphic function of z in the entire strip $S_{c,d}$ by using formula (1.33) with small γ_1 ; shifting the contour of integration to $\operatorname{Re} v = c - \operatorname{Re} z + \delta$ or $d - \operatorname{Re} z - \delta$ and taking into account the poles $v^N m_{\infty}(v, a)$ at $v = 1, \dots, (d - \operatorname{Re} z]$, we may show that

$$c_s(t, z) \in \mathcal{E}_{c-c'}^0(\mathcal{O}_{c',d'}^{m+m'-N-1})$$

if $(c', d') \subset (c, d)$. Hence $R_{\infty}^N(t, z) \in \Sigma_{c,d}^{m+m'-N-1}$.

A similar, but simpler argument shows that if $a_0(t, z)$ is defined as in (1.24) then

$$(1.34) \quad b(t, \partial) \circ a_0(t, \partial) f(t) = c_0(t, \partial) f(t),$$

where for $\max(0, c) < \operatorname{Re} z < d$

$$(1.35) \quad c_0(t, z) = \sum_{k=0}^N \frac{1}{k!} \frac{\partial^k}{\partial z^k} b(t, z) (\partial_t^k a_0)(t, z) + R_0^N(t, z).$$

and

$$(1.36) \quad R_0^N(t, z)$$

$$= \frac{1}{N!} \int_0^1 (1-s)^N \frac{1}{2\pi i} \int_{\operatorname{Re} v = \gamma_1} t^{-v} \frac{\partial^{N+1}}{\partial z^{N+1}} b(t, z+sv) v^N a_0^\sim(v, z) dv ds,$$

where $\gamma_1 \in (0, d - \operatorname{Re} z)$ and $a_0^\sim(v, z) = \int_0^\infty t^{v-1} a_0(t, z) dt$.

Again it follows that $R_0^N(t, z) \in \Sigma_{c,d}^{m+m'-N-1}$.

q.e.d.

We consider the transpose and the adjoint of an operator $A = a(t, \partial) \in \operatorname{Op} - \Sigma_{c,d}^m$. The transpose of the operator tA , and the adjoint of the operator, A^* , are defined by the relations:

$$(1.37) \quad \begin{aligned} &\text{For } f \in \mathcal{F}_{c,d}, \varphi \in C_0^\infty(\mathbb{R}^+), \\ &\int_0^\infty Af(t)\varphi(t)dt = \int_0^\infty f(t)({}^tA\varphi)(t)dt, \\ &\int_0^\infty Af(t)\overline{\varphi(t)}dt = \int_0^\infty f(t)\overline{A^*\varphi(t)}dt. \end{aligned}$$

THEOREM 1.3. Suppose that $(c, d) \cap (0, 1) \neq \emptyset$, and
that $a(t, z) \in \operatorname{Op} - \Sigma_{c,d}^m$. Then there are symbols
 ${}^ta(t, z)$ and $a^*(t, z) \in \Sigma_{1-d, 1-c}^m$, such that

$$(1.38) \quad {}^t a(t, z) - \sum_{k=0}^N \frac{(-1)^k}{k!} \left(\frac{\partial}{\partial w} \right)^k \left(\partial_t^k a \right)(t, w) \Big|_{w=1-z} \in \Sigma_{1-d, 1-c}^{m-N-1}$$

$$(1.39) \quad a^*(t, z) - \sum_{k=0}^N \frac{(-1)^k}{k!} \left\{ \left(\frac{\partial}{\partial w} \right)^k \left(\partial_t^k a \right)(t, w) \Big|_{w=1-\bar{z}} \right\}^* \in \Sigma_{1-d, 1-c}^{m-N-1}$$

and if $A = a(t, \partial) \in \text{Op} - \Sigma_{c, d}^m$, then

$$(1.40) \quad {}^t A = {}^t a(t, \partial) \in \text{Op} - \Sigma_{1-d, 1-c}^m,$$

$$(1.41) \quad A^* = a^*(t, \partial) \in \text{Op} - \Sigma_{1-d, 1-c}^m.$$

We sketch the proof of Theorem 5.2. Split $a(t, z)$ as in (1.24). If $m \leq -2$ and $\varphi \in C_0^\infty(\mathbb{R}^+)$ show that with $\max(0, c) < \gamma < 1$,

$$(1.42) \quad {}^t A_0 \varphi(s) = \frac{1}{2\pi i} \int_{\text{Re } w = 1-\gamma} s^{-w} \int_0^\infty t^{w-1} a_0(t, 1-w) \varphi(t) dt dw.$$

The proof then proceeds in the spirit of the proof of Theorem 5.1. See also Theorem 4 of [LP2]. In the case $m > -2$, choose $\lambda_0 > d$ and write $a(t, z) = a_1(t, z)(z - \lambda_0)^{m+2}$ with $a_1(t, z) \in \Sigma_{c, d}^{-2}$. Then $a(t, \partial) = a_1(t, \partial)(\partial - \lambda_0)^{m+2}$. The transpose of the differential operator $(\partial - \lambda_0)^{m+2}$ may be calculated directly and ${}^t a(t, \partial)$ is handled by the first part of the proof. We leave the details to the reader.

2. L^p estimates for pseudodifferential operators on \mathbb{R}^+ .

Let $1 < p < \infty$. We shall prove estimates for the operators considered in Chapter 1 on $L^p(\mathbb{R}^+)$ and on

various weighted and Sobolev spaces on R^+ . Let

$$\|f\|_p = \|f; L^p(R^+)\| = \left(\int_0^\infty |f(t)|^p dt \right)^{1/p}.$$

THEOREM 2.1. Suppose that $1 < p < \infty$ and that
 $c < 1/p < d$. If $a(t, z) \in \Sigma_{c,d}^0$ and

$$(2.1) \quad \begin{aligned} Af(t) &= a(t, \partial)f(t) \\ &= \frac{1}{2\pi i} \int_{\operatorname{Re} z = 1/p} t^{-z} a(t, z) \tilde{f}(t) dt, \end{aligned}$$

for $f \in \mathcal{S}'_{c,d}$, then there is a constant $C = C(a, p)$
such that

$$(2.2) \quad \|Af\|_p \leq C \|f\|_p.$$

PROOF: For $f \in \mathcal{S}'_{c,d}$ define

$$(2.3) \quad T_p f(x) = F_p(x) = e^{-(1/p)x} f(e^{-x}) \in \mathcal{S}(R).$$

Then $\|F_p; L^p(R)\| = \|f; L^p(R^+)\|$ and

$$T_p Af(x) = \frac{1}{2\pi} \int_{-\infty}^{\infty} e^{ix\xi} a(e^{-x}, 1/p + i\xi) \tilde{F}_p(\xi) d\xi.$$

The symbol $b_p(x, \xi) = a(e^{-x}, 1/p + i\xi)$ satisfies

$$\frac{\partial^{j+k}}{\partial x^j \partial \xi^k} b_p(x, \xi) = O((1 + |\xi|)^{-k})$$

and hence $T_p A T_p^{-1}$ is a classical Calderon-Zygmund

operator of class $S_{1,0}^0$, and hence

$$(2.4) \quad \|T_p A T_p^{-1} F_p; L^p(R)\| \leq C_p \|F_p; L^p(R)\|.$$

Cf. Meyer [M] or Coifman and Meyer [CM]. Inequality

(2.3) now follows from (2.4).

q.e.d.

Before studying the continuity of operator $a(t, \partial) \in Op - \Sigma_{c,d}^m$ on Sobolev spaces we remark that if $a(t, z) \in \Sigma_{c,d}^m$, then $a(t, z-1) \in \Sigma_{c+1,d+1}^m$ and that if $c+1 < d$, $(\partial/\partial t)a(t, z) \in \Sigma_{c+1,d}^m$.

For $k \in \mathbb{Z}_+ = \{0, 1, 2, \dots\}$, we let $W^{p,k}$ be the usual Sobolev space on R^+ :

$$W^{p,k} = \{f \in L^p(R^+); f^{(j)} \in L^p(R^+), j = 0, \dots, k\};$$

$$W^{p,k} = \{f \in W^{p,k}; f^{(j)}(0) = 0, j = 0, \dots, k-1\}.$$

If $k \in \mathbb{Z}_- = \{0, -1, -2, \dots\}$, then $W^{p,k}$ is the dual of $W^{q,-k}$, $1/p + 1/q = 1$, under the bilinear form

$$\langle f, \varphi \rangle = \int_0^\infty f(t) \varphi(t), \quad f \in C^\infty(\overline{R^+}), \quad \varphi \in C_0^\infty(R^+).$$

THEOREM 2.2. Suppose that $a(t, z) \in \Sigma_{c,d}^0$ and that $A = a(t, \partial)$ is defined by (2.1). Then

1. If $k \geq 0$ and $c+k < 1/p < d$,

$$(2.5) \quad a(t, \partial): W^{p,k} \rightarrow W^{p,k}$$

is continuous; moreover,

(2.6) $a(t,\partial): \dot{W}^{p,k} \rightarrow \dot{W}^{p,k}$

is continuous.

2. If $j \geq 0$ and $c < 1/p < d-j$, then

(2.7) $a(t,\partial): W^{p,-j} \rightarrow W^{p,-j}$

is continuous.

PROOF: If $c+1 < 1/p < d$ and $f \in \mathcal{F}'_{c,d}$, then

(2.8)
$$\begin{aligned} \frac{d}{dt} a(t,\partial)f(t) &= \frac{1}{2\pi i} \int_{\operatorname{Re} z = 1/p} t^{-z-1} a(t,z) (-z\tilde{f}(z)) dz \\ &= \frac{1}{2\pi i} \int_{\operatorname{Re} z = 1/p} t^{-z} (\partial/\partial t) a(t,z) \tilde{f}(z) dz. \end{aligned}$$

The first term on the r.h.s. of (2.8) is

$$\begin{aligned} &\frac{1}{2\pi i} \int_{\operatorname{Re} z = 1/p+1} t^{-z} a(t,z-1) [f']^{\sim}(t) dt \\ &= \frac{1}{2\pi i} \int_{\operatorname{Re} z = 1/p} t^{-z} a(t,z-1) [f']^{\sim}(z) dz. \end{aligned}$$

The

second term on the r.h.s. of (2.8) is $(\partial/\partial t) a(t,\partial)f$.
Hence by Theorem 2.1,

(2.9) $\|\frac{d}{dt} a(t,\partial)f\|_p \leq C\|f'\|_p + C\|f\|_p.$

The continuity of (2.5) now follows by induction on k ; (2.6) follows since $a(t,\partial): \mathcal{F}'_{c,d} \rightarrow \mathcal{F}'_{c,d}$.

To prove 2., fix $f \in C_0^\infty(\mathbb{R})$ and $\varphi \in C_0^\infty(\mathbb{R}^+)$. By Theorem 1.3,

$$\langle Af, \varphi \rangle = \langle f, {}^tA\varphi \rangle,$$

with ${}^tA = {}^t a(t, \partial)$, where ${}^t a(t, z) \in \Sigma_{1-d, 1-c}^0$. If $1/p + 1/q = 1$ and $1-d+j < 1/q < 1-c$, i.e., $c < 1/p < d-j$, then ${}^tA: W^{q,j} \rightarrow W^{q,j}$ is continuous by part 1 of the Theorem. Part 2 follows.

q.e.d.

To study operator $a(t, \partial) \in \text{Op} - \Sigma_{c,d}^m$ with $m \neq 0$, we introduce weighted Sobolev spaces following the ideas and notation of Nourrigat [N2].

DEFINITION 2.1. For $\lambda \in \mathbb{Z}_+$ define

$$(2.10) \quad W_\lambda^{p,k} = \{f \in W^{p,k}; \partial^j f \in W^{p,k}, j = 0, \dots, \lambda\},$$

and

$$(2.11) \quad \dot{W}_\lambda^{p,k} = \{f \in W^{p,k}; \partial^j f \in \dot{W}^{p,k}, j = 0, \dots, \lambda\}.$$

To define $W_\lambda^{p,k}$ with $\lambda \in \mathbb{Z}_-$, we introduce an invertible pdo $\Lambda = \lambda(\partial)$ of order 1. Let d_0 be a large positive number and define

$$(2.11) \quad \lambda(z) = z - d_0 \in \Sigma_{-\infty, \infty}^{-1},$$

$$\lambda^{-1}(z) = (z - d_0)^{-1} \in \Sigma_{-\infty, d_0}^{-1}$$

and let $\Lambda = \lambda(\partial)$, $\Lambda^{-1} = \lambda^{-1}(\partial)$. If $f \in \mathcal{S}'_{c,d}$, $d \leq d_0$

$$(2.12) \quad \Lambda^{-1}f(t) = \int_0^t \left(\frac{t}{s}\right)^{-d_0} f(s) \frac{ds}{s}.$$

and

$$(2.13) \quad \Lambda\Lambda^{-1}f = \Lambda^{-1}\Lambda f = f.$$

LEMMA 2.1. If $l \in \mathbb{Z}_+$, then $f \in W_l^{p,k}$ iff $\Lambda^l f \in W^{p,k}$; moreover, $f \in W_l^{p,k}$ iff $\Lambda^l f \in W^{p,k}$. Also

$$\|f; W_l^{p,k}\| \sim \|\Lambda^l f; W^{p,k}\|.$$

PROOF: If $f \in W_l^{p,k}$, then $\|\Lambda^l f; W^{p,k}\| \leq C \sum_{j=0}^l \|\partial^j f; W^{p,k}\|$
 $= C\|f; W_l^{p,k}\|.$

If $f \in C_0^\infty(\mathbb{R})$, then for $j = 0, \dots, l$, $\partial^j f = a_j(\partial)\Lambda^l f$
 where $a_j(z) = z^j(z - d_0)^{-l} \in \Sigma_{-\infty, d_0}^{j-l}$. By Theorem 2.2,
 $\|\partial^j f; W^{p,k}\| \leq C\|\Lambda^l f; W^{p,k}\|.$

q.e.d.

DEFINITION 2.2. For $l \in \mathbb{Z}_-$, a distribution
 $f \in \mathcal{S}'(\mathbb{R}^+)$ is in $W_l^{p,k}$ (respectively $\dot{W}_l^{p,k}$) iff
 $f = \Lambda^{-l}g$, $g \in W^{p,k}$ (respectively $\dot{W}^{p,k}$). The norm
on $W_l^{p,k}$ is

$$(2.15) \quad \|f; W_{\ell}^{p,k}\| = \|\Lambda^{\ell} f; W^{p,k}\|.$$

THEOREM 2.3. Let $m \in \mathbb{Z}$ and $d \leq d_0$. Let $a(t, z) \in \Sigma_{c,d}^m$ and define $A = a(t, \partial)$. Then

1. If $k \in \mathbb{Z}_+$ and $c+k \leq 1/p < d$, then
for all $\ell \in \mathbb{Z}$,

$$(2.16) \quad A : W_{\ell}^{p,k} \rightarrow W_{\ell-m}^{p,k}$$

$$A : \dot{W}_{\ell}^{p,k} \rightarrow \dot{W}_{\ell-m}^{p,k}$$

is continuous.

2. If $j \geq 0$ and $c < 1/p < d-j$, then

$$(2.17) \quad A : W_{\ell}^{p,-j} \rightarrow W_{\ell}^{p,-j}$$

is continuous.

PROOF: If $f = \Lambda^{-\ell} g$, $g \in W^{p,k}$, then

$\Lambda^{\ell-m} a(t, \partial) f = \Lambda^{\ell-m} a(t, \partial) \Lambda^{-\ell} g$ and the operator

$$b(t, \partial) = \Lambda^{\ell-m} a(t, \partial) \Lambda^{-\ell}$$

is in $Op - \Sigma_{c,d}^0$ by the Symbolic Calculus Theorem

1.2. Hence by Theorem 2.2, $\|A f; W_{\ell-m}^{p,k}\| \sim \|\Lambda^{\ell-m} A f; W^{p,k}\|$

$$\leq C \|b(t, \partial) g; W^{p,k}\| \leq C \|g; W^{p,k}\| \sim \|f; W_{\ell}^{p,k}\|.$$

q.e.d.

3. An algebra of pseudodifferential operators on $L^p(\mathbb{R}^+)$.

Let $a(t,\partial) \in \text{Op} - \Sigma_{c,d}^0$ with $c < 1/p < d$. We wish to give conditions on the symbol $a(t,z)$ which determine whether the operator $a(t,\partial)$ is a Fredholm operator on $L^p(\mathbb{R}^+)$ and to link the index of a Fredholm operator to topological properties of the symbol. It may happen that the operator $a(t,\partial)$ is Fredholm on $L^p(\mathbb{R}^+)$ for some choices of $1/p \in (c,d)$ and not for others; the index of the operator (on $L^p(\mathbb{R}^+)$) will change as we cross a non-Fredholm value of $1/p \in (c,d)$.

Before defining the notion of principal symbol, we introduce the functions in $\mathcal{O}_{0,1}^0$;

(3.1)

$$\begin{aligned} h(z) &= -\cot \pi z, \\ \theta(z) &= (1 - e^{2\pi iz})^{-1}, \\ 1 - \theta(z) &= -e^{2\pi iz}(1 - e^{2\pi iz})^{-1}. \end{aligned}$$

Note that $\theta(1/p + i\infty) = 1$, $\theta(1/p - i\infty) = 0$, and that for $j \geq 1$, $(d/dz)^j \theta(z) \in \mathcal{O}_{0,1}^{-\infty}$. The corresponding operators in $\text{Op} - \Sigma_{0,1}^0$ have the representations:

(3.2)

$$\begin{aligned} Hf(t) &= h(\partial)f(t) = \text{p.v.} \frac{1}{\pi} \int_0^\infty \frac{f(s)}{t-s} ds, \\ \theta(\partial) &= \frac{1}{2} (I - iH), \\ I - \theta(\partial) &= \frac{1}{2} (I + iH); \end{aligned}$$

H is the Hilbert transform on $L^p(\mathbb{R}^+)$: Note that if \mathcal{H} is the Hilbert transform on $L^p(\mathbb{R}), (1/2)(I \pm i\mathcal{H})$ form a basis for singular integral operators on \mathbb{R} .

DEFINITION 3.1. A symbol $a(t, z)$ is in the class $\Sigma_{1/p}$ iff for some c, d with $0 \leq c < 1/p < d \leq 1$,

$$1. \quad a(t, z) \in \Sigma_{c,d}^0,$$

$$2. \quad \text{There are functions } a_+(t), a_-(t) \in \Sigma_{c-d, d-c} \text{ such that}$$

$$(3.3) \quad a(t, z) - a_+(t)\theta(z) - a_-(t)(1 - \theta(z)) \in \Sigma_{c,d}^{-1}.$$

If $a(t, z) \in \Sigma_{1/p}$, the functions $a_+(t), a_-(t)$ in (3.3) are uniquely determined by the relations

$$(3.4) \quad a_+(t) = a(t, 1/p + i\infty), \quad a_-(t) = a(t, 1/p - i\infty).$$

The function $a(t, z)$ is a continuous function defined on the compact rectangle:

$$(3.5) \quad R_{1/p} = \{(t, z) : 0 \leq t \leq \infty, \quad z = 1/p + i\xi, \quad -\infty \leq \xi \leq \infty\}.$$

DEFINITION 3.2. If $a(t, z) \in \Sigma_{1/p}$ and

$$(3.6) \quad \begin{aligned} Af(t) &= a(t, \partial)f(t) \\ &= \frac{1}{2\pi i} \int_{\operatorname{Re} z = 1/p} t^{-z} a(t, z) \tilde{f}(z) dz, \end{aligned}$$

for $f \in C_0^\infty(\mathbb{R})$, then the principal symbol, $\sigma(A)$, of the operator A is the function $a(t, z)$ restricted to the boundary of the compact rectangle $R_{1/p}$.

REMARKS

1. $\sigma(A) = 0$ iff $a(t, z) \in \mathcal{F}_{c,d}^{-1}$ for some c, d .
2. If for some $\delta > 0$ we are given functions $a_+(t), a_-(t) \in \mathcal{F}_{-\delta, \delta}$, and $a_0(z), a_\infty(z)$ such that the functions

$$(3.7) \quad \begin{aligned} b_0(z) &= a_0(z) - a_+(0)\theta(z) - a_-(0)(1 - \theta(z)), \\ b_\infty(z) &= a_\infty(z) - a_+(\infty)\theta(z) - a_-(\infty)(1 - \theta(z)), \end{aligned}$$

are in $\mathcal{O}_{1/p - \delta, 1/p + \delta}^{-1}$, we select a cutoff function $\chi(t) \in C_0^\infty(\mathbb{R})$, $\chi(t) = 1$ near $t = 0$, and define

$$(3.8) \quad \begin{aligned} a(t, z) &= \chi(t)b_0(z) + (1 - \chi(t))b_\infty(z) + a_+(t)\theta(z) \\ &\quad + a_-(t)(1 - \theta(z)). \end{aligned}$$

Then $a(t, z) \in \Sigma_{1/p}$ and

$$(3.9) \quad \begin{cases} a(0, z) = a_0(z), \quad a(\infty, z) = a_\infty(z), \\ a(t, 1/p + i\infty) = a_+(t), \quad a(t, 1/p - i\infty) = a_-(t). \end{cases}$$

3. That the principal symbol, $\sigma(A)$, of an operator $A = a(t, \partial)$ is uniquely determined by the operator A is a consequence of the following observations. Fix

$f \in C_0^\infty(\mathbb{R}^+)$. Then for $\lambda \in \mathbb{R}$,

$$(3.10) \quad t^{i\lambda} A(t^{-i\lambda} f)(t) = \\ = \frac{1}{2\pi i} \int_{\operatorname{Re} z = 1/p} t^{-z} a(t, z + i\lambda) \tilde{f}(z) dz.$$

As $\lambda \rightarrow \pm\infty$, $t^{i\lambda} A(t^{-i\lambda} f)$ converges in $L^p(\mathbb{R}^+)$ to $a_\pm(t)f(t)$. Next, for $\lambda > 0$, let $T_\lambda f(t) = \lambda^{-1/p} f(t/\lambda)$. Then $\|T_\lambda f\|_p = \|f\|_p$ and

$$(3.11) \quad (T_{1/\lambda} A T_\lambda) f(t) = \frac{1}{2\pi i} \int_{\operatorname{Re} z = 1/p} t^{-z} a(\lambda t, z) \tilde{f}(z) dz.$$

Hence in $L^p(\mathbb{R}^+)$,

$$(3.12) \quad \lim_{\epsilon \rightarrow 0} (T_{1/\epsilon} A T_\epsilon f)(t) = \frac{1}{2\pi i} \int_{\operatorname{Re} z = 1/p} t^{-z} a(0, z) \tilde{f}(z) dz,$$

$$\lim_{\lambda \rightarrow \infty} (T_{1/\lambda} A T_\lambda f)(t) = \frac{1}{2\pi i} \int_{\operatorname{Re} z = 1/p} t^{-z} a(\infty, z) \tilde{f}(z) dz.$$

The following result is fundamental in studying the Fredholm properties on $L^p(\mathbb{R}^+)$ operators defined by (3.6).

THEOREM 3.1. If $a(t, z) \in \Sigma_{1/p}$, then

$$A = a(t, \partial) : L^p \rightarrow L^p$$

is a compact operator iff the principal symbol, $\sigma(A)$, is identically 0; i.e., $a(t, z) \in \Sigma_{c,d}^{-1}$ for some c, d , $c < 1/p < d$.

PROOF: We first show that if $a(t, z) \in \Sigma_{c,d}^{-1}$ then $a(t, \partial)$ is compact on $L^p(\mathbb{R}^+)$. If $a(t, z) \in \Sigma_{c,d}^{-1}$, then $f \rightarrow -t \frac{\partial}{\partial t} A f$ is bounded on $L^p(\mathbb{R}^+)$ by Theorems 1.2 and 2.1. Hence if $a(t, z) = 0$ for t outside $[\epsilon, T]$, $0 < \epsilon < T < \infty$, A maps L^p into $W^{p,1}(\epsilon, T)$, which is compactly embedded in L^p . If $a(t, z) \in \Sigma_{c,d}^{\alpha-1}$, and $\epsilon > 0$, let $\chi(t)$ be a cutoff function, and define

$$a_\epsilon(t, z) = (1 - \chi(t/\epsilon))\chi(\epsilon t)a(t, z);$$

$a_\epsilon(t, z)$ converges to $a(t, z)$ in $\Sigma_{c,d}^{\alpha-1}$ as $\epsilon \rightarrow 0^+$. The corresponding operators $a_\epsilon(t, \partial)$ are compact on L^p and converge in operator norm to $a(t, \partial)$.

Conversely, suppose $A = a(t, \partial) : L^p \rightarrow L^p$ is compact. We use the constructions of Remark 3. following Definition 3.2. Fix $f \in C_0^\infty(\mathbb{R}^+)$. Then as $\lambda \rightarrow \pm\infty$, $t^{-i\lambda}f(t)$ converges to 0 weakly in L^p , by the Riemann-Lebesgue Lemma, so that $t^{i\lambda}A(t^{-i\lambda}f)$ converges to 0 strongly in L^p . Hence $a_+(t) = a_-(t) = 0$. If $T_\lambda f(t) = \lambda^{-1/p}f(t/\lambda)$, as $\lambda \rightarrow 0^+$ or $\lambda \rightarrow \infty$, $T_\lambda f$ converges to 0 weakly in L^p , and $A(T_\lambda f)$ converges to 0 in L^p . Each of the operators in (3.11) is 0 so that $a(0, z) = a(\infty, z) = 0$.

q.e.d.

DEFINITION 3.4. Let $a(t, z) \in \Sigma_{1/p}$ and let $a(t, \partial) \in \text{Op} - \Sigma_{1/p}$ be defined by (3.6). Then A

is said to be elliptic on $L^p(\mathbb{R}^+)$ iff its principal symbol, $\sigma(A)$, satisfies

$$(3.13) \quad \sigma(A)(t, z) \neq 0, \quad (t, z) \in \partial R_{1/p}.$$

If $a(t, z)$ is the symbol of an elliptic operator on $L^p(\mathbb{R}^+)$, then for $0 \leq t \leq \infty$,

$$(3.14) \quad \begin{aligned} a_+(t) &= a(t, 1/p + i\infty) \neq 0, \\ a_-(t) &= a(t, 1/p - i\infty) \neq 0, \end{aligned}$$

which are conditions independent of p , and reflect the fact that $a(t, \partial)$, considered as a classical pdo of order 0 on the noncompact manifold \mathbb{R}^+ is elliptic. If $a(t, z) \in \Sigma_{1/p}$, the conditions

$$(3.15) \quad a(0, z) \neq 0, \quad a(\infty, z) \neq 0, \quad \operatorname{Re} z = 1/p$$

are then satisfied for all p such that $1/p$ lies outside a discrete set in (c, d) .

We now relate "ellipticity on $L^p(\mathbb{R}^+)$ " to the Fredholm properties of the operator $A = a(t, \partial)$.

THEOREM 3.2. Let $A = a(t, \partial) \in \mathcal{O}p - \Sigma_{1/p}$. The following conditions are equivalent:

1. $a(t, \partial)$ is elliptic on $L^p(\mathbb{R}^+)$.
2. For some c, d , $0 \leq c < 1/p < d \leq 1$, there is a symbol $b(t, z) \in \Sigma_{c,d}^0$ such that

$$a(t, \partial)b(t, \partial) - I, b(t, \partial)a(t, \partial) - I \in \text{Op} - \Sigma_{c,d}^{-1}.$$

3. There is a bounded operator B on $L^p(\mathbb{R}^+)$ such that $AB - I$ and $BA - I$ are compact operators on $L^p(\mathbb{R}^+)$.

4. For $f \in C_0^\infty(\mathbb{R}^+)$, there is an a priori estimate

$$(3.16) \quad \|f\|_p \leq C\|Af\|_p + \|Kf\|_p,$$

where K is a compact operator on $L^p(\mathbb{R}^+)$.

PROOF: 1 implies 2: Choose $c < 1/p < d$ such that $a(0, z)$ and $a(\infty, z)$ have no zeroes in the strip $S_{c,d}$. Let $b(t, z) \in \Sigma_{c,d}^0$ be such that for $c < 1/q < d$, $b(t, z)a(t, z) = a(t, z)b(t, z) = 1$ on the boundary of $R_{1/q}$; $b(t, \partial)$ could be constructed as in Remark 2 following Definition 3.2. Condition 2 now follows from Theorem 1.2.

2 implies 3: This follows from Theorem 3.1.

3 implies 4: This is obvious.

4 implies 1: We use the constructions of Remark 3 following Definition 3.2. Fix $f \in C_0^\infty(\mathbb{R}^+)$. Then if K is compact on L^p , $\|t^{i\lambda}K(t^{-i\lambda}f)\|_p \rightarrow 0$ as $\lambda \rightarrow \pm\infty$. Inequality (3.16) implies that

$$\|f\|_p \leq C\|t^{i\lambda}A(t^{-i\lambda}f)\|_p + \|t^{i\lambda}K(t^{-i\lambda}f)\|_p$$

and letting $\lambda \rightarrow \pm\infty$, we obtain the a priori inequality

$$(3.17) \quad \|f\|_p \leq C \|a_{\pm}(t)f\|_p$$

which implies that $|a_{\pm}(t)| \geq \frac{1}{C}$. If $T_{\lambda}f(t) = \lambda^{-1/p}f(t/\lambda)$, then

$$(3.18) \quad \|f\|_p \leq C \|T_{1/\lambda} A(T_{\lambda}f)\|_p + \|T_{1/\lambda} K T_{\lambda} f\|_p.$$

Letting $\lambda \rightarrow 0$ or $\lambda \rightarrow \infty$ in (3.18) and applying (3.12) we obtain

$$(3.19) \quad \|f\|_p \leq C \|a(0, \partial)f\|_p, \quad \|f\|_p \leq C \|a(\infty, \partial)f\|_p,$$

for $f \in C_0^{\infty}(R^+)$. Inequalities (3.19) imply that the Mellin multiplier operators $a(0, \partial)$ and $a(\infty, \partial)$ have closed range and hence the symbols $a(0, z)$ and $a(\infty, z)$ have no zeroes on $\operatorname{Re} z = 1/p$.

q.e.d.

For an $N \times N$ system of operators in $Op - \Sigma_{1/p}$, the notion of ellipticity on $L^p([L^p(R^+)]^N)$ is that

$$(3.20) \quad \det[\sigma(A_{ij})] \neq 0, \quad (t, z) \in \partial R_{1/p},$$

where $[\sigma(A_{ij})]$ is the matrix of principal symbols. A theorem analogous to Theorem 3.2 may be proved for such a system.

Since $\partial R_{1/p}$ is homeomorphic to the unit circle S^1 , if A is an $N \times N$ system of operators which is

elliptic on L^p , then $[\sigma(A_{1j})]$, considered as a map from $\partial R_{1/p}$ to $GL(N, \mathbb{C})$, has a topological degree, $n_{1/p}$, defined as

$$(3.21) \quad \begin{cases} n_{1/p} = n_{1/p}(\sigma(A)) = \text{winding number } \det[\sigma(A_{1j})] \\ \quad = \frac{1}{2\pi} \Delta_{\partial R_{1/p}} \text{Arg } \det[\sigma(A_{1j})], \end{cases}$$

where the change in argument is calculated as $\partial R_{1/p}$ is traversed in the clockwise direction.

We emphasize that $n_{1/p}$ depends on $1/p$ and that if p is changed so that the line $\text{Re } z = 1/p$ crosses a zero of $\det[\sigma(A_{1j})](0, z)$ or $\det[\sigma(A_{1j})](\infty, z)$, the jump in $n_{1/p}$ may be calculated by the Argument Principle.

The analytical index of an $N \times N$ system $A = [a_{ij}(t, \partial)]$ of operators in $Op - \Sigma_{1/p}$ which is elliptic on L^p is defined as

$$(3.22) \quad \text{ind}_p(A) = \dim \ker A - \dim \ker A^*,$$

where in (3.22) we have considered A as an operator on $[L^p(\mathbb{R}^+)]^N$ and $A^* = [a_{ji}^*(t, \partial)]$ as an operator on $[L^q(\mathbb{R}^+)]^N$, $1/p + 1/q = 1$.

The index theorem we shall develop in Theorem 3.3 is that

$$(3.23) \quad \text{ind}_p(A) = n_{1/p}(\sigma(A)).$$

The first step in the proof of (3.23) is the construction of a canonical elliptic system A for which

$$(3.24) \quad \text{ind}_p(A) = 1 = n_{1/p}(\sigma(A)).$$

For this see the example of Cordes and Herman [CH,C], or consider the following example.

For $f \in L^p(\mathbb{R})$ define the (column) vector function $f = (f_1, f_2) \in [L^p(\mathbb{R}^+)]^2$ by

$$(3.25) \quad f_1(t) = f(t), \quad f_2(t) = f(-t).$$

The Hilbert transform on $L^p(\mathbb{R})$

$$(3.26) \quad \mathcal{H}f(x) = \text{p.v.} \frac{1}{\pi} \int_{-\infty}^{\infty} \frac{f(y)}{x-y} dy,$$

may be represented as the 2×2 system of operators in $\mathcal{O}_p - \Sigma_{1/p}$:

$$(3.27) \quad \begin{aligned} (\mathcal{H}f)_1(t) &= Hf_1(t) + Sf_2(t) \\ (\mathcal{H}f)_2(t) &= -Sf_1(t) - Hf_2(t), \end{aligned}$$

when $H = h(\partial)$ is a Hilbert transform on $L^p(\mathbb{R}^+)$ defined in (3.1-2) and

$$(3.28) \quad Sf(t) = s(\partial)f(t) = \frac{1}{\pi} \int_0^{\infty} \frac{f(s)}{t+s} ds$$

is the Stieltjes transform on $L^p(\mathbb{R}^+)$ with symbol $s(z) = \csc \pi z \in \mathcal{O}_{0,1}^{-\infty}$. For $f \in L^p(\mathbb{R})$, define

$$(3.29) \quad f(z) = \frac{1}{2\pi i} \int_{-\infty}^{\infty} \frac{f(x)}{x-z} dx, \quad \text{Im } z \neq 0,$$

$$f^{\pm}(x) = \lim_{\epsilon \rightarrow 0^+} f(x \pm i\epsilon) = \frac{1}{2} (I \pm iN)f(x).$$

For $x \in \mathbb{R}$ let $w = w(x) = \frac{x-1}{x+1}$; then $f(x) \rightarrow M_w f(x) = w(x)f(x)$ has the system representation

$$(3.30) \quad (M_w f)_1(t) = w(t)f_1(t), \quad (M_w f)_2(t) = \bar{w}(t)f_2(t).$$

It is well known that if on $L^p(\mathbb{R})$ we define $Gf(x) = w(x)f^-(x) + f^+(x)$, G has kernel spanned by $f(x) = \frac{1}{w(x)} - 1$ and cokernel 0. When G is written as a 2×2 system of pseudodifferential operators we obtain the matrices of principal symbols

$$(3.31) \quad \sigma(G)\left(t, \frac{1}{p} + i\infty\right) = \begin{bmatrix} w(t) & 0 \\ 0 & 1 \end{bmatrix}$$

$$\sigma(G)(0, z) = \begin{bmatrix} -ih & is \\ -is & ih \end{bmatrix}, \quad \sigma(G)(\infty, z) = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

$$\sigma(G)\left(t, \frac{1}{p} - i\infty\right) = \begin{bmatrix} 1 & 0 \\ 0 & \bar{w}(t) \end{bmatrix}.$$

Using (3.31) we have that $\text{ind}_p(G) = 1 = n_{1/p}(\sigma(G))$.

The second step in the proof of (3.23) is, given a system A of operators which is elliptic on L^p and such that $n_{1/p}(\sigma(A)) = 0$, to construct a

homotopy through elliptic operators to the identity operator.

Suppose that A_0 and A_1 are systems which are elliptic on L^p and there is a homotopy between the matrices of principal symbols

$$(3.32) \quad \sigma(A_s) : \partial R_{1/p} \rightarrow GL(N, \mathbb{C}), \quad 0 \leq s \leq 1$$

such that for each s , $\sigma(A_s)$ is a matrix of principal symbols, and

1. If $A_{s\pm}(t) = \sigma(A_s)(t, 1/p \pm i\infty)$, then

$$(3.33) \quad \lim_{s \rightarrow s'} \sup_{0 \leq t \leq \infty} \|A_{s\pm}(t) - A_{s'\pm}(t)\| = 0.$$

2. $\lim_{s \rightarrow s'} \sup_{\operatorname{Re} z = 1/p} \|\sigma(A_s)(0, z) - \sigma(A_{s'})(0, z)\| = 0,$

$$(3.34) \quad \lim_{s \rightarrow s'} \sup_{\operatorname{Re} z = 1/p} \left\| z \frac{d}{dz} [\sigma(A_s)(0, z) - \sigma(A_{s'})(0, z)] \right\| = 0.$$

3. $\lim_{s \rightarrow s'} \sup_{\operatorname{Re} z = 1/p} \|\sigma(A_s)(\infty, z) - \sigma(A_{s'})(\infty, z)\| = 0,$

$$(3.35) \quad \lim_{s \rightarrow s'} \sup_{\operatorname{Re} z = 1/p} \left\| z \frac{d}{dz} [\sigma(A_s)(\infty, z) - \sigma(A_{s'})(\infty, z)] \right\| = 0.$$

A homotopy (3.32) satisfying (3.33-35) will be called a homotopy through elliptic symbols; for each s we use the construction of (3.8) for a matrix of operators, A_s , with the given symbols $\sigma(A_s)$; by the 1-

dimensional Marcinkiewicz Theorem we have that

$s \rightarrow A_s$ is continuous from $[0,1]$ into the bounded operators on $[L^p(\mathbb{R}^+)]^N$. Hence $\text{ind}_p(A_0) = \text{ind}_p(A_1)$.

We now outline the proof of the Index Theorem. As a first step consider the scalar case.

LEMMA 3.1. Let $\sigma(A)$ be the symbol of an operator $A = a(t, \partial) \in \text{Op} - \Sigma_{1/p}$ which is elliptic on $L^p(\mathbb{R}^+)$ and suppose that

$$n_{1/p} = 0$$

where $n_{1/p} = n_{1/p}(\sigma(A))$ is defined by (3.21) with $N = 1$. Then

$$\text{ind}_p(A) = 0.$$

PROOF: The required homotopy through elliptic symbols is given by

$$\sigma(A_s)(t, z) = e^{s \log \sigma(A)}, \quad 0 \leq s \leq 1, \quad (t, z) \in \partial R_{1/p}.$$

q.e.d.

We now introduce a nonzero scalar function

$\varphi \in \mathcal{F}_{-\delta, \delta}$ such that $\varphi(0) = \varphi(\infty) = 1$ and $n(\varphi) = \frac{1}{2\pi} \Delta \int_0^\infty \text{Arg } \varphi = 1$. For $k \in \mathbb{Z}$ define the operator

$$(3.36) \quad A_k = a_k(t, \partial) = \varphi(t)^k \theta(\partial) + (I - \theta(\partial)).$$

Then A_k is elliptic on L^p for $1 < p < \infty$, and $n_{1/p}(\sigma(A_k)) = n(\sigma^k) = k$. Define

$$(3.37) \quad J = \text{ind}_p(A_1).$$

Then

$$(3.38) \quad \text{ind}_p(A_k) = k \text{ind}_p(A_1) = Jk.$$

(We will show in Theorem 3.3 that in fact, $J = 1$.)

The next Lemma is evident.

LEMMA 3.2. If $A = [\delta_{ij} a_{ij}(t, \partial)]$ is a diagonal sys-
tem of operators which is elliptic on L^p . Then

$$(3.39) \quad \text{ind}_p(A) = J n_{1/p}(\sigma(A)),$$

where J is defined as in (3.37).

We next calculate the index of a system whose symbols is the identity matrix on $\partial R_{1/p}$.

LEMMA 3.3. Let $\sigma(A) = [\sigma(A_{ij})]$ be an $N \times N$ matrix
of principal symbols a system A which is elliptic
on L^p . Suppose moreover that:

$$(3.40) \quad \begin{aligned} \sigma(A)(t, 1/p + i\infty) &= \sigma(A)(t, 1/p - i\infty) = I, \\ \sigma(A)(\infty, z) &= I. \end{aligned}$$

Then

$$(3.41) \quad \text{ind}_p(A) = J n_{1/p}(\sigma(A)).$$

PROOF: Multiplying A by a diagonal system, B , of operators satisfying the hypotheses of Lemma 3.2, condition (3.40), and such that $n_{1/p}(\sigma(B)) = -n_{1/p}(\sigma(A))$, we may assume that $n_{1/p}(\sigma(A)) = 0$.

From condition (3.40) we may choose c, d , $0 \leq c < 1/p < d \leq 1$ such that $a_{ij}(0, z) = \delta_{ij} + g_{ij}(z)$, where $g_{ij} \in \mathcal{O}_{c,d}^{-1}$ and $[a_{ij}(0, z)]$ has nonvanishing determinant in $S_{c,d}$.

Given $\epsilon > 0$, shrinking (c, d) if necessary, by Runge's Theorem there are rational functions

$g_{ij\epsilon} \in \mathcal{O}_{c,d}^{-1}$ such that

$$\sup_{S_{c,d}} |g_{ij} - g_{ij\epsilon}| < \epsilon, \quad \sup_{S_{c,d}} \left| z \frac{d}{dz} \{g_{ij} - g_{ij\epsilon}\} \right| < \epsilon.$$

Let $a_{ij\epsilon}(0, z) = \delta_{ij} + g_{ij\epsilon}(z)$ and let A_ϵ be a system of operators with principal symbol, $\sigma(A_\epsilon)$

which satisfies (3.40) and such that

$\sigma(A_\epsilon)(0, z) = [a_{ij\epsilon}(0, z)]$. If ϵ is small, then $n_{1/p}(\sigma(A_\epsilon)) = 0$ and $\text{ind}_p A_\epsilon = \text{ind}_p A$. We have reduced the problem to the case that $\sigma(A_\epsilon)(0, z)$ consists of rational functions.

Make a fractional linear transformation, $z \rightarrow w$, which maps the half plane $\{\text{Re } z < 1/p\}$ onto the interior of the unit circle $\{|w| < 1\}$ and such that as z traverses $\text{Re } z = 1/p + i\xi$, $-\infty \leq \xi \leq \infty$, w tra-

verses $\{|w| = 1\}$ in the clockwise direction. Define $G(w) = [a_{ij}(0, z)]$. Then $G: S^1 \rightarrow GL(N, \mathbb{C})$ is a rational function. Now (Cf., e.g., [V, pp. 45-49]) there are matrices of rational functions, $G_0(w)$, $G_\infty(w)$ holomorphic in $\{|w| \leq 1\}$ and $\{|w| \geq 1\}$ respectively such that

$$(3.42) \quad G(w) = G_0(w) D G_\infty(w)$$

where

$$(3.43) \quad D = D(n_1, \dots, n_N) = [\delta_{ij} w^{n_i}]$$

is a diagonal matrix and

$$n_1 + \dots + n_N = 0 = \frac{1}{2\pi} \Delta_{|w|=1} \operatorname{Arg} \det G(w).$$

For $0 \leq s \leq 1$, define

$$(3.44) \quad G_s(w) = G_0^{-1}(s) G_0(sw) D G_\infty(w/s) G_\infty^{-1}(1/s).$$

Formula (3.44) gives a homotopy through elliptic symbols to a diagonal matrix of operators which satisfies the hypotheses of Lemma 3.2.

q.e.d.

It is now straightforward to prove the formula

$$(3.45) \quad \operatorname{ind}_p(A) = J n_{1/p}(\sigma(A))$$

in the case that $\sigma(A)$ is the principal symbol of an

elliptic operator whose principal symbol is the identity matrix I at the corners of $\partial R_{1/p}$; i.e.,

$$(3.46) \quad \sigma(A)(0, 1/p \pm i\infty) = \sigma(A)(\infty, 1/p \pm i\infty) = I.$$

To handle the general case we use the following Lemma whose proof we leave to the reader.

LEMMA 3.4. Given 4 matrices A_{0+} , A_{0-} , $A_{\infty+}$, $A_{\infty-}$ in $GL(N, \mathbb{C})$, there is a symbol, $\sigma(A)$, of an elliptic system on L^p such that

$$(3.47) \quad \begin{aligned} \sigma(A)\left(0, \frac{1}{p} \pm i\infty\right) &= A_{0, \pm} \\ \sigma(A)\left(0, \frac{1}{p} \pm i\infty\right) &= A_{\infty, \pm} \end{aligned}$$

and such that

$$(3.48) \quad \text{ind}_p(A) = J n_{1/p}(\sigma(A)).$$

We are now ready to prove (3.23).

THEOREM 3.3. (The Index Theorem). If A is a system of operators in $\text{Op } \Sigma_{1/p}$ which is elliptic on L^p , then

$$(3.23) \quad \text{ind}_p(A) = n_{1/p}(\sigma(A)).$$

PROOF: By Lemmas 3.1 to 3.4 we have that

$\text{ind}_p(A) = J n_{1/p}(\sigma(A))$. The example studied in (3.25) to (3.31) shows that $J = 1$. q.e.d.

4. Pseudodifferential operators on a finite interval.

We shall develop an algebra of pdo's on a finite interval $I = [0, \pi]$. For $f \in L^p(I)$, define

$$Tf(t) = f(\pi - t).$$

Then $T^* = T$ and $TT = I$. If A is a bounded operator on $L^p(I)$, the essential requirement for A to be in our class of pdo's is that, when acting on functions supported in $[0, \pi)$, both A and TAT , modulo compact operators, are in $Op - \Sigma_{1/p}$ on R^+ .

DEFINITION 4.1. A bounded operator A on $L^p(I)$ is a pdo of class $Op - \Sigma_{1/p}(I)$ iff

1. If $\varphi, \psi \in C_0^\infty(R)$ have disjoint supports then the map

$$(4.1) \quad f \rightarrow (\varphi A \psi)f = \varphi \chi_{[0, \pi]} A(\chi_{[0, \pi]} \psi f)$$

is a compact operator on $L^p(R^+)$.

2. If $\varphi, \psi \in C_0^\infty([0, \pi))$, there is an operator $A_{\varphi\psi} \in Op - \Sigma_{1/p}$ and a compact operator $K_{\varphi\psi}$ on $L^p(I)$ such that

$$(4.2) \quad \varphi A \psi = A_{\varphi\psi} + K_{\varphi\psi}.$$

3. The operator

$$(4.3) \quad f \rightarrow TATf = A_T f$$

with $Tf(t) = f(\pi - t)$ satisfies conditions 1 and 2.

An example of an operator of class $Op - \Sigma_{1/p}(I)$ is the finite Hilbert transform

$$(4.4) \quad Hf(t) = \text{p.v.} \frac{1}{\pi} \int_0^\pi \frac{f(s)}{t-s} ds, \quad (THT = -H),$$

and the finite Stieltjes transform

$$(4.5) \quad Sf(t) = \text{p.v.} \frac{1}{\pi} \int_0^\pi \frac{f(s)}{t+s} ds.$$

For $\varphi, \psi \in C_0^\infty([0, \pi))$, $\varphi TST\psi$ is a compact operator on $L^p(I)$.

To define the principal symbol of a pdo of class $Op - \Sigma_{1/p}(I)$, we first note that if $\varphi, \psi \in C_0^\infty([0, \pi))$ and $a_{\varphi\psi}(t, \partial) = A_{\varphi\psi} \in Op - \Sigma_{1/p}$ is defined so that

$$(4.6) \quad \varphi A\psi = A_{\varphi\psi} + K_{\varphi\psi}$$

as in (4.2), then there is a well defined function $a_0(t, z)$ defined on

$$(4.7) \quad \Lambda_{1/p} = \{(t, z) : (t, z) \in \partial R_{1/p}, 0 \leq t < \pi\}$$

such that

$$(4.8) \quad \sigma(a_{\varphi\psi}(t, \partial)) = \varphi(t)a_0(t, z)\psi(t)$$

for $(t, z) \in \Lambda_{1/p}$. The function $a_0(t, z)$ is defined by choosing $\varphi, \psi \in C_0^\infty([0, \pi))$ which are $= 1$ in a neighborhood of t and letting

(4.9) $a_0(t,z) = \sigma(a_{\varphi\psi})(t,z).$

The formula (4.9) uniquely determines $a_0(t,z)$ on $\Lambda_{1/p}$; if φ_1 and ψ_1 are also = 1 near t then $\varphi_1\varphi A\psi_1 = \varphi_1\psi_1 a_{\varphi\psi}(t,\partial) + K = \varphi\psi a_{\varphi_1\psi_1} + K_1$ where K and K_1 are compact operators on $L^p(\mathbb{R}^+)$, and by Theorem 3.1, $\varphi_1\psi_1 a_{\varphi\psi}$ and $\varphi\psi a_{\varphi_1\psi_1}$ have the same principal symbols.

The operator $A_T = TAT$ has a well defined principal symbol on $\Lambda_{1/p}$ which we denote by $a_\pi(t,z).$

To glue together the symbols $a_0(t,z)$ and $a_\pi(t,z)$ we will show that for $0 < t < \pi$, the following compatibility conditions are satisfied:

(4.10)
$$\begin{aligned} a_0(t, 1/p + i\infty) &= a_\pi(\pi - t, 1/p - i\infty), \\ a_0(t, 1/p - i\infty) &= a_\pi(\pi - t, 1/p + i\infty). \end{aligned}$$

To prove (4.10) we introduce the operators

(4.11)
$$\begin{aligned} \Theta &= \frac{1}{2} (I - iH) \\ I - \Theta &= \frac{1}{2} (I + iH) = T\Theta T \end{aligned}$$

with H the finite Hilbert transform given by (4.4).

Fix $\varphi \in C^\infty_c((0,\pi))$ such that $T\varphi(t) = \varphi(t)$ and $\varphi(t) = 1$ on $(\delta, \pi - \delta)$. As operators on $L^p(\mathbb{R}^+)$,

(4.12)
$$\begin{aligned} \varphi A\varphi &= \varphi\{a_0(t, 1/p + i\infty)\Theta(\partial) \\ &\quad + a_0(t, 1/p - i\infty)\} \varphi + K_0, \end{aligned}$$

and

(4.13)
$$\varphi T A T \varphi = \varphi \{ a_{\pi}(t, 1/p + i\infty) \theta(\partial) + a_{\pi}(t, 1/p - i\infty) \} \varphi + K_{\pi},$$

with K_0 and K_{π} compact on $L^p(\mathbb{R}^+)$ and $\theta(\partial)$ defined by (3.2). Modulo compact operators, on $L^p(I)$, we have the representation:


$$\begin{aligned} \varphi A \varphi &= \varphi (a_0(t, 1/p + i\infty) \Theta \\ &\quad + a_0(t, 1/p - i\infty) (I - \Theta)) \end{aligned}$$

and since $T \Theta T = I - \Theta$,

(4.14)
$$\begin{aligned} T \varphi A \varphi T &= \varphi \{ a_0(\pi - t, 1/p + i\infty) (I - \Theta) \\ &\quad + a_0(\pi - t, 1/p - i\infty) \Theta \} \varphi. \end{aligned}$$

Comparing (4.14) to (4.13) we obtain (4.10).

If A is a pdo of class $Op - \Sigma_{1/p}(I)$, the principal symbol of A , $\sigma(A)$, is defined as the pair of functions $a_0(t, z)$, $a_{\pi}(t, z)$ defined on $\wedge_{1/p}$ which satisfy the compatibility conditions (4.10). The natural setting for the principal symbol as a continuous function defined on the boundary of a compact rectangle $R_{1/p, I}$ in (4.15):



It can be shown that a pdo of class $Op - \Sigma_{1/p}(I)$ is compact iff its principal symbol is $\equiv 0$.

The principal symbol of an $N \times N$ system of pdo's of class $Op - \Sigma_{1/p}(I)$ is defined as the matrix of principal symbols.

DEFINITION 4.2. An $N \times N$ system of pdo's of class $Op - \Sigma_{1/p}(I)$ is elliptic on $L^p = [L^p(I)]^N$ iff the determinant of the matrix of principal symbols does not vanish on $\partial R_{1/p, I}$.

THEOREM 4.1. Let $A = [A_{ij}]$ be an $N \times N$ system of pdo's of class $Op - \Sigma_{1/p}(I)$. The following are equivalent:

1. A is elliptic on L^p .
2. There is an $N \times N$ system of pdo's of class $Op - \Sigma_{1/p}(I)$ such that

$$AB - I \text{ and } BA - I$$

are compact operators on $[L^p(I)]^N$.

3. There is a bounded operator B on $[L^p(I)]^N$ such that $BA - I$ is a compact operator on $[L^p(I)]^N$.

4. For $\vec{f} = (f_1, \dots, f_N)$ and N -tuple of func-
tions in $C_0^\infty((0,1))$, there is an a priori esti-
mate

(4.16) $\|f; L^p(I)\| \leq c \|Af: L^p(I)\| + \|Kf: L^p(I)\|$

where K is a compact operator on $[L^p(I)]^N$.

PROOF: That 1. implies 2. implies 3. implies 4. is obvious. If (4.16) holds, let $f \in [C_0^\infty((\delta, \pi - \delta))]^N$, and let $\varphi \in C_0^\infty([0, \pi])$ be such that $\varphi = 1$ on $[0, \pi - \delta)$. Then $Af = \varphi A_0 \varphi f + K$, where K is a compact operator on $L^p(\mathbb{R}^+)$. Using the techniques in the proof of Theorem 3.2, we obtain the apriori estimates

(4.17) $\|f; L^p(\mathbb{R}^+)\| \leq c \|A_0(t; 1/p \pm i\epsilon)f; L^p(\mathbb{R}^+)\|,$

(4.18) $\|f; L^p(\mathbb{R}^+)\| \leq c \|A_0(0, \delta)f; L^p(\mathbb{R}^+)\|.$

Inequalities (4.17) and (4.18) are valid for all $\vec{f} \in [C_0^\infty((\delta, \pi - \delta))]^N$, with c independent of δ . Since $A_0(0, \delta)$ commutes with dilations, (4.18) is valid for all $\vec{f} \in [C_0^\infty(\mathbb{R}^+)]^N$ and therefore $\det[a_{1j0}(0, z)] \neq 0$, $z = 1/p + i\epsilon$, $-\infty \leq \epsilon \leq \infty$. The continuity of $\sigma(A)(t, z)$ and (4.17) show that $[a_{1j0}(t, z)]$ is nonsingular on $\Lambda_{1/p}$.

q.e.d.

If A is a system of pdo's of class $Op - \Sigma_{1/p}(I)$ which is elliptic on L^p we define

(4.19) $n_{1/p}(\sigma(A)) = \frac{1}{2\pi} \Delta_{\partial R_{1/p, I}} \text{Arg } \det(\sigma(A));$

the change in argument being taken as $\partial R_{1/p, I}$ in (4.15) is traversed in the clockwise direction. The analytical index of the system A is defined as

$$(4.20) \quad \text{ind}_p(A) = \dim \ker A - \dim \ker A^*,$$

where A is considered as an operator on $[L^p(I)]^N$ and A^* is considered as an operator on $[L^q(I)]^N$, $1/p + 1/q = 1$.

Before proving the Index Theorem 4.2, i.e., that

$$(4.21) \quad \text{ind}_p(A) = n_{1/p}(\sigma(A)),$$

we consider a canonical system A for which we show directly that

$$(4.22) \quad \text{ind}_p(A) = 1 = n_{1/p}(\sigma(A)).$$

Let $S^1 = \{w \in \mathbb{C}; |w| = 1\}$ be the unit circle and for $f(w) = f(e^{i\varphi}) \in C^\infty(S^1)$ define for $z = e^{i\theta}$

$$(4.23) \quad \begin{aligned} Bf(z) &= \text{p.v.} \frac{1}{\pi i} \int_{|\zeta|=1} \frac{f(\zeta)}{\zeta - z} d\zeta \\ &= \frac{1}{2\pi} \int_{-\pi}^{\pi} f(e^{i\varphi}) d\varphi - \text{p.v.} \frac{1}{2\pi i} \int_{-\pi}^{\pi} f(e^{i\varphi}) \cot\left(\frac{\theta - \varphi}{2}\right) d\varphi. \end{aligned}$$

Note that $f \rightarrow \frac{1}{2\pi} \int_{-\pi}^{\pi} f(e^{i\varphi}) d\varphi$ is a compact operator on $L^p(-\pi, \pi)$. Given $f \in L^p(S^1)$ we define the column vector function $\vec{f} = (f_1, f_2) \in [L^p(I)]^2$ by

$$\begin{aligned}
 (4.24) \quad f_1(t) &= f(e^{1t}) & 0 \leq t \leq \pi \\
 f_2(t) &= f(e^{-1t}) & 0 \leq t \leq \pi.
 \end{aligned}$$

We consider the operator

$$(4.25) \quad B_{11}f(t) = \text{p.v.} \frac{1}{\pi} \int_0^\pi f(s) \frac{1}{2} \cot\left(\frac{t-s}{2}\right) ds, \quad t \in [0, \pi].$$

If $\varphi \in C_0^\infty([0, \pi])$, $\varphi B_{11}\varphi f = \varphi H\varphi f + Kf$, where K is a compact operator on $L^p(0, \pi)$ and H is the finite Hilbert transform defined by (4.4). Now

$$TB_{11}T = -B_{11}$$

so that B_{11} differs from H by a compact operator.

Now consider

$$(4.26) \quad B_{12}f(t) = \frac{1}{\pi} \int_0^\pi f(s) \frac{1}{2} \cot\left(\frac{t+s}{2}\right) ds, \quad t \in [0, \pi].$$

The singularity of the kernel $\frac{1}{2} \cot\left(\frac{t+s}{2}\right)$ occurs when $t+s$ is near 0 or near 2π . If $\varphi \in C_0^\infty([0, \pi])$,

$$\begin{aligned}
 (4.27) \quad \varphi B_{12}\varphi f(t) &= \frac{1}{\pi} \int_0^\pi \frac{\varphi(t)\varphi(s)}{t+s} f(s) ds + Kf(t), \\
 &= (\varphi S\varphi)f + Kf,
 \end{aligned}$$

where K is compact on $L^p(I)$ and S is the finite Stieltjes transform defined by (4.5). Also

$$(4.28) \quad TB_{12}T = -B_{12}.$$

The operator B defined in (4.23) on $L^p(S^1)$, using the correspondence (4.24) may be represented,

modulo a compact operator, as a system on $L^p(I) \times L^p(I)$:

$$(4.29) \quad \begin{aligned} (Bf)_1(t) &= B_{11}f_1(t) + B_{12}f_2(t) \\ (Bf)_2(t) &= -B_{12}f_1(t) - B_{11}f_2(t), \end{aligned}$$

which is a 2×2 system of pdo's of class $Op - \Sigma_{1/p}(I)$

whose principal symbols are given in (4.30). (In

$$(4.30) \quad h(z) = -\cot \pi z, \quad s(z) = \csc \pi z.)$$

$$(4.30) \quad \begin{array}{ccc} & \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix} & \\ & \downarrow & \\ \begin{bmatrix} h & s \\ -s & h \end{bmatrix} & \xrightarrow{\partial R_{1/p, I}} & \begin{bmatrix} -h & -s \\ s & h \end{bmatrix} \\ & \uparrow & \\ & \begin{bmatrix} -1 & 0 \\ 0 & 1 \end{bmatrix} & \end{array}$$

Given $f(e^{i\theta}) \in C^\infty(S^1)$, define for $z = e^{i\theta}$, let

$$f^+(z) = \lim_{r \uparrow 1} \frac{1}{2\pi i} \oint \frac{f(\zeta)}{\zeta - rz} d\zeta, \quad \text{and} \quad f^-(z) = \lim_{r \uparrow 1} \oint \frac{f(\zeta)}{\zeta - rz} d\zeta.$$

Modulo a compact operator,

$$f^\pm(e^{i\theta}) = \frac{1}{2} \{f(e^{i\theta}) \pm iBf(e^{i\theta})\}.$$

Finally, note that the multiplication operator

$f(w) \rightarrow wf(w)$ has the representation as a system:

$$f_1 \rightarrow e^{it}f_1; \quad f_2 \rightarrow e^{-it}f_2.$$

The operator

$$(4.31) \quad Af(w) = wf^-(w) + f^+$$

has a one dimensional kernel spanned by $1/w - 1$ and cokernel $\{0\}$. As a system of pdo's of class $Op - \Sigma_{1/p}(I)$, a direct calculation shows that $n_{1/p}(\sigma(A)) = 1$ so that

$$(4.32) \quad \text{ind}_p(A) = 1 = n_{1/p}(\sigma(A)).$$

We are now ready to prove the Index Theorem.

THEOREM 4.2. If A is an $N \times N$ matrix of pdo's of class $\Sigma_{1/p}(I)$ which is elliptic on L^p , then

$$\text{ind}_p(A) = n_{1/p}(\sigma(A)).$$

PROOF: We may follow the proofs of Lemmas 3.1-3.4 with only a small change in notation to prove the same special cases. Hence there is an integer J_I such that

$$\text{ind}_p(A) = J_I n_{1/p}(\sigma(A)).$$

By the example considered above, $J_I = 1$.

q.e.d.

5. Single and double layer potentials in polygonal domains.

As an application of pdo's of class $\Sigma_{1/p}(I)$ we use single and double layer potentials to study the

Dirichlet, Neumann, and Oblique Derivative Problems
for Laplace's equation in a polygon.

Let Ω be a simply connected polygon in R^2 with N successive vertices labeled as $P_1, P_3, \dots, P_{2N-1} = P_1$ as $\partial\Omega$ is traversed in the counter clockwise direction. Label the interior angles at P_{2k-1} as θ_{2k-1} , $k = 1, \dots, N$. At the midpoint of $\overline{P_{2k-1}P_{2k+1}}$ we introduce a false vertex P_{2k} with interior angle $\theta_{2k} = \pi$ and parametrize the new half sides with $t \in [0, 1]$ so that $t = 0$ at P_{2k-1} or P_{2k+1} and $t = 1$ at P_{2k} ; i.e.:

For $t \in [0, 1]$, define

$$P_t = P_{t, 2k-1} = tP_{2k} + (1-t)P_{2k-1} \in \overline{P_{2k-1}P_{2k}}, \quad (5.1)$$

$$P_t = P_{t, 2k-2} = tP_{2k-2} + (1-t)P_{2k-1} \in \overline{P_{2k-1}P_{2k-2}}$$

If l_1 is the length of $\overline{P_1P_{1+1}}$, the arclength $d\sigma$ is given by

$$(5.2) \quad d\sigma = (-1)^{1+l_1} l_1 dt.$$

For $\phi \in L^p(\partial\Omega)$ define the double layer potential

$$(5.3) \quad u(X) = \frac{1}{\pi} \int_{\partial\Omega} \frac{\langle X-Q, n_Q \rangle}{|X-Q|^2} \phi(Q) d\sigma_Q,$$

where n_Q is the interior unit normal to a point $Q \in \partial\Omega$.

For $P \in \partial\Omega$, let

$$\begin{aligned}
 (5.4) \quad u^+(P) &= \lim_{X \rightarrow P, X \in \Omega} u(X) = \phi(P) + K\phi(P), \\
 u^-(P) &= \lim_{X \rightarrow P, X \in \mathbb{R}^2 - \Omega} u(X) = -\phi(P) + K\phi(P),
 \end{aligned}$$

where

$$(5.5) \quad K\phi(P) = \frac{1}{\pi} \int_{\partial\Omega} \frac{\langle P-Q, n_Q \rangle}{|P-Q|^2} \phi(Q) d\sigma_Q,$$

and the limits in (5.4) are taken nontangentially in $L^p(\partial\Omega)$ as $X \rightarrow P \in \partial\Omega$. Using the correspondences given by (5.1), consider ϕ and $K\phi$ as functions in $[L^p(0,1)]^{2N}$, where $\phi_1(t) = \phi(P_t)$; $P_t \in \overline{P_1 P_{1+1}}$. Then K may be represented as a $2N \times 2N$ system of pdo's of class $Op - \Sigma_{1/p}(I)$, $I = [0,1]$. We obtain the following operators for $K = [K_{ij}]$:

$$\begin{aligned}
 (5.6) \quad K_{11} &= 0, \quad K_{2k-1,2k} = K_{2k,2k-1} = 0, \\
 K_{ij} &\text{ is compact if } \overline{P_i P_{i+1}} \text{ and } \overline{P_j P_{j+1}} \text{ do not touch,} \\
 K_{2k-2,2k-1}\varphi(t) &= \int_0^1 k_\theta(l_{2k-2}t/l_{2k-1}s)\varphi(s) \frac{ds}{s}, \\
 K_{2k-1,2k-2}\varphi(t) &= \int_0^1 k_\theta(l_{2k-1}t/l_{2k-2}s)\varphi(s) \frac{ds}{s}, \\
 &\text{with } \theta = \theta_{2k-1}
 \end{aligned}$$

where

$$(5.7) \quad k_\theta(t) = \frac{1}{\pi} \frac{t \sin \theta}{t^2 + 1 - 2t \cos \theta}.$$

The Mellin transform of the kernel $k_\theta(t)$ is given by

$$(5.8) \quad \tilde{k}_\theta(z) = \frac{\sin(\pi-\theta)z}{\sin \pi z} \in \mathcal{O}_{0,1}^{-\infty}.$$

The matrix of principal symbols of the operator $[b_{ij}I + K_{ij}]$ is the identity on the top, bottom, and r.h.s. of $\partial R_{1/p,I}$; at $t = 0$ its determinant can be calculated by using an even number of row and column transpositions to obtain a matrix with 2×2 blocks on the diagonal; the vertex P_{2k-1} contributes

$$(5.8) \quad \begin{bmatrix} 1 & b^z \tilde{k}_\theta(z) \\ b^{-z} \tilde{k}_\theta(z) & 1 \end{bmatrix}$$

where $b = \mu_{2k-2}/\mu_{2k-1}$ and $\theta = \theta_{2k-1}$. Hence

$$\det(\sigma(I+K))(0,z) = \prod_{j=1}^N (1 - [\tilde{k}_{\theta_{2j-1}}(z)]^2).$$

The zeroes of $1 - [\tilde{k}_\theta(z)]^2$ occur at $z = n\pi/\theta$ or $z = n\pi/(2\pi-\theta)$. If $\theta \neq \pi$ we define p_θ by

$$(5.9) \quad 1/p_\theta = \begin{cases} \pi/(2\pi-\theta), & 0 < \theta < \pi, \\ \pi/\theta, & \pi < \theta < 2\pi, \end{cases}$$

so that $z_\theta = 1/p_\theta$ is the simple root of $1 - k_\theta^2 = 0$ satisfying $0 < \operatorname{Re} z < 1$; in fact $1 < p_\theta < 2$. By the Argument Principle

$$(5.10) \quad n_{1/p}(\theta) = \frac{1}{2\pi} \Delta_{z=1/p+i\epsilon}^{\operatorname{Arg} (1 - [k_\theta]^2)} \quad -\infty \leq \epsilon \leq +\infty$$

$$= \begin{cases} 0, 0 < 1/p < 1/p_\theta \\ 1, 1/p_\theta < 1/p < 1 \end{cases}.$$

From the results of Chapter 4, the system

$$(5.11) \quad (I+K)\phi = \psi$$

is Fredholm on $L^p(\partial\Omega)$ iff $p \neq p_{\theta_1}, \dots, p_{\theta_{2N-1}}$; in case (5.11) is Fredholm on $L^p(\partial\Omega)$ we have

$$(5.12) \quad \text{ind}_p(I+K) = \sum_{p < p_{\theta_{2k-1}}} 1.$$

In particular, since $p_\theta < 2$, for $p \geq 2$, $\text{ind}_p(I+K) = 0$.

To study the kernel and cokernel of (5.11) we introduce the single layer potential

$$(5.13) \quad v(X) = \frac{1}{2\pi} \int_{\partial\Omega} |X-Q|^2 \psi(Q) d\sigma_Q.$$

Then

$$(5.14) \quad \frac{\partial v}{\partial n^+}(P) = \lim_{X \rightarrow P, X \in \Omega} \frac{\partial v}{\partial n_P}(X) = (I-K^*)\psi$$

$$\frac{\partial v}{\partial n^-}(P) = \lim_{X \rightarrow P, X \in \mathbb{R}^2 - \Omega} \frac{\partial v}{\partial n_P}(X) = -(I+K^*)\psi,$$

where

$$(5.15) \quad K^*\psi(P) = \frac{1}{\pi} \int_{\partial\Omega} \frac{\langle Q-P, n_P \rangle}{|P-Q|^2} \psi(Q) d\sigma_Q.$$

Let M be the (compact) mean value operator

$$M\phi = \frac{1}{|\partial\Omega|} \int_{\partial\Omega} \phi(Q) d\sigma_Q.$$

From, e.g., Petrovsky [Pe, §34], we recall the following facts for these operators on $L^p(\partial\Omega)$:

$\ker(I-M) = \{\text{constant functions}\}$; $(I+K)M = 2M$; and $\ker(I+K^*)(I-M) = \{\text{constant functions}\}$.

Define

$$(5.16) \quad p_0 = \max p_{\theta_{2k-1}}$$

with p_{θ} given by (5.9). Note that $p_0 < 2$.

LEMMA 5.1. For $p > p_0$ the kernel of $(I-M)(I+K)$ consists of constant functions.

PROOF: Constant functions are in the kernel of

$$(I-M)(I+K) \quad \text{and} \quad \text{ind}_p(I-M)(I+K) = \text{ind}_p(I+K) = 0.$$

Moreover, $(I+K^*)(I-M)$ has kernel of dimension 1.

q.e.d.

LEMMA 5.2. For $p > p_0$,

$$(5.17) \quad \ker(I+K) = \{0\}.$$

PROOF: If $(I+K)\phi = 0$ then $\phi = \text{constant}$ by Lemma

5.1. But then $(I+K)\phi = 2\phi = 0$.

q.e.d.

LEMMA 5.3. For $1 < q < \infty$

$$\ker(I+K^*) \cap L^q(\partial\Omega) = \{0\}.$$

PROOF: If $1 < q \leq 2$ and $1/p + 1/q = 1$, then $p > p_0$. Hence $\text{ind}_q(I+K^*) = 0$ and $\ker(I+K) \cap L^p(\partial\Omega) = \{0\}$. Hence $\ker(I+K^*) \cap L^q(\partial\Omega) = \{0\}$. If $2 \leq q$, $\ker(I+K^*) \cap L^q(\partial\Omega) \subset \ker(I+K^*) \cap L^2(\partial\Omega) = \{0\}$.
q.e.d.

THEOREM 5.1. If K is defined by (5.5) then the integral equation

$$(5.11) \quad (I+K)\phi = \psi$$

considered on $L^p(\partial\Omega)$ has the following properties:

1. (5.11) is Fredholm on $L^p(\partial\Omega)$ iff

$$(5.16) \quad p \neq p_{\theta_1}, \dots, p_{\theta_{2N-1}},$$

2. For $1 < q < \infty$, $\ker(I+K^*) \cap L^q(\partial\Omega) = \{0\}$.

3. For every p satisfying (5.16),

$$(5.17) \quad \dim \ker(I+K) \cap L^p(\partial\Omega) = \begin{cases} 0, & p \geq 2 \\ \text{no. of vertices with} \\ \text{angle } \theta \text{ such that} \\ p < p_\theta \end{cases}$$

(p_θ given by (5.9)).

The proof of Theorem 5.1 follows from the preceding discussion.

We next seek to resolve the Neumann problem $\Delta v = 0$ in Ω , $\frac{\partial v}{\partial n^+}(P) = g(P) \in L^q(\partial\Omega)$ in the form of a single layer potential given by (5.13); using (5.14) we study the integral equation

$$(5.18) \quad (I-K^*)\psi = g$$

on $L^q(\partial\Omega)$. The operator $I-K$ is a Fredholm operator on $L^p(\partial\Omega)$ iff $p \neq p_{\theta_1}, \dots, p_{\theta_{2N-1}}$ and for such values of p has the same index on $L^p(\partial\Omega)$ as the operator $I+K$. We show that the kernel of $(I-K^*)$ consists of constants.

We show that the kernel of $I-K^*$ has dimension 1. Let q be near 1 and $S_q = \ker(I-K^*) \cap L^q(\partial\Omega)$. Now $S_q \neq \{0\}$ since $\{\text{constants}\} \subset \ker(I-K)$ and $\text{ind}_q(I-K) = 0$. It is easy to show that if $\psi \in S_q$ and $M\psi = 0$, then $\psi = 0$. Hence there is a $\psi_1 \in S_q$ such that $M\psi_1 = 1$. Then if $\psi \in S_q$, $\psi = (M\psi)\psi_1$.

For (5.18) we have the following theorem.

THEOREM 5.2. The integral equation

$$(5.18) \quad (I-K^*)\psi = g$$

on $L^q(\partial\Omega)$ has the following properties.

1. For $1 < q < \infty$,

(5.19) $\dim \ker(I-K^*) \cap L^q(\partial\Omega) = 1.$

2. (5.18) is Fredholm on $L^q(\partial\Omega)$ iff

(5.20) $q \neq q_{\theta_1}, \dots, q_{\theta_{2N-1}}; (1/q_{\theta_1} + 1/p_{\theta_1} = 1).$

$(p_{\theta}$ given by (5.9)).

3. If q satisfies (5.20),

$$\operatorname{ind}_q(I-K^*) = \begin{cases} 0, & q \leq 2, \\ -(\text{no. of vertices with} \\ \text{angle } \theta \text{ for which} \\ q > q_{\theta}). \end{cases}$$

COROLLARY 5.1. The integral equation

(5.21) $(I-K)\phi = \psi$

on $L^p(\partial\Omega)$ has the following properties.

1. If $p > p_0$, $\ker(I-K) \cap L^p(\partial\Omega) = \{\text{constant functions}\}.$
2. For $p \neq p_{\theta_1}, \dots, p_{\theta_{2N-1}}$
- $$\dim \ker(I-K) \cap L^p(\partial\Omega) = 1 + (\text{no. of vertices with} \\ \text{angle } \theta \text{ for which} \\ p < p_{\theta}).$$


Finally we apply the pdo theory to study the
Oblique Derivative Problem or a Mixed Boundary Value

Problem in a polygon. We use the same notation as before, and use single layer potentials to study:

$$(5.22) \quad \begin{aligned} \Delta v &= 0 \quad \text{in } \Omega, \\ \frac{\partial v}{\partial \nu} &= g(P) \in L^p(\partial\Omega), \end{aligned}$$

where on each segment $\overline{P_{2k-1}P_{2k+1}}$, $v = v(P)$ is a smooth nonvanishing vector field. Dirichlet data can be handled by taking v tangent to $\partial\Omega$; jump discontinuities in v may be considered by introducing a new vertex with angle $\theta = \pi$ at the jump.

For simplicity we assume that each segment $\overline{P_iP_{i+1}}$ has unit length. On each segment $\overline{P_iP_{i+1}}$ we introduce unit tangent vectors τ_i in the direction of increasing t as in (5.23):



Normalize so that $|v(P)| = 1$ and define smooth angles $\gamma_i = \gamma_i(t)$ so that on $\overline{P_iP_{i+1}}$;

$$v(P_t) = \cos \gamma_i(t) \tau_i + \sin \gamma_i(t) n_i.$$

The continuity of $v(P)$ at the even vertices may be expressed as:

$$(5.24) \quad \gamma_{2k-1}(1) + \gamma_{2k}(1) = \pi \pmod{2\pi}.$$

Consider the operators

$$(5.25) \quad A^\tau \psi(P) = \lim_{X \rightarrow P, X \in \Omega} \frac{\partial v}{\partial \tau_P}(X)$$

$$= \text{p.v.} \frac{1}{\pi} \int_{\partial \Omega} \frac{\langle P-Q, \tau_P \rangle}{|P-Q|^2} \psi(Q) d\sigma_Q,$$

$$(5.26) \quad A^n \psi(P) = (I-K^*)\psi(P) = \lim_{\epsilon \rightarrow 0^+} \frac{\partial v}{\partial n_P}(P + \epsilon n_P),$$

with v the single layer potential given by (5.13).

To represent the operators A^τ and A^n as systems of pdo's of class $\text{Op} - \Sigma_{1/p}(I)$, we introduce the Hardy kernels (and their Mellin transforms):

$$k_{\tau\theta}(t) = \frac{1}{\pi} \frac{t - \cos \theta}{t^2 + 1 - 2t \cos \theta}$$

$$\tilde{k}_{\tau\theta}(z) = - \frac{\cos((\pi-\theta)z + \theta)}{\sin \pi z}$$

$$(5.27) \quad k_{n\theta}(t) = \frac{1}{\pi} \frac{-\sin \theta}{t^2 + 1 - 2t \cos \theta}$$

$$\tilde{k}_{n\theta}(z) = - \frac{\sin((\pi-\theta)z + \theta)}{\sin \pi z}$$

$$s(t) = k_{\tau\pi}(t) = \frac{1}{\pi} \frac{1}{t+1}$$

$$\tilde{s}(z) = \frac{1}{\sin \pi z}$$

and the corresponding Hardy kernel operations:

(5.28)
$$K\psi(t) = \int_0^1 k\left(\frac{t}{s}\right) f(s) \frac{ds}{s}$$

which, as operators in $Op - \Sigma_{1/p}(I)$ have principal symbols, $\sigma(K)$, given by

(5.29)
$$\tilde{K}(z) \begin{array}{c} 0 \\ \boxed{R_{1/p,I}} \\ 0 \end{array} 0 \ .$$

For the operator $A^T = [A^T_{ij}]$, we obtain

1. $A^T_{11}\psi_1(t) = H\psi_1(t) = p.v. \frac{1}{\pi} \int_0^1 \frac{\psi_1(s)}{t-s} ds.$
2. $A^T_{2k-1,2k}\psi_{2k}(t) = \frac{1}{\pi} \int_1^2 \frac{1}{t-s} \psi_{2k}(2-s)ds$
 $= -TST\psi_{2k}(t).$
3. $A^T_{2k,2k-1}\psi_{2k-1}(t) = \frac{1}{\pi} \int_0^1 \frac{-1}{2-t-s} \psi_{2k-1}(s)ds$
 $= -TST\psi_{2k-1}(t).$
4. $A^T_{2k-2,2k-1} = A^T_{2k-1,2k-2} = K_{\tau\theta}, \theta = \theta_{2k-1}.$
5. A^T_{ij} is compact if $\overline{P_iP_{i+1}}$ and $\overline{P_jP_{j+1}}$ do not touch.

For the operator $A^n = [A_{ij}^n] = (I - K^*)$ we have

$$6. \quad A_{ii}^n = I.$$

$$7. \quad A_{2k-1, 2k}^n = A_{2k, 2k-1}^n = 0.$$

$$8. \quad A_{2k-2, 2k-1}^n = A_{2k-1, 2k-2}^n = K_{n\theta}.$$

$$9. \quad A_{ij}^n \text{ is compact if } \overline{P_1 P_{i+1}} \text{ and } \overline{P_j P_{j+1}} \text{ do not touch.}$$

The system which arises to study (5.22) is now

$$(5.30) \quad (\nu \cdot \tau_P) A^\tau \psi(P) + (\nu \cdot n_P) A^n \psi(P) = g(P).$$

on $L^p(\partial\Omega)$.

We now calculate the determinant of the matrices of principal symbols of (5.40) defined on $\partial R_{1/p, I}$.

On the r.h.s. of $\partial R_{1/p, I}$; i.e., when $t = 1$, we make an even number of row and column transpositions to obtain N 2×2 blocks along the diagonal; each even numbered vertex P_{2k} contributes with $\gamma = \gamma_{2k-1}(1) = \pi - \gamma_{2k}(1) \pmod{2\pi}$:

$$(5.31) \quad \begin{bmatrix} -\cos \gamma \cot \pi z + \sin \gamma & \cos \gamma \csc \pi z \\ -\cos \gamma \csc \pi z & \cos \gamma \cot \pi z + \sin \gamma \end{bmatrix}$$

which has determinant $= 1$ for $0 < \operatorname{Re} z < 1$. Hence on the r.h.s. of $\partial R_{1/p, I}$:

$$\det[\sigma(A)] = \prod_{j=1}^N 1 = 1.$$

To calculate the determinant of the matrix of principal symbols on $\partial R_{1/p, I}$, with $0 \leq t < 1$, we are again reduced to studying 2×2 blocks which arise from considering the operator near a vertex P_{2k-1} . Calling $i = 2k-1$, $\theta = \theta_i$, $\gamma_{i-1} = \gamma_{2k-2}(t)$, $\gamma_i = \gamma_{2k-1}(t)$ the matrix which comes from P_i is given by

$$(5.32) \quad \frac{-1}{\sin \pi z} \begin{bmatrix} \cos(\pi z + \gamma_{i-1}) & \cos((\pi - \theta)z + \theta - \gamma_{i-1}) \\ \cos((\pi - \theta)z + \theta - \gamma_i) & \cos(\pi z + \gamma_i) \end{bmatrix}$$

which has determinant

$$(5.33) \quad \frac{1}{2} \frac{\cos(2\pi z + 2\gamma) - \cos(2(\pi - \theta)z + 2\theta + 2\gamma)}{\sin^2 \pi z}$$

$$2\gamma = \gamma_{i-1} + \gamma_i.$$

The function in (5.33) may be expressed as

$$(5.34) \quad F_{\theta 2\gamma}(z) = \frac{\sin((2\pi - \theta)z + \theta + 2\gamma)\sin(\theta(1-z))}{\sin^2 \pi z}.$$

$F_{\theta 2\gamma}$ satisfies

$$(5.35) \quad F_{\theta 2\gamma}(1/p + i\infty) = -e^{-12\gamma}, \quad F_{\theta 2\gamma}(1/p - i\infty) = -e^{-12\gamma}$$

and has zeroes when

$$(5.36) \quad 1 - z = \frac{n\pi}{\theta}, \quad 1 - z = \frac{n\pi + 2\gamma}{2\pi - \theta}.$$

Hence

$$\det[\sigma(A)](t, 1/p + i\infty) = (-1)^N \prod_{j=1}^N e^{-1(\gamma_{2k-2}(t) + \gamma_{2k-1}(t))}$$

(5.37)

$$\det[\sigma(A)](t, 1/p - i\infty) = (-1)^N \prod_{j=1}^N e^{+1(\gamma_{2k-2}(t) + \gamma_{2k-1}(t))}.$$

Note that $\det[\sigma(A)](1, 1/p \pm i\infty) = 1$ by the continuity condition (5.24). We conclude that (5.40) is Fredholm on $L^p(\partial\Omega)$ iff $1/q = 1 - 1/p$ satisfies

$$\frac{1}{q} \neq \frac{n\pi}{\theta}, \quad \frac{1}{q} \neq \frac{n\pi + 2\gamma}{2\pi - \theta}$$

(5.38)

$$\theta = \theta_{2k-1}, \quad 2\gamma = \gamma_{2k-2}(0) + \gamma_{2k-1}(0), \quad k = 1, \dots, N.$$

If (5.38) is satisfied, the index of (5.30) on $L^p(\partial\Omega)$ can be readily calculated using the Index Theorem 4.2 and (5.37).

For an Index Theorem for the Oblique Derivative Problem in the case of a domain with smooth boundary, see Hormander [H]. We also refer the reader to the work of Costabel [Cos] who develops an algebra of singular integral operators on curves with corners which includes the double layer potential and proves an Index Theorem. His approach also allows weights such as t^β at the vertices of the polygon.

ACKNOWLEDGEMENTS

The first author was partially supported by the National Science Foundation and the Italian National Research Council, Gruppo G.N.A.F.A.

The second author was partially supported by the Italian National Research Council, Gruppo G.N.A.F.A., and the University of Illinois at Chicago.

BIBLIOGRAPHY

- Av1 - A. Avantiaggiati, Problemi al contorno in aperti di R^n con punti conici, Rend. del Sem. Mat e Fisico di Milano 47 (1979), 9-43.
- Av2 - A. Avantiaggiati, Sistemi di equazione a derivate parizale a coefficienti costante in due variabili e problemi ellitici, Ricerche Mat. 24 (1975), 203-279.
- B - L. Boutet de Monvel, Boundary value problems for pseudodifferential operators, Acta Math. 126 (1971), 11-51.
- CM - R.R. Coifman and Y. Meyer, Au dela des Opérateurs Pseudo-differentiels, Asterisque 57 (1978), Soc. Math. France, Paris.
- C - H.O. Cordes, Elliptic Pseudo-Differential Operators - An Abstract Theory, Lecture Notes in Mathematics 756 (1979), Springer, Berlin.
-

- CH - H.O. Cordes and E.A. Herman, Singular integral operators on a half-line, P.N.A.S.U.S.A. 56 (1966), 1668-1673.
- Cos - M. Costabel, Singular integral operators on curves with corners, Int. Eq. and Op. Theory 2 (1980), 323-349.
- DL - L. Diomeda and B. Lisena, Boundary value problems for Δ^2 in a square with L^p data. To appear.
- E - G.I. Eskin, General mixed boundary problems for elliptic differential equations, CIME Course on Pseudodifferential Operators with Applications, Ligouri, Naples, 1978.
- FJL - E.B. Fabes, M. Jodeit, Jr., and J.E. Lewis, Double layer potentials in domains with corners and edges, Indiana U. Math. J. 26 (1977), 95-114.
- G - P. Grisvard, Boundary Value Problems in Non-smooth Domains, University of Maryland Lecture Notes 19 (1980).
- H - L. Hörmander, Linear Partial Differential Operators, Springer, Berlin, 1963.
- K - V.A. Kondratiev, Boundary value problems for elliptic equations in domains with conical points, Trudy Mosk. Mat. Obch 16 (1967), 209-292 (Transactions Moscow Math. Soc. (1967), 227-313.)
- LP1 - J.E. Lewis and C. Parenti, Pseudodifferential operators and Hardy kernels on $L^p(\mathbb{R}^+)$, Ann. Sc. Norm. Sup. Pisa, Ser. IV, 7 (1980), 481-503.

- LP2 - J.E. Lewis and C. Parenti, Abstract singular parabolic equations, *Comm. P.D.E.* 7 (1982), 279-324.
- Mel - R.B. Melrose, Differential boundary value problems of principal type, Seminar on Singularities of Solutions of Linear Partial Differential Equations, Princeton Annals of Mathematics Studies 91 (1979).
- Mer - M. Merigot, Etude du probleme $\Delta u = f$ dans un polygone plan. Inegalites a priori, *Boll. U.M.I.* (4) 10 (1974), 577-597.
- Mey - Y. Meyer, Sharp estimates for pseudo-differential operators, Harmonic Analysis in Euclidean Spaces, A.M.S.P.S.P.M. 35 (part 2) (1979), 105-114.
- N1 - J. Nourrigat, Une classe d'opérateurs pseudo-différentiels a une variable, *C.R.A.S. Series A.* 283 (1976), 457-460.
- N2 - J. Nourrigat, Parametrices for a class of pseudodifferential operators of Fuchs type, preprint.
- Pe - I. Petrovsky, Lectures on Partial Differential Equations, Interscience, New York, 1954.
- Pl - B.A. Plamenevskii, An algebra of pseudo-differential operators in spaces with weight (in Russian), *Math. Sb.* 106 (1978), 297-320. (*Math. USSR-Sbornik* 24 (1978), 841-865).
- S1 - E. Shamir, Reduced Hilbert transforms and singular integral equations, *J. Analyse Math.* 12 (1964), 277-305.
-

- S2 - E. Shamir, Elliptic systems of singular
integral operators I, Trans. A.M.S.
127 (1967), 107-123.
- V - N.P. Vekua, Systems of Singular Integral
Equations, P. Noordhof, Groningen, 1967.

Revised November 1982