

# RESEARCH STATEMENT

## DR. RER. NAT. MARTIN BIES

By upbringing, I am a physicist, searching for solutions to **string theory**, which resemble the physics of our everyday experience. This is an involved geometric task and my research relies heavily on **algebraic geometry**, in particular **constructive and enumerative methods**. The resulting computations can often only be handled by modern **computer implementations**. Therefore, I have made major contributions to computer algebra systems, most recently the **OSCAR Computer Algebra System**.

**Outline** Section 1 provides an introduction to *string theory* and *F-theory*, which experts may skip. In section 2, I formulate a criterion to tell if solutions to *F-theory* contain certain exotic particles – *vector-like exotics* – which have not been observed in experiments. This leads to the quest for line bundles on certain curves subject to physical constraints, in particular their sheaf cohomologies. Section 3 summarizes my past contributions towards finding *F-theory* solutions without *vector-like exotics* and section 4 my current efforts.

## 1 Motivation

### 1.1 Why *string theory*?

It is experimentally established that four “rules” are necessary to describe all phenomena of our physical world: the *electromagnetic, weak and strong interaction* [1–3] and *gravity* [4]. The first three admit a unified description in the *standard model of particle physics* [5–7]. *General relativity* follows different rules and cannot be unified with the *standard model of particle physics* by means of perturbative quantum field theory [8]. The search for a unified description of *quantum gravity* has lead to a remarkable candidate for a theory of *quantum gravity*, namely *superstring theory* [9–12].

For reasons of consistency, *superstring theory* requires a 10 (real) dimensional spacetime  $\mathcal{S}$  and *supersymmetry* [9–12]. This is in contrast to our everyday experience of 4 spacetime directions. To account for this mismatch, one performs a compactification. We write

$$\mathcal{S} = \mathcal{E} \times_w \mathcal{M}_6,$$

where “ $w$ ” reminds us that this can in general be a warp-product and  $\mathcal{E}$  represents the 4 (real) dimensional spacetime of our everyday experience. The so-called *internal space*  $\mathcal{M}_6$  is, in the simplest case, a *compact Calabi-Yau manifold* of real dimension 6.<sup>1</sup>

Experiments have not found evidence for additional spacetime dimension. For this reason, we take  $\mathcal{M}_6$  sufficiently small. Similarly, *supersymmetry* has not been found in experiments – at least not yet. While recent experiments put a lot of stresses on

---

<sup>1</sup>In absence of fluxes, a necessary condition for conformal invariance of quantized *superstring theory* is  $0 = \alpha' R_{\mu\nu} + \frac{1}{2} (\alpha')^2 R_{\mu\lambda\rho\sigma} R_{\nu}^{\lambda\rho\sigma} + \dots$  [13], where  $R_{\mu\lambda\rho\sigma}$  is the *Riemann* tensor of the metric  $g_{\mu\nu}$  on  $\mathcal{S}$ . For the special case of a direct product and “small”  $\alpha'$ , i.e. in the SUGRA limit of *string theory*, this implies a Ricci flat  $\mathcal{M}_6$ . This is equivalent to the *Calabi-Yau* condition  $c_1(T_{\mathcal{M}_6}) = 0$  [14].

*supersymmetric theories*, we must not forget that the detection of the Higgs boson took a half-century, and that of *gravitational waves* an entire century. Therefore, I remain optimistic that *supersymmetry* can eventually be established at high energy scales and is broken at low energies. An exposition of mechanism for *supersymmetry* breaking in the context of my research can be found e.g. in [15, 16].

String phenomenology goes by the assumption that *superstring theory* is the correct unification of *gravity* and the *standard model of particle physics*. The goal is to find solutions to the string equations of motion such that upon compactification and *supersymmetry* breaking, only the observed 4-dimensional physics remains. To tell if a compactification represents the experimentally observed physics, among others the exact massless spectrum of this compactification must be computed.<sup>2</sup> This is what my research focuses on.

## 1.2 Why *F-theory*?

*Superstring theory* in 10 dimensions can be phrased in five consistent formulations [9, 10, 13]. We focus on so-called type IIB *superstring theory* for which the following are important:

- Dilaton  $\phi$ : A scalar field (massless bosonic string excitation) on the spacetime  $\mathcal{S}$ .
- D7-branes: 8 (real) dimensional, dynamic objects in the spacetime  $\mathcal{S}$ , which are electrically charged under the 8-form RR gauge potential  $C_8$ .

The dilaton  $\phi$  is of ample importance since  $\phi_0 := \lim_{x \rightarrow \infty} \phi(x)$  fixes the string coupling far away from D-branes via  $g_s = e^{\phi_0}$  [9–12]. D7-branes are crucial since they realize  $U(N)$  gauge theories [15]. The consistency of the field equation for the  $C_8$  gauge potential requires that the  $C_8$ -charges of all D7-branes in a type IIB compactification cancel. Therefore, non-trivial and stable compactifications with D7-branes must contain orientifold planes  $O7$ , which are negatively charged under  $C_8$  [9, 10]. To achieve this, the internal space  $\mathcal{M}_6$  must be modified by the operation of an involution  $\sigma$  via

$$\mathcal{M}_6 \rightarrow \mathcal{B}_6 := \mathcal{M}_6 / \sigma.$$

*Superstring theory* on  $\mathcal{B}_6$  is obtained by removing the orientation of the strings. To this end, one focuses on string excitations which are invariant under the operation of the parity operator  $\Omega$  and  $(-1)^{F_L}$ .<sup>3</sup> This defines the *orientifold theory* on  $\mathcal{B}_6$  [15, 17, 18].

While type IIB orientifold theories with D7-branes are phenomenologically desirable, the dilaton  $\phi$  then develops a profile which challenges the perturbative descriptions and often requires non-perturbative techniques [19]. This can be seen as the origin of *F-theory* – the exact description of strongly coupled type IIB *superstring theory* [20]. The key is to combine the dilaton  $\phi: \mathcal{E} \times_w \mathcal{B}_6 \rightarrow \mathbb{R}$  with the 0-form valued RR gauge field  $C_0: \mathcal{E} \times_w \mathcal{B}_6 \rightarrow \mathbb{R}$  to the axio-dilaton  $\tau$ :

$$\tau: \mathcal{E} \times_w \mathcal{B}_6 \rightarrow \mathbb{C}, \quad x^\mu \mapsto C_0(x^\mu) + ie^{-\phi(x^\mu)}.$$

<sup>2</sup>Those massless states are rendered massive by spontaneous symmetry breaking via the Higgs mechanism.

<sup>3</sup> $F_L$  is the number operator for left-handed, fermionic string excitations.

By *Lorentz invariance*,  $\tau$  is constant on  $\mathcal{E}$  and the section of a holomorphic  $\mathrm{SL}(2, \mathbb{Z})$  line bundle over  $\mathcal{B}_6$  [21, 22]. We interpret the value of  $\tau$  at  $x^\mu \in \mathcal{B}_6$  as the complex structure modulus of an elliptic curve  $\mathbb{C}_{1, \tau(x^\mu)}$ . A geometry which keeps track of the axio-dilaton profile is thus obtained by "attaching" the elliptic curve  $\mathbb{C}_{1, \tau(x^\mu)}$  at  $x^\mu \in \mathcal{B}_6$ . This leads to the study of elliptically fibered 4-folds  $\pi: Y_4 \rightarrow \mathcal{B}_6$  and fibre  $\mathbb{C}_{1, \tau(x^\mu)} := \pi^{-1}(x^\mu)$ . The geometry of  $Y_4$  contains information about the physics. The phenomenological implications are summarized e.g. in [23–26]. For non-trivial physics,  $Y_4$  must be singular. Since the analysis of smooth geometries is typically simpler, it is standard to resolve the singularities [27], most commonly by a crepant blowup resolution. For simplicity, I will not make a strict distinction between the singular  $Y_4$  and its resolved cousin  $\widehat{Y}_4$ .

An elliptic curve is topologically isomorphic to a torus surface with a marked point [28]. Hence, in considering an elliptically fibered 4-fold  $Y_4$ , there exists for each  $x^\mu \in \mathcal{B}_6$  a marked point  $P(x^\mu) \in \pi^{-1}(x^\mu) = \mathbb{C}_{1, \tau(x^\mu)}$  and we can consider  $s: \mathcal{B}_6 \rightarrow Y_4$ ,  $x^\mu \mapsto P(x^\mu)$ . For *F-theory* applications, we demand that this map is (at least) continuous, i.e.  $Y_4$  admits a (continuous) section  $s$ . While our applications focus on *F-theory with a section*, it is worth mentioning that *F-theory without section* has been described e.g. in [29–32].

### 1.3 Why *F-theory standard models*?

Enormous efforts have been undertaken to demonstrate the particle spectrum of the *standard model* from *string theory*. The earliest studies focus on the  $E_8 \times E_8$  heterotic string [33–40] and were later extended by intersecting branes models [41–48]. While these compactifications realize the gauge sector and chiral spectrum of the *standard model*, they are limited to the perturbative regime in the string coupling. Typically, they also suffer from vector-like exotics. The first globally consistent, perturbative MSSM constructions are [36, 37] (see [49, 50] for more details).

As *F-theory* describes strongly coupled IIB *superstring theory* and provides a coherent approach to analyze the relations among the geometry of  $Y_4$  and the 4-dimensional physics [20–22], efforts were undertaken to realize the *standard model* in *F-theory* [51–59]. These works have recently led to the discovery of the largest, currently-known, class of one quadrillion globally consistent *F-theory standard models (QSMs)* with gauge coupling unification and no chiral exotics [60]. I aim towards finding even better solutions and analyze the exact matter content to tell if unobserved vector-like particles are predicted.

## 2 Towards *F-theory vacua* without vector-like exotics

### 2.1 Counting exact massless spectra in *F-theory*

In [61], the study of massless spectra in *F-theory* was initiated. The matter fields relevant for our study localize along so-called *matter curves*  $C_{\mathbf{R}} \subset \mathcal{B}_6$  [62–64].<sup>4</sup> Mathematically,  $C_{\mathbf{R}}$

---

<sup>4</sup>The matter fields relevant for our study are, in type IIB language, the strings stretching between two D7-branes. Other matter fields stem from open strings on the same D7-brane and closed strings. In quantum field theory, particles are specified by a representation of a group  $G$  [65–68]. In *F-theory*, the representation  $\mathbf{R}$  of particles localized on  $C_{\mathbf{R}}$  is determined by the geometry of  $Y_4$  [23–26].

is a complex curve/Riemann surface [27]. The number of matter fields on  $C_{\mathbf{R}}$  is specified by gauge information. In part, this information is given by the 4-form valued gauge flux  $G_4 \in \Omega^4(Y_4)$ , which is induced by  $F$ -theory/M-theory duality from the M-Theory 3-form valued gauge potential  $C_3$  [23–26].  $G_4$  is a "field strength" (curvature) and can therefore not provide the complete gauge information (connection). In [69, 70],  $F$ -theory gauge fields were identified with elements of the Deligne cohomology  $H_D^4(Y_4, \mathbb{Z}(2))^5$ , which can be understood via the short exact sequence [75, 76]

$$0 \rightarrow J^2(Y_4) \rightarrow H_D^4(Y_4, \mathbb{Z}(2)) \xrightarrow{\hat{c}_2} H_{\mathbb{Z}}^{2,2}(Y_4) \rightarrow 0.$$

For  $A \in H_D^4(Y_4, \mathbb{Z}(2))$ , we interpret  $\hat{c}_2(A)$  as  $G_4$ -flux and the kernel of  $\hat{c}_2$  as the  $C_3$ -flat configurations.<sup>6</sup> We argue in [61] that  $A$  leads to a line bundle  $L_{\mathbf{R}}$  on the matter curve  $C_{\mathbf{R}}$ . In analogy with classical results on type IIB *superstring theory* [79–85], we then propose to count the massless matter on  $C_{\mathbf{R}}$  as follows:

- $\mathcal{N} = 1$  chiral multiplets are given by elements of  $H^0(C_{\mathbf{R}}, L_{\mathbf{R}} \otimes \sqrt{K_{C_{\mathbf{R}}}})$ .
- $\mathcal{N} = 1$  anti-chiral multiplets are given by elements of  $H^1(C_{\mathbf{R}}, L_{\mathbf{R}} \otimes \sqrt{K_{C_{\mathbf{R}}}})$ .<sup>7</sup>

$\sqrt{K_{C_{\mathbf{R}}}}$  is the spin bundle on  $C_{\mathbf{R}}$  which is fixed by the holomorphic embedding  $\iota: C_{\mathbf{R}} \hookrightarrow Y_4$  in a  $\mathcal{N} = 1$  supersymmetric configuration. In the following, I write  $\mathcal{L}_{\mathbf{R}} = L_{\mathbf{R}} \otimes \sqrt{K_{C_{\mathbf{R}}}}$ .

## 2.2 Absence of vector-like exotic particles

Experiments have found 3 types of leptons<sup>8</sup> [86]. To include them in an  $F$ -theory vacuum, there must be a matter curve  $C_{\mathbf{R}}$  which realizes the representation  $(\mathbf{1}, \mathbf{2})_{-1/2}$  of the leptons under the *standard model gauge group*  $\mathrm{SU}(3)_C \times \mathrm{SU}(2)_L \times \mathrm{U}(1)_Y$ . In addition, the  $F$ -theory gauge field  $A \in H_D^4(Y_4, \mathbb{Z}(2))$  must induce  $\mathcal{L}_{(\mathbf{1}, \mathbf{2})_{-1/2}} \in \mathrm{Pic}(C_{(\mathbf{1}, \mathbf{2})_{-1/2}})$  with

$$h^0(C_{(\mathbf{1}, \mathbf{2})_{-1/2}}, \mathcal{L}_{(\mathbf{1}, \mathbf{2})_{-1/2}}) = 3 + a, \quad h^1(C_{(\mathbf{1}, \mathbf{2})_{-1/2}}, \mathcal{L}_{(\mathbf{1}, \mathbf{2})_{-1/2}}) = a.$$

The chiral index – a topological invariant – is given by the Riemann-Roch theorem

$$\chi = \int_{C_{\mathbf{R}}} c_1(L_{\mathbf{R}}) = h^0(C_{\mathbf{R}}, L_{\mathbf{R}}) - h^1(C_{\mathbf{R}}, L_{\mathbf{R}}).$$

This famous result cannot predict the number  $a$  above. If  $a \neq 0$ , there is a non-trivial pair – known as *vector-like pair* – consisting of an element of  $h^0(C_{\mathbf{R}}, L_{\mathbf{R}})$  and  $h^1(C_{\mathbf{R}}, L_{\mathbf{R}})$ . *Vector-like pairs* have not been observed for leptons [86] and we must satisfy  $a = 0$ .

Variations of the complex structure of  $Y_4$  change  $\mathcal{L}_{\mathbf{R}}$ ,  $C_{\mathbf{R}}$  and, as is well-known e.g. in Brill-Noether theory [87] (see [88] for a more modern exposition and [89] for an earlier

<sup>5</sup>An equivalent approach can be found in [71–74].

<sup>6</sup>Equivalently, different gauge potentials in  $H_D^4(Y_4, \mathbb{Z}(2))$  differ by their Wilson lines [77, 78].

<sup>7</sup>A *supersymmetric field theory* only contains chiral fields. What we count are chiral superfields in the charge conjugate representation  $\bar{\mathbf{R}}$ . The terminology "anti-chiral" is inspired from low energy physics.

<sup>8</sup>These include the electron, which drives all our electric devices.

application of Brill-Noether theory in *F-theory*), can lead to jumps in  $h^i(C_{\mathbf{R}}, L_{\mathbf{R}})$ . This means that one could find  $h^i(C_{\mathbf{R}}, L_{\mathbf{R}}) = (3, 0)$  for generic and  $(4, 1)$  for special complex structure of  $Y_4$ . For leptons the generic outcome is desired. In contrast, a non-generic configuration is necessary for the Higgs field – the particle that gives mass to all other matter particles [90–92] and lead to the *Nobel Prize in Physics 2013*: A matter curve  $C_{\text{Higgs}}$  and  $\mathcal{L}_{\text{Higgs}} \in \text{Pic}(C_{\text{Higgs}})$  – induced by the above *F-theory* gauge field – with

$$h^0(C_{\text{Higgs}}, \mathcal{L}_{\text{Higgs}}) = h^1(C_{\text{Higgs}}, \mathcal{L}_{\text{Higgs}}) = 1.$$

By extending this task to all particles in the *standard model*, we arrive at the quest for *F-theory* realizations of *minimally supersymmetric standard models* (*F-theory* MSSMs).

## 3 Contributions towards *F-theory* MSSMs

### 3.1 F-theory and coherent sheaf cohomologies

In [61], we argued that for special gauge fields  $A$  the line bundle  $\mathcal{L}_R$  is the restriction of a bundle defined on a simple (typically toric) ambient space. While arithmetically favorable, this is not a generic feature of *F-theory* vacua. Consider for example *F-theory*  $\text{SU}(5)$  GUT models [63, 64, 93–96]. The gauge group  $\text{SU}(5)$  is realized on a D7-brane  $\mathcal{E} \times \Sigma^9$  and must be broken by a hypercharge flux. The latter is given by a holomorphic  $\text{U}(1)$  line bundle  $\mathcal{L}_Y$  on  $\mathcal{E} \times \Sigma$ . The absence of exotic massive particles requires<sup>10</sup>  $c_1(\mathcal{L}_Y) \in H^{1,1}(\Sigma) - \iota^* H^{1,1}(\mathcal{B}_6)$ . Consequently, the physics in  $\mathcal{E}$  observes the breaking  $\text{SU}(5) \rightarrow \text{SU}(3)_C \times \text{SU}(2)_L \times \text{U}(1)_Y$  iff  $\mathcal{L}_Y^\Sigma = \mathcal{L}_Y|_\Sigma$  is not pullback of a line bundle on  $\mathcal{B}_6$ . While our study in [97] focused on exactly this type of application, it could also show that in the presence of *matter surface fluxes* the massless spectra are typically encoded in non-pullback line bundles. Consequently, the computation of cohomologies of pullback line bundles – e.g. via the famous *cohomCalc* algorithm [98–102] – is insufficient to determine the massless spectra of most *F-theory* vacua. This makes the quest for *F-theory standard models* without vector-like exotics very challenging.

We can extend every line bundle  $\mathcal{L}_{\mathbf{R}}$  by “zero” outside of  $C_{\mathbf{R}}$  and obtain a coherent sheaf. On toric varieties, the cohomologies of coherent sheaves can be computed [103, 104]. In [105], I put forward an algorithm for such computations. It combines *cohomCalc* [98–102] with [106]. This algorithm was developed in collaboration with *Prof. Dr. Mohamed Barakat* at the *university of Siegen*, whose previous works establish a constructive approach to coherent sheaves on toric spaces is achieved by *Serre Quotient categories* [107–111]. The new algorithm is implemented in the **gap**-package [112]. It is written in the language of *categorical programming* (**CAP**) [113] and models coherent sheaves as objects of *Freyd categories* [114, 115]. The latter were first implemented in **CAP** in [116–118].

This new algorithm in [112] employs *cohomCalc* [98–102] to computed vanishing sets, which refine the semigroup  $K_{\text{sat}}$  introduced in [119] and used in [120] (see also [121] for

<sup>9</sup> $\mathcal{E}$  is the spacetime of our everyday experience and  $\Sigma \subset \mathcal{B}_6$  the complex surface wrapped in  $\mathcal{B}_6$ .

<sup>10</sup> $\mathcal{L}_Y$  defines an embedding  $\mathfrak{u}(1)_Y \hookrightarrow \mathfrak{su}(5)$ . The commutant of  $\mathfrak{u}(1)_Y$  does not obtain masses from the *Dirac-Born-Infeld*-action. A mass for the  $\text{U}(1)_Y$  gauge boson can be induced from the *Chern-Simons* action. An explicit study shows that this mass vanishes iff  $c_1(\mathcal{L}_Y) \in H^{1,1}(\Sigma) - \iota^* H^{1,1}(\mathcal{B}_6)$ .

more background). In [97, 105], we made use of these implementations and analyzed zero modes in *F-Theory* GUTs. While our results could not satisfy the experimental requirements, we gave the first examples of jumps in massless spectra of *F-theory vacua*.<sup>11</sup>

### 3.2 Promonoidal structures in Freyd categories

As outlined above, the computations necessary to determine massless spectra in *F-theory* are very challenging. The motivation for [123] was motivated by the desire for further improvements for the algorithm in [112]. The latter models a coherent sheaf  $\mathcal{F}$  on a toric space  $X_\Sigma$  by an object  $F$  in the Freyd category  $\mathbf{A}(S)$  [114, 115] associated to the Cox ring  $S$  of  $X_\Sigma$ . The sheaf cohomology  $H^0(X_\Sigma, \mathcal{F})$  is computed by  $H^0(X_\Sigma, \mathcal{F}) \cong \underline{\mathrm{Hom}}_S(I, F)_0$  where  $I$  is a suitable other object of  $\mathbf{A}(S)$ ,  $\underline{\mathrm{Hom}}_S$  denotes the so-called *Internal-Hom* of this category and the subscript 0 indicates the truncation to degree 0. The *Internal-Hom* is part of the *monoidal structure* of  $\mathbf{A}(S)$ . Consequently, the desire for quick determination of sheaf cohomologies sparks an interest in *monoidal structures* in *Freyd categories*.

In [123] we show that such *monoidal structures* derive from *pro-monoidal structures*. This insight allows us to formulate a constructive approach to tensor products in the category of finitely presented functors. We argue that our construction is equivalent to the restriction of the so-called *Day convolution of arbitrary functors* to finitely-presented functors [124, 125]. The software packages [116–118, 126] were subsequently remodeled and are now available in [127]. Accordingly, also [112] has been completely remodeled.

### 3.3 Systematics of jumps in exact massless spectra of *F-theory*

The geometry of  $Y_4$  and choice of gauge field  $A$  determine the matter curves  $C_{\mathbf{R}}$  and line bundles  $L_{\mathbf{R}}$ . As we vary the complex structure  $Y_4$ , it is expected that jumps in the massless spectra occur. Such jumps are crucial to match the experimental data – e.g. to accommodate the Higgs field at non-generic complex structure – we conducted a systematic study of such jumps in [128]. To simplify the computations, our study was inspired by an *F-theory*  $\mathrm{SU}(5)$ -GUT model. We considered a D7 GUT-brane wrapped on  $\mathcal{E} \times \Sigma$ , where the complex surface  $\Sigma \subset \mathcal{B}_6$  was chosen as a del Pezzo surface  $\Sigma = dP_3$ . We further focused on a gauge field  $A \in H_D^4(Y_4, \mathbb{Z}(2))$  such that  $L_{\mathbf{R}} \in \mathrm{Pic}(C_{\mathbf{R}})$  on the hypersurface matter curve  $C_{\mathbf{R}} \subseteq dP_3$  was the restriction of a line bundle on  $dP_3$  to  $C_{\mathbf{R}}$ .

This setup could not satisfy the experimental constraints, but is computationally rather simple. By employing the algorithms in [112], we therefore managed to generate a large database [129], which we then analyzed with decision trees – an interpretable machine learning tool. We found that the majority of jumps could be predicted from factorization of the curve  $C_{\mathbf{R}}$ . For example, we notice that once a  $\mathbb{P}^1$  factors-off, there is a good chance that the vector-like spectrum increases. Along this rational, we engineered an additional vector-like pair in a geometry which is – with the current technology – arithmetically almost inaccessible. Our database also contains jumps for which the curve remains smooth

---

<sup>11</sup>The study in [97, 105] was facilitated by modelling *F-theory* gauge potentials with elements of the Chow group  $\mathrm{CH}^2(Y_4, \mathbb{Z})$ . While putting the finishing touches to [97], we found that (local) anomalies in *F-Theory* follow similar rules and can be understood from Chow groups of  $Y_4$  [122].

and irreducible. Those are Brill-Noether jumps [87] – the line bundle divisor moves into a fairly non-generic configuration so that it admits additional sections.

A Koszul resolution of  $L_{\mathbf{R}}$  allowed us to analytically determine the cohomologies as the (co)kernel of a complex-structure valued matrix [128]. By analyzing this matrix, we stratified the parameter space of  $C_{\mathbf{R}}$  with respect to the massless spectra and verified that jumps arise from a combination of curve factorization and Brill-Noether jumps.

### 3.4 (Limit) Roots in the F-theory QSMs

Recently, one quadrillion ( $10^{15}$ ) *F-theory* vacua were found which have many attractive physical features for *standard models* (e.g. gauge coupling unification and no chiral exotics) [60].<sup>12</sup> The vector-like spectra for these compactifications were not identified in [60]. Since only the  $G_4$ -flux was specified – it is sufficient to determine the chiral index [51–53, 74, 94, 130–134] but does not uniquely determine the *F-theory* gauge field  $A$  – we only found the fairly non-trivial and necessary constraint that  $L_{\mathbf{R}}$  is a root bundle [135]. For the geometries in [60], the triple intersection number  $\overline{K}_{\mathcal{B}_6}^3$  of the anticanonical class of  $\mathcal{B}_6$  satisfies  $\overline{K}_{\mathcal{B}_6}^3 \in \{6, 10, 18, 30\}$  and there are five matter curves:<sup>13</sup>

curve	genus	root bundle constraint	$h^i$ constraint
$C_{(\mathbf{3},\mathbf{2})_{1/6}}$	$\frac{\overline{K}_{\mathcal{B}_6}^3+2}{2}$	$P_{(\mathbf{3},\mathbf{2})_{1/6}}^{\otimes 2\overline{K}_{\mathcal{B}_6}^3} = K_{(\mathbf{3},\mathbf{2})_{1/6}}^{\otimes (6+\overline{K}_{\mathcal{B}_6}^3)}$	$(3, 0)$
$C_{(\mathbf{1},\mathbf{2})_{-1/2}}$	$\frac{9\overline{K}_{\mathcal{B}_6}^3+2}{2}$	$P_{(\mathbf{1},\mathbf{2})_{-1/2}}^{\otimes 2\overline{K}_{\mathcal{B}_6}^3} = K_{(\mathbf{1},\mathbf{2})_{-1/2}}^{\otimes (4+\overline{K}_{\mathcal{B}_6}^3)} \otimes \mathcal{O}_{C_{(\mathbf{1},\mathbf{2})_{-1/2}}}(-30 \cdot Y_1)$	$(3, 0) \oplus (1, 1)$
$C_{(\overline{\mathbf{3}},\mathbf{1})_{-2/3}}$	$\frac{\overline{K}_{\mathcal{B}_6}^3+2}{2}$	$P_{(\overline{\mathbf{3}},\mathbf{1})_{-2/3}}^{\otimes 2\overline{K}_{\mathcal{B}_6}^3} = K_{(\overline{\mathbf{3}},\mathbf{1})_{-2/3}}^{\otimes (6+\overline{K}_{\mathcal{B}_6}^3)}$	$(3, 0)$
$C_{(\overline{\mathbf{3}},\mathbf{1})_{1/3}}$	$\frac{9\overline{K}_{\mathcal{B}_6}^3+2}{2}$	$P_{(\overline{\mathbf{3}},\mathbf{1})_{1/3}}^{\otimes 2\overline{K}_{\mathcal{B}_6}^3} = K_{(\overline{\mathbf{3}},\mathbf{1})_{1/3}}^{\otimes (4+\overline{K}_{\mathcal{B}_6}^3)} \otimes \mathcal{O}_{C_{(\overline{\mathbf{3}},\mathbf{1})_{1/3}}}(-30 \cdot Y_3)$	$(3, 0)$
$C_{(\mathbf{1},\mathbf{1})_1}$	$\frac{\overline{K}_{\mathcal{B}_6}^3+2}{2}$	$P_{(\mathbf{1},\mathbf{1})_1}^{\otimes 2\overline{K}_{\mathcal{B}_6}^3} = K_{(\mathbf{1},\mathbf{1})_1}^{\otimes (6+\overline{K}_{\mathcal{B}_6}^3)}$	$(3, 0)$

For a specific choice of  $\mathcal{B}_6$ , we argue in [135] that  $C_{(\mathbf{3},\mathbf{2})_{1/6}}$  admits  $2\overline{K}_{\mathcal{B}_6}^3$ -th root  $P_{(\mathbf{3},\mathbf{2})_{1/6}}$  of the  $(6 + \overline{K}_{\mathcal{B}_6}^3)$ -th power of the canonical bundle of  $C_{(\mathbf{3},\mathbf{2})_{1/6}}$  with cohomologies  $(3, 0)$ :

1. We deform  $C_{(\mathbf{3},\mathbf{2})_{1/6}}$  into a canonical nodal curve  $C_{(\mathbf{3},\mathbf{2})_{1/6}}^\bullet$ .<sup>14</sup>
2. An explicit construction proves the existence of a so-called *limit root*  $P_{(\mathbf{3},\mathbf{2})_{1/6}}^\bullet$  on  $C_{(\mathbf{3},\mathbf{2})_{1/6}}^\bullet$  with cohomologies  $(3, 0)$  [141] (see also [142–144]).
3. The number of global sections cannot increase along  $C_{(\mathbf{3},\mathbf{2})_{1/6}}^\bullet \rightarrow C_{(\mathbf{3},\mathbf{2})_{1/6}}$  by upper semicontinuity. It cannot decrease neither since it is already minimal on  $C_{(\mathbf{3},\mathbf{2})_{1/6}}^\bullet$ . Hence, the root  $P_{(\mathbf{3},\mathbf{2})_{1/6}}$  on  $C_{(\mathbf{3},\mathbf{2})_{1/6}}$  with cohomologies  $(3, 0)$  exists.

<sup>12</sup>See [52, 56, 58, 59] for earlier works in this *F-theory* program.

<sup>13</sup>The Higgs and lepton matter curves coincide in the models in [60].

<sup>14</sup> $\mathcal{B}_6$  is obtained from the desingularization of a toric K3-surface [136] (see also [104, 137]). By classical results such as [138, 139], the Picard lattice of said K3-surface is closely related to  $C_{(\mathbf{3},\mathbf{2})_{1/6}}^\bullet$  [140].

This last step fails for  $C_{(1,2)_{-1/2}}$  since we need a *non-minimal* number of global sections to realize the Higgs field. This sparks part of my current research efforts.

### 3.5 Enumeration of limit root bundles

The techniques in [141] allow constructing limit roots on nodal curves by solving certain combinatorics tasks. To gain deeper insights into the root bundle constraints outlined above, we have thus conducted a systematic study in [140]. A simple estimation shows that even in the simplest geometries, we expect about  $10^8$  root bundles. Indeed, the resulting combinatorics tasks are huge. To complete them in a finite time, we have therefore developed C++-code (available as part of [145]) which – after several performance optimizations and imposing simplifying assumptions – was able to complete this task.

A tremendous simplification arises because, the curve  $C_{(3,2)_{1/6}}^\bullet$  is identical for many of the geometries in [60], which are obtained from desingularizations of toric K3-surfaces [136, 137]. It then turns out the Picard lattice of the K3-surface in question is closely related to the dual graph of  $C_{(3,2)_{1/6}}^\bullet$ . This allowed us to scan over the majority over these quadrillion geometries [60, 146]. Our computations find the largest fraction of root bundles  $P_{(3,2)_{1/6}}$  with cohomologies  $(3, 0)$  for a particular family of spaces  $\mathcal{B}_6$ , namely the toric varieties obtained from fine regular star triangulations of the 8-th 3-dimensional reflexive polytope  $\Delta_8^\circ$  in the Kreuzer-Skarke list [137].<sup>15</sup> At the current state-of-affairs, these spaces are the most promising base spaces  $\mathcal{B}_6$  for *F-theory MSSMs*.

## 4 Current research efforts

My current and – to the extent that I can anticipate – future research efforts will focus on further elucidating the quest for *F-theory standard models* without vector-like exotics. I believe that we made a lot of progress in this direction over the last years, and I hope to find more insights in the foreseeable future. My current efforts are as follows.

**Equidimensional deformations** A key towards extending our work [135] to exactly one Higgs field is to understand by how much the cohomologies of limit root line bundles on  $C_{(1,2)_{-1/2}}^\bullet$  differ from those on  $C_{(1,2)_{-1/2}}$ . From the physics, we expect interactions among fields to happen in the nodes of  $C_{(1,2)_{-1/2}}^\bullet$ . This effect is accounted for by the complex structure valued mass matrix  $M$ . The physics predicts that the rank  $\text{rk}(M)$  tells the difference between the vector-like spectra on  $C_{(1,2)_{-1/2}}^\bullet$  and  $C_{(1,2)_{-1/2}}$ . The key challenges are to compute the mass matrix  $M$  as pioneered in [148] and recently extended in [149], and then to relate the physics prediction to the corresponding mathematics, e.g. Brill-Noether jumps [87] or *limit linear series* [150–152].

---

<sup>15</sup>Star means that every simplex in the triangulation contains the origin, fine that every lattice point of  $\Delta^\circ$  is used as ray generator and regular implies that the resulting variety is projective [103, 147].



**Counting limit root bundles on the Higgs curve** The genus of  $C_{(\mathbf{3},\mathbf{2})_{1/6}}$  is much smaller than  $C_{(\mathbf{1},\mathbf{2})_{-1/2}}$  for the *F-theory QSMs* [60]. For example, for bases  $\mathcal{B}_6$  with  $\overline{K}_{\mathcal{B}_6}^3 = 6$ , it holds  $g(C_{(\mathbf{3},\mathbf{2})_{1/6}}) = 4 < 28 = g(C_{(\mathbf{1},\mathbf{2})_{-1/2}})$ . Consequently, we have to analyze  $12^8$  limit roots on  $C_{(\mathbf{3},\mathbf{2})_{1/6}}^\bullet$  and  $12^{56}$  on  $C_{(\mathbf{1},\mathbf{2})_{-1/2}}^\bullet$  (cf. section 3.4). Clearly, the combinatorics task on  $C_{(\mathbf{1},\mathbf{2})_{-1/2}}$  is huge compared to  $C_{(\mathbf{3},\mathbf{2})_{1/6}}$ . This is the reason why, for simplicity, our analysis in [140] counted limit root line bundles on curves  $C_{(\mathbf{3},\mathbf{2})_{1/6}}$  with  $g = 4, 6$ . Still, the enumeration of limit root bundles on  $C_{(\mathbf{1},\mathbf{2})_{-1/2}}$  is crucial to gain further insights in the accommodation of the Higgs field in the *F-theory QSMs* [60]. Therefore, we have refined and extended our algorithm. Significant computational efforts have led to partial results in tandem with our findings in [140]. We are currently investigating further extensions of our computational powers to gain further insights.

**Brill-Noether theory for roots on nodal curves** Under simplifying assumptions, one can order nodal curves via their dual graphs from "simple" to "involved". Our algorithms in [145] approximate their Brill-Noether theory in that they find a lower bound to the number  $n(h^0)$  of root bundle with  $h^0$  global sections.<sup>16</sup> Generally speaking, these computations require a lot of computational resources. Inspired by applications to the F-theory QSMs [60], one may wonder if  $n(h^0)$  can be predicted/approximated from the dual graph of the nodal curve in question. It would be interesting to study this question – in spirit similar to [128] – driven both by insights from *machine learning tools* and *analytic/algebraic insights*.

**OSCAR Computer Algebra System** I have developed software for the computation of vector-like spectra since around 2015. My main focus was on the *ToricVarieties\_project* [145], which by now consists of more than 40,000 lines of code. I have also contributed to [111, 113, 155]. According to *github*, I have added/modified almost 1,000,000 lines of code. For comparison, the code for this L<sup>A</sup>T<sub>E</sub>X-document consists of less than 300 lines of code.

While [145] is a nice software collection, improvements are necessary. This project was mainly written in the programming language *gap* [156], which is – measured in time-scales of computer sciences – somewhat antique by now. For this reason, I have contributed to the computer algebra system *Oscar* [155] which uses the modern language of *Julia*. I am one of the main contributors on *toric geometry* and hope to refactor the entire functionality of the [145]. Thereby, this code should be modernized and subsequently available to a much broader audience.

---

<sup>16</sup>By classical results, roughly half of the spin bundles on a smooth curve have even and the other half odd number of global sections [153, 154]. We are trying to refine this classical statement.

## References

- [1] Sheldon L. Glashow, *Partial-symmetries of weak interactions*, *Nuclear Physics* **22**, no. 4 (1961) 579–588, URL [https://dx.doi.org/10.1016/0029-5582\(61\)90469-2](https://dx.doi.org/10.1016/0029-5582(61)90469-2). 1
- [2] Weinberg, Steven, *A Model of Leptons*, *Phys. Rev. Lett.* **19** (1967) 1264–1266, URL <http://dx.doi.org/10.1103/PhysRevLett.19.1264>. 1
- [3] Abdus Salam, *Weak and electromagnetic interactions*, pp. 244–254, URL [http://dx.doi.org/10.1142/9789812795915\\_0034](http://dx.doi.org/10.1142/9789812795915_0034). 1
- [4] Einstein, Albert, *Die Feldgleichungen der Gravitation*, chapter , pp. 88–92, John Wiley & Sons, Ltd, 2005, ISBN 9783527608959, URL <https://dx.doi.org/10.1002/3527608958.ch5>. 1
- [5] Halzen, F. and Martin, A.D. and John Wiley & Sons, *Quarks and Leptones: An Introductory Course in Modern Particle Physics*, Wiley, 1984, ISBN 9780471887416, URL <https://books.google.de/books?id=zwDvAAAAMAAJ>. 1
- [6] Griffiths, D., *Introduction to Elementary Particles*, Physics textbook, Wiley, 2008, ISBN 9783527618477, URL <https://books.google.de/books?id=Wb9DYrjcoKAC>. 1
- [7] Schwartz, M.D., *Quantum Field Theory and the Standard Model*, Quantum Field Theory and the Standard Model, Cambridge University Press, 2014, ISBN 9781107034730, URL <https://books.google.de/books?id=HbdEAgAAQBAJ>. 1
- [8] A. Shomer, *A Pedagogical explanation for the non-renormalizability of gravity* [0709.3555]. 1
- [9] Polchinski, J., *String Theory: Volume 1, An Introduction To The Bosonic String*, Cambridge monographs on mathematical physics, Cambridge University Press, 1998, ISBN 9781139457408, URL [http://books.google.de/books?id=jbM3t\\_usmX0C](http://books.google.de/books?id=jbM3t_usmX0C). 1, 2
- [10] Polchinski, J., *String Theory: Volume 2, Superstring Theory and Beyond*, Cambridge Monographs on Mathematical Physics, Cambridge University Press, 2005, ISBN 9780521672283, URL <http://books.google.de/books?id=tJtOMAEACAAJ>. 1, 2
- [11] Green, M.B. and Schwarz, J.H. and Witten, E., *Superstring Theory: Volume 1, Introduction*, Cambridge Monographs on Mathematical Physics, Cambridge University Press, 1988, ISBN 9780521357524, URL <http://books.google.co.in/books?id=ItVsHqjJo4gC>. 1, 2
- [12] Green, M.B. and Schwarz, J.H. and Witten, E., *Superstring Theory: Volume 2, Loop Amplitudes, Anomalies and Phenomenology*, Cambridge Monographs on Mathematical Physics, Cambridge University Press, 1987, ISBN 9780521357531, URL <http://books.google.de/books?id=Z-uz4svcl0QC>. 1, 2

- [13] Blumenhagen, R. and Lüst, D. and Theisen, S., *Basic Concepts of String Theory*, Theoretical and Mathematical Physics, Springer, 2012, ISBN 9783642294969, URL <http://books.google.de/books?id=-3PNFQn6AzcC>. 1, 2
- [14] Yau, Shing-Tung, *Calabi's conjecture and some new results in algebraic geometry*, *Proceedings of the National Academy of Sciences* **74**, no. 5 (1977) 1798–1799, URL <http://dx.doi.org/10.1073/pnas.74.5.1798>. 1
- [15] Blumenhagen, Ralph and Körs, Boris and Lüst, Dieter and Stieberger, Stephan, *Four-dimensional string compactifications with D-branes, orientifolds and fluxes*, *Physics Reports* **445**, no. 1-6 (2007) 1–193, URL <http://dx.doi.org/10.1016/j.physrep.2007.04.003>. 2
- [16] Ibáñez, L.E. and Uranga, A.M., *String Theory and Particle Physics: An Introduction to String Phenomenology*, Cambridge University Press, 2012, ISBN 9780521517522, URL <http://books.google.de/books?id=vAUUu6DpVkUC>. 2
- [17] F. G. Marchesano Buznego, *Intersecting D-brane models*, Ph.D. thesis, 5 2003, [hep-th/0307252]. 2
- [18] Cremades, D. and Ibáñez, L.E. and Marchesano, F., *Intersecting Brane Models of Particle Physics and the Higgs Mechanism*, *Journal of High Energy Physics* **2002**, no. 07 (2002) 022–022, URL <http://dx.doi.org/10.1088/1126-6708/2002/07/022>. 2
- [19] Brian R. Greene and Alfred Shapere and Cumrun Vafa and Shing-Tung Yau, *Stringy cosmic strings and noncompact Calabi-Yau manifolds*, *Nuclear Physics B* **337**, no. 1 (1990) 1–36, URL [http://dx.doi.org/10.1016/0550-3213\(90\)90248-C](http://dx.doi.org/10.1016/0550-3213(90)90248-C). 2
- [20] Vafa, Cumrun, *Evidence for F-theory*, *Nuclear Physics B* **469**, no. 3 (1996) 403–415, URL [http://dx.doi.org/10.1016/0550-3213\(96\)00172-1](http://dx.doi.org/10.1016/0550-3213(96)00172-1). 2, 3
- [21] Morrison, David R. and Vafa, Cumrun, *Compactifications of F-theory on Calabi-Yau threefolds. (I)*, *Nuclear Physics B* **473**, no. 1-2 (1996) 74–92, URL [http://dx.doi.org/10.1016/0550-3213\(96\)00242-8](http://dx.doi.org/10.1016/0550-3213(96)00242-8). 3
- [22] Morrison, David R. and Vafa, Cumrun, *Compactifications of F-theory on Calabi-Yau threefolds (II)*, *Nuclear Physics B* **476**, no. 3 (1996) 437–469, URL [http://dx.doi.org/10.1016/0550-3213\(96\)00369-0](http://dx.doi.org/10.1016/0550-3213(96)00369-0). 3
- [23] Denef, Frederik, *Les Houches Lectures on Constructing String Vacua*, *Les Houches* **87** (2008) 483–610, [0803.1194]. 3, 4
- [24] Weigand, Timo, *Lectures on F-theory compactifications and model building*, *Classical and Quantum Gravity* **27**, no. 21 (2010) 214004, URL <http://dx.doi.org/10.1088/0264-9381/27/21/214004>. 3, 4

- [25] M. Cvetič and L. Lin, *TASI Lectures on Abelian and Discrete Symmetries in F-theory*, *PoS TASI2017* (2018) 020, URL <http://dx.doi.org/10.22323/1.305.0020>. 3, 4
- [26] T. Weigand, *F-theory*, *PoS TASI2017* (2018) 016, URL <http://dx.doi.org/10.22323/1.305.0016>. 3, 4
- [27] Griffiths, P. and Harris, J., *Principles of Algebraic Geometry*, Wiley Classics Library, Wiley, 2011, ISBN 9781118030776, URL <http://books.google.de/books?id=Sny48qKdW40C>. 3, 4
- [28] Connell, Ian, *Elliptic Curve Handbook*, 1999, URL <https://webs.ucm.es/BUCM/mat/doc8354.pdf>. 3
- [29] Witten, Edward, *Non-perturbative superpotentials in string theory*, *Nuclear Physics B* **474**, no. 2 (1996) 343–360, URL [http://dx.doi.org/10.1016/0550-3213\(96\)00283-0](http://dx.doi.org/10.1016/0550-3213(96)00283-0). 3
- [30] Braun, Volker and Morrison, David R., *F-theory on genus-one fibrations*, *Journal of High Energy Physics* **2014**, no. 8, URL [http://dx.doi.org/10.1007/JHEP08\(2014\)132](http://dx.doi.org/10.1007/JHEP08(2014)132). 3
- [31] Morrison, David R. and Taylor, Washington, *Sections, multisections, and  $U(1)$  fields in F-theory*, *Journal of Singularities* **15** (2016) 126–149, URL <http://dx.doi.org/10.5427/jsing.2016.15g>. 3
- [32] Anderson, Lara B. and García-Etxebarria, Iñaki and Grimm, Thomas W. and Keitel, Jan, *Physics of F-theory compactifications without section*, *Journal of High Energy Physics* **2014**, no. 12, URL [http://dx.doi.org/10.1007/JHEP12\(2014\)156](http://dx.doi.org/10.1007/JHEP12(2014)156). 3
- [33] P. Candelas and Gary T. Horowitz and Andrew Strominger and Edward Witten, *Vacuum configurations for superstrings*, *Nuclear Physics B* **258** (1985) 46–74, URL [http://dx.doi.org/10.1016/0550-3213\(85\)90602-9](http://dx.doi.org/10.1016/0550-3213(85)90602-9). 3
- [34] Brian R. Greene and Kelley H. Kirklin and Paul J. Miron and Graham G. Ross, *A superstring-inspired standard model*, *Physics Letters B* **180**, no. 1 (1986) 69–76, URL [http://dx.doi.org/10.1016/0370-2693\(86\)90137-1](http://dx.doi.org/10.1016/0370-2693(86)90137-1). 3
- [35] Braun, Volker and He, Yang-Hui and Ovrut, Burt A. and Pantev, Tony, *A heterotic standard model*, *Physics Letters B* **618**, no. 1-4 (2005) 252–258, URL <http://dx.doi.org/10.1016/j.physletb.2005.05.007>. 3
- [36] Bouchard, Vincent and Donagi, Ron, *An  $SU(5)$  heterotic standard model*, *Physics Letters B* **633**, no. 6 (2006) 783–791, URL <http://dx.doi.org/10.1016/j.physletb.2005.12.042>. 3
- [37] Bouchard, Vincent and Cvetič, Mirjam and Donagi, Ron, *Tri-linear couplings in an heterotic minimal supersymmetric Standard Model*, *Nuclear Physics B* **745**, no. 1-2 (2006) 62–83, URL <http://dx.doi.org/10.1016/j.nuclphysb.2006.03.032>. 3

- [38] L. B. Anderson, J. Gray, Y.-H. He and A. Lukas, *Exploring positive monad bundles and a new heterotic standard model*, *Journal of High Energy Physics* **2010**, no. 2, URL [http://dx.doi.org/10.1007/JHEP02\(2010\)054](http://dx.doi.org/10.1007/JHEP02(2010)054). 3
- [39] *Two hundred heterotic standard models on smooth Calabi-Yau threefolds*, *Physical Review D* **84**, no. 10, URL <http://dx.doi.org/10.1103/PhysRevD.84.106005>. 3
- [40] L. B. Anderson, J. Gray, A. Lukas and E. Palti, *Heterotic line bundle standard models*, *Journal of High Energy Physics* **2012**, no. 6, URL [http://dx.doi.org/10.1007/JHEP06\(2012\)113](http://dx.doi.org/10.1007/JHEP06(2012)113). 3
- [41] M. Berkooz, M. R. Douglas and R. G. Leigh, *Branes intersecting at angles*, *Nuclear Physics B* **480**, no. 1-2 (1996) 265–278, URL [http://dx.doi.org/10.1016/S0550-3213\(96\)00452-X](http://dx.doi.org/10.1016/S0550-3213(96)00452-X). 3
- [42] G. Aldazabal, S. Franco, L. E. Ibáñez, R. Rabadan and A. M. Uranga,  *$D=4$  chiral string compactifications from intersecting branes*, *Journal of Mathematical Physics* **42**, no. 7 (2001) 3103–3126, URL <http://dx.doi.org/10.1063/1.1376157>. 3
- [43] G. Aldazabal, S. Franco, L. E. Ibáñez, R. Rabadan and A. M. Uranga, *Intersecting brane worlds*, *Journal of High Energy Physics* **2001**, no. 02 (2001) 047–047, URL <http://dx.doi.org/10.1088/1126-6708/2001/02/047>. 3
- [44] L. E. Ibáñez, F. Marchesano and R. Rabadan, *Getting just the standard model at intersecting branes*, *Journal of High Energy Physics* **2001**, no. 11 (2001) 002–002, URL <http://dx.doi.org/10.1088/1126-6708/2001/11/002>. 3
- [45] R. Blumenhagen, B. Kors, D. Lüst and T. Ott, *The Standard Model from stable intersecting brane world orbifolds*, *Nuclear Physics B* **616**, no. 1-2 (2001) 3–33, URL [http://dx.doi.org/10.1016/S0550-3213\(01\)00423-0](http://dx.doi.org/10.1016/S0550-3213(01)00423-0). 3
- [46] M. Cvetič, G. Shiu and A. M. Uranga, *Three-Family Supersymmetric Standardlike Models from Intersecting Brane Worlds*, *Physical Review Letters* **87**, no. 20, URL <http://dx.doi.org/10.1103/PhysRevLett.87.201801>. 3
- [47] M. Cvetič, G. Shiu and A. M. Uranga, *Chiral four-dimensional  $N=1$  supersymmetric type IIA orientifolds from intersecting  $D6$ -branes*, *Nuclear Physics B* **615**, no. 1-3 (2001) 3–32, URL [http://dx.doi.org/10.1016/S0550-3213\(01\)00427-8](http://dx.doi.org/10.1016/S0550-3213(01)00427-8). 3
- [48] R. Blumenhagen, M. Cvetič, P. Langacker and G. Shiu, *Toward realistic intersecting  $D$ -brane models*, *Annual Review of Nuclear and Particle Science* **55**, no. 1 (2005) 71–139, URL <http://dx.doi.org/10.1146/annurev.nucl.55.090704.151541>. 3
- [49] Gómez, Tomás L. and Lukic, Sergio and Sols, Ignacio, *Constraining the Kähler Moduli in the Heterotic Standard Model*, *Communications in Mathematical Physics* **276**, no. 1 (2007) 1–21, URL <http://dx.doi.org/10.1007/s00220-007-0338-8>. 3

- [50] V. Bouchard and R. Donagi, *On heterotic model constraints*, *Journal of High Energy Physics* **2008**, no. 08 (2008) 060–060, URL <http://dx.doi.org/10.1088/1126-6708/2008/08/060>. 3
- [51] T. W. Grimm and H. Hayashi, *F-theory fluxes, chirality and Chern-Simons theories*, *Journal of High Energy Physics* **2012**, no. 3, URL [http://dx.doi.org/10.1007/JHEP03\(2012\)027](http://dx.doi.org/10.1007/JHEP03(2012)027). 3, 7
- [52] Krause, Sven and Mayrhofer, Christoph and Weigand, Timo, *G<sub>4</sub>-flux, chiral matter and singularity resolution in F-theory compactifications*, *Nuclear Physics B* **858**, no. 1 (2012) 1–47, URL <http://dx.doi.org/10.1016/j.nuclphysb.2011.12.013>. 3, 7
- [53] S. Krause, C. Mayrhofer and T. Weigand, *Gauge Fluxes in F-theory and Type IIB Orientifolds*, *Journal of High Energy Physics* **2012**, no. 8, URL [http://dx.doi.org/10.1007/JHEP08\(2012\)119](http://dx.doi.org/10.1007/JHEP08(2012)119). 7
- [54] V. Braun, T. W. Grimm and J. Keitel, *Geometric Engineering in Toric F-Theory and GUTs with U(1) Gauge Factors*, *Journal of High Energy Physics* **2013**, no. 12, URL [http://dx.doi.org/10.1007/JHEP12\(2013\)069](http://dx.doi.org/10.1007/JHEP12(2013)069). 3
- [55] M. Cvetič, A. Grassi, D. Klevers and H. Piragua, *Chiral four-dimensional F-theory compactifications with SU(5) and multiple U(1)-factors*, *Journal of High Energy Physics* **2014**, no. 4, URL [http://dx.doi.org/10.1007/JHEP04\(2014\)010](http://dx.doi.org/10.1007/JHEP04(2014)010). 3
- [56] M. Cvetič, D. Klevers, D. K. M. Peña, P.-K. Oehlmann and J. Reuter, *Three-family particle physics models from global F-theory compactifications*, *Journal of High Energy Physics* **2015**, no. 8, URL [http://dx.doi.org/10.1007/JHEP08\(2015\)087](http://dx.doi.org/10.1007/JHEP08(2015)087). 3, 7
- [57] L. Lin, C. Mayrhofer, O. Till and T. Weigand, *Fluxes in F-theory Compactifications on Genus-One Fibrations*, *Journal of High Energy Physics* **2016**, no. 1, URL [http://dx.doi.org/10.1007/JHEP01\(2016\)098](http://dx.doi.org/10.1007/JHEP01(2016)098). 3
- [58] L. Lin and T. Weigand, *G<sub>4</sub>-flux and standard model vacua in F-theory*, *Nuclear Physics B* **913** (2016) 209–247, URL <http://dx.doi.org/10.1016/j.nuclphysb.2016.09.008>. 3, 7
- [59] M. Cvetič, L. Lin, M. Liu and P.-K. Oehlmann, *An F-theory Realization of the Chiral MSSM with  $\mathbb{Z}_2$ -Parity*, *Journal of High Energy Physics* **2018**, no. 9, URL [http://dx.doi.org/10.1007/JHEP09\(2018\)089](http://dx.doi.org/10.1007/JHEP09(2018)089). 3, 7
- [60] M. Cvetič, J. Halverson, L. Lin, M. Liu and J. Tian, *Quadrillion F-Theory Compactifications with the Exact Chiral Spectrum of the Standard Model*, *Physical Review Letters* **123**, no. 10, URL <http://dx.doi.org/10.1103/PhysRevLett.123.101601>. 3, 7, 8, 9

- [61] M. Bies, C. Mayrhofer, C. Pehle and T. Weigand, *Chow groups, Deligne cohomology and massless matter in F-theory* [1402.5144]. 3, 4, 5
- [62] S. H. Katz and C. Vafa, *Matter from geometry*, *Nuclear Physics B* **497**, no. 1-2 (1997) 146–154, URL [http://dx.doi.org/10.1016/S0550-3213\(97\)00280-0](http://dx.doi.org/10.1016/S0550-3213(97)00280-0). 3
- [63] C. Beasley, J. J. Heckman and C. Vafa, *GUTs and exceptional branes in F-theory — I*, *Journal of High Energy Physics* **2009**, no. 01 (2009) 058–058, URL <http://dx.doi.org/10.1088/1126-6708/2009/01/058>. 3, 5
- [64] Donagi, Ron and Wijnholt, Martijn, *Gluing Branes I*, *Journal of High Energy Physics* **2013**, no. 05 (2013) 068–068, URL [http://dx.doi.org/10.1007/JHEP05\(2013\)068](http://dx.doi.org/10.1007/JHEP05(2013)068). 3, 5
- [65] Peskin, M.E. and Schroeder, D.V., *An Introduction To Quantum Field Theory*, Frontiers in Physics, Avalon Publishing, 1995, ISBN 9780813345437, URL <https://books.google.de/books?id=EVeNNcslvX0C>. 3
- [66] Weinberg, S., *The Quantum Theory of Fields*, Cambridge University Press, 1995, ISBN 9780521550017, URL [https://books.google.de/books?id=doeDB3\\_WLvwc](https://books.google.de/books?id=doeDB3_WLvwc). 3
- [67] Weinberg, S., *The Quantum Theory of Fields: Volume 2, Modern Applications*, Cambridge University Press, 1996, ISBN 9781139643252, URL <https://books.google.de/books?id=J2EhAwAAQBAJ>. 3
- [68] Weinberg, S., *The Quantum Theory of Fields: Volume 3, Supersymmetry*, Cambridge University Press, 2005, ISBN 9781139643436, URL <https://books.google.de/books?id=QMkgAwAAQBAJ>. 3
- [69] R. Donagi, *Heterotic / F theory duality: ICMP lecture*, in *12th International Congress of Mathematical Physics (ICMP 97)*, pp. 206–213, 2 1998, [hep-th/9802093]. 4
- [70] Curio, Gottfried and Donagi, Ron, *Moduli in  $N = 1$  heterotic/F-theory duality*, *Nuclear Physics B* **518**, no. 3 (1998) 603–631, URL [http://dx.doi.org/10.1016/S0550-3213\(98\)00185-0](http://dx.doi.org/10.1016/S0550-3213(98)00185-0). 4
- [71] Diaconescu, Emanuel and Freed, Daniel S. and Moore, Gregory, *The M-theory 3-form and E8 gauge theory*, p. 44–88, London Mathematical Society Lecture Note Series, Cambridge University Press, 2007, URL <http://dx.doi.org/10.1017/CB09780511721489.005>. 4
- [72] Freed, Daniel S. and Moore, Gregory W., *Setting the Quantum Integrand of M-Theory*, *Communications in Mathematical Physics* **263**, no. 1 (2006) 89–132, URL <http://dx.doi.org/10.1007/s00220-005-1482-7>. 4

- [73] Moore, Gregory W., *Anomalies, Gauss laws, and Page charges in M-theory*, *Comptes Rendus Physique* **6**, no. 2 (2005) 251–259, URL <http://dx.doi.org/10.1016/j.crhy.2004.12.005>. 4
- [74] Intriligator, Kenneth and Jockers, Hans and Mayr, Peter and Morrison, David R. and Plesser, M. Ronen, *Conifold transitions in M-theory on Calabi-Yau fourfolds with background fluxes*, *Advances in Theoretical and Mathematical Physics* **17**, no. 3 (2013) 601–699, URL <http://dx.doi.org/10.4310/ATMP.2013.v17.n3.a2>. 4, 7
- [75] Eisenbud, David and Harris, Joe, *3264 and All That: A Second Course in Algebraic Geometry*, Cambridge University Press, 2016, URL <http://dx.doi.org/10.1017/CB09781139062046>. 4
- [76] Fulton, W., *Intersection Theory*, *Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge / A Series of Modern Surveys in Mathematics*, Springer Berlin Heidelberg, 2013, ISBN 9783662024218, URL <https://books.google.de/books?id=gCXsCAAAQBAJ>. 4
- [77] Greiner, Sebastian and Grimm, Thomas W., *On Mirror Symmetry for Calabi-Yau Fourfolds with Three-Form Cohomology*, *Journal of High Energy Physics* **2016**, no. 9, URL [http://dx.doi.org/10.1007/JHEP09\(2016\)073](http://dx.doi.org/10.1007/JHEP09(2016)073). 4
- [78] Greiner, Sebastian and Grimm, Thomas W., *Three-form periods on Calabi-Yau fourfolds: Toric hypersurfaces and F-theory applications*, *Journal of High Energy Physics* **2017**, no. 5, URL [http://dx.doi.org/10.1007/JHEP05\(2017\)151](http://dx.doi.org/10.1007/JHEP05(2017)151). 4
- [79] E. Witten, *Topological sigma models*, *Communications in Mathematical Physics* **118**, no. 3 (1988) 411–449, URL <https://dx.doi.org/10.1007/BF01466725>. 4
- [80] Witten, Edward, *Mirror manifolds and topological field theory*, *AMS/IP Stud. Adv. Math.* **9** (1991) 121–160, [hep-th/9112056]. 4
- [81] B. R. Greene, *String theory on Calabi-Yau manifolds*, in *Theoretical Advanced Study Institute in Elementary Particle Physics (TASI 96): Fields, Strings, and Duality*, pp. 543–726, 6 1996, [hep-th/9702155]. 4
- [82] Sharpe, Eric, *D-branes, derived categories, and Grothendieck groups*, *Nuclear Physics B* **561**, no. 3 (1999) 433–450, URL [http://dx.doi.org/10.1016/S0550-3213\(99\)00535-0](http://dx.doi.org/10.1016/S0550-3213(99)00535-0). 4
- [83] Katz, Sheldon and Sharpe, Eric, *D-branes, open string vertex operators, and Ext groups*, *Advances in Theoretical and Mathematical Physics* **6**, no. 6 (2002) 979–1030, URL <http://dx.doi.org/10.4310/ATMP.2002.v6.n6.a1>. 4
- [84] E. Sharpe, *Lectures on D-branes and sheaves*, 7 2003, [hep-th/0307245]. 4
- [85] P. S. Aspinwall, *D-branes on Calabi-Yau manifolds*, *Progress In String Theory* pp. 1–152, URL [http://dx.doi.org/10.1142/9789812775108\\_0001](http://dx.doi.org/10.1142/9789812775108_0001). 4



- [86] Zyla, P.A. and others, *Review of Particle Physics*, *PTEP* **2020**, no. 8 (2020) 083C01, URL <http://dx.doi.org/10.1093/ptep/ptaa104>. 4
- [87] Brill, Alexander and Noether, Max, *Ueber die algebraischen Functionen und ihre Anwendung in der Geometrie*, *Mathematische Annalen* **7**, no. 2 (1874) 269–310, URL <http://dx.doi.org/10.1007/BF02104804>. 4, 7, 8
- [88] D. Eisenbud, M. Green and J. Harris, *Cayley-Bacharach theorems and conjectures*, *Bulletin of the American Mathematical Society* **33**, no. 03 (1996) 295–325, URL <http://dx.doi.org/10.1090/s0273-0979-96-00666-0>. 4
- [89] Watari, Taizan, *Vector-like pairs and Brill–Noether theory*, *Physics Letters B* **762** (2016) 145–150, URL <http://dx.doi.org/10.1016/j.physletb.2016.09.006>. 4
- [90] Englert, F. and Brout, R., *Broken Symmetry and the Mass of Gauge Vector Mesons*, *Phys. Rev. Lett.* **13** (1964) 321–323, URL <http://dx.doi.org/10.1103/PhysRevLett.13.321>. 5
- [91] P.W. Higgs, *Broken symmetries, massless particles and gauge fields*, *Physics Letters* **12**, no. 2 (1964) 132–133, URL [https://dx.doi.org/10.1016/0031-9163\(64\)91136-9](https://dx.doi.org/10.1016/0031-9163(64)91136-9). 5
- [92] Higgs, Peter W., *Broken Symmetries and the Masses of Gauge Bosons*, *Phys. Rev. Lett.* **13** (1964) 508–509, URL <https://dx.doi.org/10.1103/PhysRevLett.13.508>. 5
- [93] Beasley, Chris and Heckman, Jonathan J and Vafa, Cumrun, *GUTs and exceptional branes in F-theory — II. Experimental predictions*, *Journal of High Energy Physics* **2009**, no. 01 (2009) 059–059, URL <http://dx.doi.org/10.1088/1126-6708/2009/01/059>. 5
- [94] Donagi, Ron and Wijnholt, Martijn, *Model building with F-theory*, *Advances in Theoretical and Mathematical Physics* **15** (2011) 1237 – 1317, URL <http://dx.doi.org/10.4310/ATMP.2011.v15.n5.a2>. 5, 7
- [95] Donagi, Ron and Wijnholt, Martijn, *Breaking GUT groups in F-theory*, *Advances in Theoretical and Mathematical Physics* **15** (2008) 1523 – 1603, URL <http://dx.doi.org/10.4310/ATMP.2011.v15.n6.a1>. 5
- [96] Donagi, Ron and Wijnholt, Martijn, *Gluing Branes II: Flavour Physics and String Duality*, *Journal of High Energy Physics* **2013** (2011) , URL [http://dx.doi.org/10.1007/JHEP05\(2013\)092](http://dx.doi.org/10.1007/JHEP05(2013)092). 5
- [97] Bies, Martin and Mayrhofer, Christoph and Weigand, Timo, *Gauge backgrounds and zero-mode counting in F-theory*, *Journal of High Energy Physics* **2017**, no. 11, URL [http://dx.doi.org/10.1007/JHEP11\(2017\)081](http://dx.doi.org/10.1007/JHEP11(2017)081). 5, 6

- [98] Blumenhagen, Ralph and Jurke, Benjamin and Rahn, Thorsten and Roschy, Helmut, *Cohomology of line bundles: A computational algorithm*, *Journal of Mathematical Physics* **51**, no. 10 (2010) 103525, URL <http://dx.doi.org/10.1063/1.3501132>. 5, 18
- [99] *cohomCalc package*, Download link, 2010, URL <http://wwwth.mppmu.mpg.de/members/blumenha/cohomcalc/>, high-performance line bundle cohomology computation based on [98]. 5
- [100] Jow, Shin-Yao, *Cohomology of toric line bundles via simplicial Alexander duality*, *Journal of Mathematical Physics* **52**, no. 3 (2011) 033506, URL <http://dx.doi.org/10.1063/1.3562523>. 5
- [101] Roschy, Helmut and Rahn, Thorsten, *Cohomology of line bundles: Proof of the algorithm*, *Journal of Mathematical Physics* **51**, no. 10 (2010) 103520, URL <http://dx.doi.org/10.1063/1.3501135>. 5
- [102] Blumenhagen, Ralph and Jurke, Benjamin and Rahn, Thorsten and Roschy, Helmut, *Cohomology of line bundles: Applications*, *Journal of Mathematical Physics* **53**, no. 1 (2012) 012302, URL <http://dx.doi.org/10.1063/1.3677646>. 5
- [103] Fulton, W., *Introduction to Toric Varieties*, Annals of mathematics studies, Princeton University Press, 1993, ISBN 9780691000497, URL <http://books.google.de/books?id=CbmaGqCRbhUC>. 5, 8
- [104] Cox, D.A. and Katz, S., *Mirror Symmetry and Algebraic Geometry*, Mathematical surveys and monographs, American Mathematical Society, 1999, ISBN 9780821821275, URL <http://dx.doi.org/http://dx.doi.org/10.1090/surv/068>. 5, 7
- [105] Bies, Martin, *Cohomologies of coherent sheaves and massless spectra in F-theory*, Ph.D. thesis, Heidelberg U., 2018, URL <http://dx.doi.org/10.11588/heidok.00024045>. 5, 6
- [106] Smith, Gregory G., *Computing Global Extension Modules*, *Journal of Symbolic Computation* **29**, no. 4-5 (2000) 729–746, URL <http://dx.doi.org/10.1006/jSCO.1999.0399>. 5
- [107] Barakat, Mohamed and Lange-Hegermann, Markus, *On the Ext-computability of Serre quotient categories*, *Journal of Algebra* **420** (2014) 333–349, URL <http://dx.doi.org/10.1016/j.jalgebra.2014.08.004>. 5
- [108] Barakat, Mohamed and Lange-Hegermann, Markus, *On monads of exact reflective localizations of Abelian categories*, *Homology, Homotopy and Applications* **15**, no. 2 (2013) 145–151, URL <http://dx.doi.org/10.4310/HHA.2013.v15.n2.a8>. 5
- [109] Barakat, Mohamed and Lange-Hegermann, Markus, *An axiomatic setup for algorithmic homological algebra and an alternative approach to localization*, *J. Algebra Appl.*

- 10**, no. 2 (2011) 269–293, URL <http://dx.doi.org/10.1142/S0219498811004562>.  
5
- [110] Barakat, Mohamed and Lange-Hegermann, Markus, *Characterizing Serre quotients with no section functor and applications to coherent sheaves*, *Appl. Categor. Struct.* **22**, no. 3 (2014) 457–466, URL <http://dx.doi.org/10.1007/s10485-013-9314-y>.  
5
- [111] The homalg project authors, *The homalg project – Algorithmic Homological Algebra*, (<https://homalg-project.github.io/>), 2003–2021. 5, 9
- [112] M. Bies, *SheafCohomologyOnToricVarieties*, ([https://github.com/homalg-project/ToricVarieties\\_project/tree/master/SheafCohomologyOnToricVarieties](https://github.com/homalg-project/ToricVarieties_project/tree/master/SheafCohomologyOnToricVarieties)), 2019–2021. 5, 6
- [113] Gutsche, Sebastian and Skartsæterhagen, Øystein and Posur, Sebastian, *The CAP project – Categories, Algorithms, Programming*, ([http://homalg-project.github.io/CAP\\_project](http://homalg-project.github.io/CAP_project)), 2013–2021. 5, 9
- [114] Freyd, Peter, *Representations in abelian categories*, in *Proc. Conf. Categorical Algebra (La Jolla, Calif., 1965)*, pp. 95–120, Springer, New York, 1966, URL [http://dx.doi.org/10.1007/978-3-642-99902-4\\_4](http://dx.doi.org/10.1007/978-3-642-99902-4_4). 5, 6
- [115] Beligiannis, Apostolos, *On the Freyd categories of an additive category.*, *Homology, Homotopy and Applications* **2** (2000) 147–185, URL <http://dx.doi.org/10.4310/HHA.2000.v2.n1.a11>. 5, 6
- [116] M. Bies, *CAPCategoryOfProjectiveGradedModules*, (<https://github.com/HereAround/CAPCategoryOfProjectiveGradedModules>), 2017. 5, 6
- [117] M. Bies, *CAPPresentationCategory*, (<https://github.com/HereAround/CAPPresentationCategory>), 2017. 5, 6
- [118] M. Bies, *PresentationsByProjectiveGradedModules*, (<https://github.com/HereAround/PresentationsByProjectiveGradedModules>), 2017. 5, 6
- [119] MacLagan, D. and Smith, G. G., *Multigraded Castelnuovo-Mumford Regularity*, *Journal für die reine und angewandte Mathematik (Crelles Journal)* **2004**, no. 571, URL <http://dx.doi.org/10.1515/crll.2004.040>. 5
- [120] Barakat, Mohamed and Coquand, Thierry and Quadrat, Alban, *Mini-Workshop: Constructive Homological Algebra with Applications to Coherent Sheaves and Control Theory*, *Oberwolfach Reports* **10** (2013) 1495–1531, URL <http://dx.doi.org/10.4171/OWR/2013/25>. 5
- [121] Eisenbud, D. and Mustata, M. and Stillman, M., *Cohomology on Toric Varieties and Local Cohomology with Monomial Supports*, *J. Symb. Comput.* **29** (2000) 583–600, URL <http://dx.doi.org/10.1006/jSCO.1999.0326>. 5

- [122] M. Bies, C. Mayrhofer and T. Weigand, *Algebraic Cycles and Local Anomalies in F-Theory*, *Journal of High Energy Physics* **2017**, no. 11, URL [http://dx.doi.org/10.1007/JHEP11\(2017\)100](http://dx.doi.org/10.1007/JHEP11(2017)100). 6
- [123] Bies, Martin and Posur, Sebastian, *Tensor products of finitely presented functors*, *Journal of Algebra and Its Applications* p. 2250186, URL <http://dx.doi.org/10.1142/s0219498822501869>. 6
- [124] Day, Brian, *On closed categories of functors*, in *Reports of the Midwest Category Seminar, IV*, Lecture Notes in Mathematics, Vol. 137, pp. 1–38, Springer, Berlin, 1970, URL <http://dx.doi.org/10.1007/BFb0060438>. 6
- [125] Day, Brian, *On closed categories of functors. II*, in *Category Seminar (Proc. Sem., Sydney, 1972/1973)*, pp. 20–54. Lecture Notes in Math., Vol. 420, Springer, Berlin, 1974, URL <http://dx.doi.org/10.1007/BFb0063098>. 6
- [126] M. Bies, *TruncationsOfPresentationsByProjectiveGradedModules*, (<https://github.com/HereAround/TruncationsOfPresentationsByProjectiveGradedModules>), 2017. 6
- [127] M. Bies and S. Posur, *FreydCategories*, ([https://github.com/homalg-project/CAP\\_project/tree/master/FreydCategoriesForCAP](https://github.com/homalg-project/CAP_project/tree/master/FreydCategoriesForCAP)), 2019-2021. 6
- [128] M. Bies, M. Cvetič, R. Donagi, L. Lin, M. Liu and F. Ruehle, *Machine Learning and Algebraic Approaches towards Complete Matter Spectra in 4d F-theory*, *Journal of High Energy Physics* **2021**, no. 1, URL [http://dx.doi.org/10.1007/JHEP01\(2021\)196](http://dx.doi.org/10.1007/JHEP01(2021)196). 6, 7, 9
- [129] M. Bies, M. Cvetič, R. Donagi, L. Lin, M. Liu and F. Ruehle, *Database*, (<https://github.com/Learning-line-bundle-cohomology>), 2020. 6
- [130] H. Hayashi, R. Tatar, Y. Toda, T. Watari and M. Yamazaki, *New Aspects of Heterotic-F Theory Duality*, *Nuclear Physics B* **806**, no. 1-2 (2009) 224–299, URL <http://dx.doi.org/10.1016/j.nuclphysb.2008.07.031>. 7
- [131] Donagi, Ron and Wijnholt, Martijn, *Higgs Bundles and UV Completion in F-Theory*, *Commun. Math. Phys.* **326** (2014) 287–327, URL <http://dx.doi.org/10.1007/s00220-013-1878-8>. 7
- [132] Heckman, Jonathan J., *Particle Physics Implications of F-Theory*, *Annual Review of Nuclear and Particle Science* **60**, no. 1 (2010) 237–265, URL <http://dx.doi.org/10.1146/annurev.nucl.012809.104532>. 7
- [133] Marsano, Joseph and Schäfer-Nameki, Sakura, *Yukawas, G-flux, and spectral covers from resolved Calabi-Yau's*, *Journal of High Energy Physics* **2011**, no. 11, URL [http://dx.doi.org/10.1007/JHEP11\(2011\)098](http://dx.doi.org/10.1007/JHEP11(2011)098). 7

- [134] Braun, A.P. and Collinucci, A. and Valandro, R., *G-flux in F-theory and algebraic cycles*, *Nuclear Physics B* **856**, no. 2 (2012) 129–179, URL <http://dx.doi.org/10.1016/j.nuclphysb.2011.10.034>. 7
- [135] Bies, Martin and Cvetič, Mirjam and Donagi, Ron and Liu, Muyang and Ong, Marielle, *Root bundles and towards exact matter spectra of F-theory MSSMs*, *Journal of High Energy Physics* **2021**, no. 9, URL [http://dx.doi.org/10.1007/JHEP09\(2021\)076](http://dx.doi.org/10.1007/JHEP09(2021)076). 7, 8
- [136] V. V. Batyrev, *Dual polyhedra and mirror symmetry for Calabi-Yau hypersurfaces in toric varieties*, *J. Alg. Geom.* **3** (1994) 493–545, [alg-geom/9310003]. 7, 8
- [137] M. Kreuzer and H. Skarke, *Classification of reflexive polyhedra in three-dimensions*, *Adv. Theor. Math. Phys.* **2** (1998) 853–871, URL <http://dx.doi.org/10.4310/ATMP.1998.v2.n4.a5>. 7, 8
- [138] E. Perevalov and H. Skarke, *Enhanced gauged symmetry in type II and F theory compactifications: Dynkin diagrams from polyhedra*, *Nuclear Physics B* **505**, no. 3 (1997) 679–700, URL [http://dx.doi.org/10.1016/S0550-3213\(97\)00477-X](http://dx.doi.org/10.1016/S0550-3213(97)00477-X). 7
- [139] F. Rohsiepe, *Lattice polarized toric K3 surfaces* [hep-th/0409290]. 7
- [140] M. Bies, M. Cvetič and M. Liu, *Statistics of limit root bundles relevant for exact matter spectra of F-theory MSSMs*, *Physical Review D* **104**, no. 6, URL <http://dx.doi.org/10.1103/PhysRevD.104.L061903>. 7, 8, 9
- [141] Lucia Caporaso and Cinzia Casagrande and Maurizio Cornalba, *Moduli of Roots of Line Bundles on Curves*, *Transactions of the American Mathematical Society* **359**, no. 8 (2007) 3733–3768, URL <http://dx.doi.org/10.1090/S0002-9947-07-04087-1>. 7, 8
- [142] T. J. Jarvis, *Geometry of the moduli of higher spin curves*, *International Journal of Mathematics* **11** (1998) 637–663, URL <http://dx.doi.org/10.1142/S0129167X00000325>. 7
- [143] Jarvis, Tyler J., *The Picard group of the moduli of higher spin curves*, *The New York Journal of Mathematics* **7** (2001) 23–47, URL <https://nyjm.albany.edu/j/2001/7-3.html>. 7
- [144] Sergey Natanzon and Anna Pratussevitch, *Higher Spin Klein Surfaces*, *Moscow Mathematical Journal* **16** (2016) 95–124, URL <http://www.mathjournals.org/mmj/2016-016-001/2016-016-001-004.html>. 7
- [145] The Toric Varieties project authors, *The ToricVarieties project*, ([https://github.com/homalg-project/ToricVarieties\\_project](https://github.com/homalg-project/ToricVarieties_project)), 2019–2021. 8, 9
- [146] J. Halverson and J. Tian, *Cost of seven-brane gauge symmetry in a quadrillion F-theory compactifications*, *Physical Review D* **95**, no. 2, URL <http://dx.doi.org/10.1103/PhysRevD.95.026005>. 8

- [147] Cox, D.A. and Little, J.B. and Schenck, H.K., *Toric Varieties*, Graduate studies in mathematics, American Mathematical Society, 2011, ISBN 9780821848197, URL <https://dx.doi.org/10.1090/gsm/124>. 8
- [148] Cecotti, Sergio and Cheng, Miranda C. N. and Heckman, Jonathan J. and Vafa, Cumrun, *Yukawa Couplings in F-theory and Non-Commutative Geometry*, *Surveys in differential geometry* **15** (2009) 37–98, [0910.0477]. 8
- [149] Cvetič, Mirjam and Lin, Ling and Liu, Muyang and Zhang, Hao Y. and Zoccarato, Gianluca, *Yukawa Hierarchies in Global F-theory Models*, *Journal of High Energy Physics* **2020**, no. 1, URL [http://dx.doi.org/10.1007/JHEP01\(2020\)037](http://dx.doi.org/10.1007/JHEP01(2020)037). 8
- [150] Eisenbud, D. and Harris, J., *Limit linear series: Basic theory*, *Inventiones mathematicae* **85** (1986) 337–372, URL <http://dx.doi.org/10.1007/BF01389094>. 8
- [151] Osserman, Brian, *A limit linear series moduli scheme*, *Annales de l’institut Fourier* **56**, no. 4 (2006) 1165–1205, URL <http://dx.doi.org/10.5802/AIF.2209>. 8
- [152] Gavril Farkas, *Theta characteristics and their moduli*, 2012, URL <http://dx.doi.org/10.1007/s00032-012-0178-7>. 8
- [153] M. F. Atiyah, *Riemann surfaces and spin structures*, in *Annales Scientifiques de L’Ecole Normale Supérieure*, volume 4, pp. 47–62, Société mathématique de France, 1971, URL <http://www.numdam.org/item/10.24033/asens.1205.pdf>. 9
- [154] D. Mumford, *Theta characteristics of an algebraic curve*, *Ann. Sci. Ecole Norm. Sup* **4**, no. 4 (1971) 181–192, URL [http://www.numdam.org/article/ASENS\\_1971\\_4\\_4\\_2\\_181\\_0.pdf](http://www.numdam.org/article/ASENS_1971_4_4_2_181_0.pdf). 9
- [155] The Oscar authors, *OSCAR Computer Algebra System*, (<https://github.com/oscar-system/Oscar.jl/>), 2019–2021. 9
- [156] The GAP Group, *GAP – Groups, Algorithms, and Programming, Version 4.11.1*, (<https://www.gap-system.org>), 2021. 9