

Nest relief in cryptically incubating semipalmated sandpiper is quick, but vocal

Martin Bulla bulla.mar@gmail.com, Christina Muck, Daniela Trischer, Bart Kempenaers

Figure S1 | Recording equipment at the nest. The camera is mounted on a little tripod. The battery is hidden at the vegetation. Here, the nest is protected by a cage and radio frequency identification system is visible next to the cage.

Table S1 | Incubating parent's calling rate and fly-off probability before nest relief and during regular incubation.

Model	Response variable	Effect type	Effect	Estimate	95% CI	
					Lower	Upper
a	Calling rate (10min ⁻¹) (Poisson scale)	Fixed	Intercept (before exchange)	-0.923	-1.284	-0.557
			Observation (regular incubation)	-0.834	-1.085	-0.58
b	Calling rate (10min ⁻¹) (Poisson scale)	Fixed	Intercept (before exchange & ♀)	-0.887	-1.266	-0.506
			Observation (regular incubation)	-0.831	-1.186	-0.454
			Sex (♂)	-0.074	-0.317	0.171
			Day of incubation	-0.041	-0.251	0.17
			Sex × Observation	-0.001	-0.506	0.512
			Day of incubation × Observation	0.02	-0.237	0.28
c	Fly-off (0, 1) (binomial scale)	Fixed	Intercept (before exchange)	-1.896	-2.392	-1.387
			Length of observation (ln(min))	-0.315	-0.582	-0.044
			Observation (regular incubation)	-0.352	-1.145	0.437
d	Fly-off (0, 1) (binomial scale)	Fixed	Intercept (before exchange & ♀)	-2.027	-2.763	-1.331
			Length of observation (ln(min))	-0.281	-0.556	0.004
			Observation (regular incubation)	-0.538	-1.781	0.662
			Sex (♂)	0.289	-0.665	1.259
			Day of incubation	0.146	-0.322	0.606
			Sex × Observation	0.29	-1.282	1.894
			Day of incubation × Observation	-0.039	-0.755	0.639

Shown are the posterior estimates (medians) of the effect sizes with the 95% credible intervals (CI) from a posterior distribution of 5,000 simulated values generated by the 'sim' function in R (Gelman and Su 2015). Variance components were estimated by the 'lmer' function in R (Bates *et al.* 2015).

$N = 265$ observations (148 of before exchange, 117 of regular incubation) from 31 nests. Continuous predictors were z-transformed (mean-centred and divided by SD). All four models contained nest ID as random intercept. Calling was entered as count and offset specified length of observation in minutes (divided by ten minutes and ln-transformed), which not only controlled for the length of observation, but also generated a calling rate 10min⁻¹.

Distribution of fly-offs was bimodal (26 observations with one fly-off, 3 observations with two fly-offs and 3 with four fly-offs, the remaining observation no fly-offs); thus, in the analyses we used fly-off intensity as binary variable (0 = no fly-off, 1 = at least one fly-off).

Table S2 | Incubating parent's timing of calls and fly-offs before the nest relief and during other times of incubation.

Model	Response variable	Effect type	Effect	Estimate	95% CI	
					Lower	Upper
a	Call time (to observation end)	Fixed	Intercept (before nest relief)	9.59	7.68	11.48
			Type (control)	4.03	1.23	6.84
		Random (variance)	Observation ID within Nest (intercept)	31%		
			Nest (intercept)	10%		
			Residual	59%		
b	Call time (to observation end)	Fixed	Intercept (♀ & before nest relief)	9.01	6.64	11.4
			Type (control)	4.4	0.1	8.57
			Sex (♂)	1.2	-1.85	4.22
			Type × Sex	-0.8	-6.54	5.02
		Random (variance)	Observation ID within Nest (intercept)	32%		
			Nest (intercept)	10%		
			Residual	59%		
c	Fly-off time (to observation end)	Fixed	Intercept (before nest relief)	18.22	14.78	21.56
			Type (control)	-5.87	-11.07	-0.45
		Random (variance)	Observation ID within Nest (intercept)	42%		
			Nest (intercept)	0%		
d	Fly-off time (to observation end)	Fixed	Intercept (♀ & before nest relief)	21.25	16.11	26.41
			Type (control)	-8.94	-17.11	-0.52
			Sex (♂)	-4.61	-11.35	1.8
			Type × Sex	3.69	-7.39	14.13
		Random (variance)	Observation ID within Nest (intercept)	24%		
			Nest (intercept)	24%		
			Residual	52%		

Shown are the posterior estimates (medians) of the effect sizes with the 95% credible intervals (CI) from a posterior distribution of 5,000 simulated values generated by the 'sim' function in R (Gelman and Su 2015). Variance components were estimated by the 'lmer' function in R (Bates *et al.* 2015).

N_{call} = 261 calling instances from 89 before nest relief observations of 26 nests and 84 calling instances from 36 control observation 18 nests.

$N_{\text{fly-off}}$ = 25 fly-offs from 20 before nest relief observations of 14 nests and 19 fly-offs from 12 control observations of 11 nests.

Note that included are only observations (from Table S1) containing at least one call or fly-off

Table S3 | Time components of exchange behaviour in relation to sex and day of incubation.

Model	Response variable	Effect type	Effect	Estimate	95% CI	
					Lower	Upper
a	Presence before nest relief (s)	Fixed	Intercept (♀)	9.74	-9.26	28.03
			Sex of returning parent (♂)	4.77	-18.65	28.32
			Day of incubation	3.79	-26.73	33.21
			Sex × Day of incubation	-6.5	-30.32	17.57
		Random (variance)	Nest ID (intercept)	75%		
			Day of incubation	1%		
			Residual	24%		
b	Nest relief length (log(s))	Fixed	Intercept (♀ & before present)	2.43	2.2	2.66
			Left (while around)	-0.03	-0.37	0.28
			Left (during exchange)	0.85	0.63	1.06
			Sex of returning parent (♂)	0.04	-0.1	0.18
			Day of incubation	-0.13	-0.25	-0.01
			Sex × Day of incubation	0.05	-0.09	0.2
		Random (variance)	Nest ID (intercept)	25%		
			Day of incubation	2%		
c	Exchange gap (log(s))	Fixed	Intercept (♀)	2.59	2.1	3.05
			Sex of returning parent (♂)	0.48	-0.08	1.06
			Day of incubation	-0.01	-0.49	0.47
			Sex × Day of incubation	-0.19	-0.77	0.39
		Random (variance)	Nest ID (intercept)	38%		
			Day of incubation	0%		
d	Nest relief start to incubating leave (s)	Fixed	Intercept (♀ & no 'please leave')	11.98	7.78	16.12
			Sex of returning parent (♂)	-0.56	-5.92	5.07
			'Please leave' (yes)	13.82	2.34	25.45
			Day of incubation	-4.47	-8.97	-0.25
			Sex × 'Please leave'	9.49	-8.93	27.64
			Sex × Day of incubation	1.95	-3.44	7.43
		Random (variance)	Bird ID (intercept)	10%		
			'Please leave' (0,1)	36%		
			Nest ID (intercept)	5%		
			Residual	49%		

Shown are the posterior estimates (medians) of the effect sizes with the 95% credible intervals (CI) from a posterior distribution of 5,000 simulated values generated by the 'sim' function in R (Gelman and Su 2015). Variance components were estimated by the 'lmer' function in R (Bates *et al.* 2015).

$N = 163$ exchange observations from 31 nests, except for model 'd' where $N = 123$ exchange observations (from 30 nests) where incubating parent left while or after the returning parent initiated the exchange. Day of incubation was z-transformed (mean-centred and divided by SD), except for random slope in 'Exchange gap' model where non-transformed day of incubation improved model convergence. Note that 'Presence before exchange' is zero inflated (see Results) and hence the model fitted the data poorly, but binomial model with 'Presence before exchange' as binary response variable (0 = not present, 1 = present) generated similar results.

To allow for ln-transformation, we specified instantaneous exchanges (i.e. without any exchange gap) as taking 0.001s.

Table S4 | Calling during exchange in relation to sex, previous calling bouts and day of incubation.

Model	Response variable	Effect type	Effect	Estimate	95% CI	
					Lower	Upper
a	Calling while arriving (0, 1) (binomial scale)	Fixed	Intercept (♀)	-0.582	-1.207	0.027
			Sex of returning parent (♂)	-0.699	-1.516	0.117
			Day of incubation	0.226	-0.374	0.824
			Sex × Day of incubation	0.018	-0.789	0.808
		Random (variance)	Bird ID (intercept)	21%		
			Nest ID (intercept)	79%		
b	Reply of incubating parent (0, 1) (binomial scale)	Fixed	Intercept (♀)	-0.583	-1.715	0.542
			Sex of incubating parent (♂)	0.268	-1.194	1.805
			Day of incubation	-0.574	-1.791	0.665
			Sex × Day of incubation	1.15	-0.506	2.865
		Random (variance)	Bird ID (intercept)	100%		
			Nest ID (intercept)	0%		
c	Calling intensity of returning parent (while both parents around)		Intercept (♀)	1.55	1.097	1.978
			Sex of incubating parent (♂)	0.289	-0.314	0.894
			Calling intensity incubating parent	0.594	0.219	0.947
			Day of incubation	0.229	-0.177	0.623
			Sex × Calling incubation parent	-0.649	-1.148	-0.142
			Sex × Day of incubation	-0.031	-0.54	0.503
		Random (variance)	Bird ID (intercept)	33%		
			Calling intensity incubating parent	3%		
			Nest ID (intercept)	3%		
			Residual	60%		
d	Calling intensity during exchange gap		Intercept (♀)	1.641	1.373	1.912
			Sex of returning parent (♂)	-0.407	-0.781	-0.034
			Calling intensity during return	0.491	0.188	0.797
			Day of incubation	0.066	-0.198	0.325
			Sex × Calling during return	0.05	-0.366	0.458
			Sex × Day of incubation	0.199	-0.177	0.574
		Random (variance)	Bird ID (intercept)	5%		
			Calling during return	2%		
			Nest ID (intercept)	0%		
			Residual	93%		
e	Calling intensity after exchange		Intercept (♀)	0.512	0.22	0.819
			Sex of returned parent (♂)	-0.096	-0.517	0.329
			Calling intensity during return	0.139	-0.108	0.386
			Day of incubation	0.08	-0.183	0.344
			Sex × Calling during return	-0.149	-0.494	0.197
			Sex × Day of incubation	0.007	-0.361	0.358
		Random (variance)	Bird ID (intercept)	12%		
			Calling during return	3%		
			Nest ID (intercept)	0%		
			Residual	85%		

Shown are the posterior estimates (medians) of the effect sizes with the 95% credible intervals (CI) from a posterior distribution of 5,000 simulated values generated by the 'sim' function in R (Gelman and Su 2015). Variance components were estimated by the 'lmer' and 'glmer' function in R (Bates *et al.* 2015). Predictors (Day of incubation and Calling intensities) were z-transformed (mean-centred and divided by SD). Note that effects of those continuous predictors that are not specified as random slopes (because of convergence issues) might be inflated. Also, whereas we specified bird and nest as random intercepts in (b), their variance was estimated as zero and hence is not shown.

$N_{\text{Calling while arriving}}$ = 132 exchange observations of 50 parent from 28 nests.

$N_{\text{Reply of incubating parent}}$ = 31 calling while arriving observations of 24 parent from 19 nests.

$N_{\text{Calling intensity of returning parent}}$ = 103 exchange observations of 46 parent from 27 nests.

$N_{\text{Calling intensity during exchange}}$ = 109 exchange observations of 49 parent from 28 nests.

$N_{\text{Calling intensity after exchange}}$ = 108 exchange observations of 48 parent from 27 nests.

Table S5 | Calling during exchange in relation to current bout length.

Model	Response variable	Effect type	Effect	Estimate	95% CI	
					Lower	Upper
a	Calling while arriving (0, 1) (binomial scale)	Fixed	Intercept (♀)	-0.473	-1.114	0.18
			Sex of returning parent (♂)	-1.067	-1.99	-0.149
			Current bout length	0.438	-0.25	1.157
			Sex × Current bout length	0.404	-0.702	1.507
		Random	Bird ID (intercept)	0%		
			Nest ID (intercept)	100%		
b	Reply of incubating parent (0, 1) (binomial scale)	Fixed	Intercept (♀)	-0.1	-1.41	1.219
			Sex of incubating parent (♂)	-0.167	-1.82	1.471
			Current bout length	-1.184	-3.149	0.856
			Sex × Current bout length	1.482	-0.817	3.791
c	Calling intensity of returning parent (while both parents around)		Intercept (♀)	1.64	1.127	2.15
			Sex of incubating parent (♂)	0.315	-0.362	1.002
			Current bout length	-0.115	-0.475	0.236
			Sex × Current bout length	0.449	-0.181	1.058
		Random (variance)	Bird ID (intercept)	33%		
			Current bout length	3%		
			Nest ID (intercept)	0%		
			Residual	64%		
d	Calling intensity of incubating parent (while both parents around)		Intercept (♀)	0.638	0.184	1.087
			Sex of incubating parent (♂)	0.146	-0.466	0.782
			Current bout length	-0.265	-0.609	0.084
			Sex × Current bout length	0.433	-0.14	1.003
		Random (variance)	Bird ID (intercept)	25%		
			Current bout length	1%		
			Nest ID (intercept)	0%		
			Residual	74%		
e	Calling intensity during exchange gap		Intercept (♀)	1.666	1.368	1.977
			Sex of returning parent (♂)	-0.505	-0.976	-0.051
			Current bout length	0.255	-0.132	0.652
			Sex × Current bout length	-0.299	-0.827	0.216
		Random (variance)	Bird ID (intercept)	12%		
			Current bout length	11%		
			Nest ID (intercept)	0%		
f	Calling intensity after exchange		Intercept (♀)	0.53	0.262	0.799
			Sex of returning parent (♂)	-0.077	-0.468	0.328
			Current bout length	-0.001	-0.24	0.23
			Sex × Current bout length	-0.066	-0.389	0.263
		Random (variance)	Bird ID (intercept)	30%		
			Current bout length	4%		
			Nest ID (intercept)	0%		
			Residual	66%		

Shown are the posterior estimates (medians) of the effect sizes with the 95% credible intervals (CI) from a posterior distribution of 5,000 simulated values generated by the 'sim' function in R (Gelman and Su 2015). Variance components were estimated by the 'lmer' and 'glmer' function in R (Bates *et al.* 2015). Current and next bout lengths were z-transformed (mean-centred and divided by SD). Note that effects of those continuous predictors that are not specified as random slopes (because of convergence issues) might be inflated. Also, whereas we specified bird and nest as random intercepts in (b), their variance was estimated as zero and hence is not shown.

*N*_{Calling while arriving} = 123 exchange observations of 47 parents from 27 nests.

*N*_{Reply of incubating parent} = 30 calling while arriving observations of 23 parents from 18 nests.

*N*_{Calling intensity of returning parent} = 101 exchange observations of 46 parents from 26 nests.

*N*_{Calling intensity of incubating parent} = 96 exchange observations of 44 parents from 26 nests.

*N*_{Calling intensity during exchange} = 124 exchange observations of 48 parents from 28 nests.

*N*_{Calling intensity after exchange} = 123 exchange observations of 47 parents from 27 nests.

Table S6 | Next incubation bout in relation to calling during exchange.

Model	Response variable	Effect type	Effect	Estimate	95% CI	
					Lower	Upper
a	Next bout (min)	Fixed	Intercept (♀ & no arrival call)	688.5	633.4	743.4
			Arrival call (yes)	44.6	-28.5	116.3
			Sex of returning parent (♂)	-26.8	-95.4	39.3
			Sex × Arrival call	-18.7	-116.5	83.9
		Random (variance)	Bird ID (intercept)	28%		
			Nest ID (intercept)	10%		
			Residual	62%		
b	Next bout (min)	Fixed	Intercept (♀ & no reply call)	596.5	504.3	686.6
			Reply call (yes)	250.1	101	406.2
			Sex of incubating parent (♂)	85	-44	215.9
			Sex × Reply call	-161.4	-369.5	43.6
		Random (variance)	Bird ID (intercept)	19%		
			Nest ID (intercept)	4%		
			Residual	77%		
c	Next bout (min)		Intercept (♀)	704.2	651.6	756.3
			Sex of returning parent (♂)	-20.2	-78.5	39.5
			Calling intensity returning parent	-12.8	-56.7	29.3
			Calling intensity incubating parent	27.4	-11.4	66.4
			Calling intensity during exchange gap	18.1	-18.7	55.8
			Calling intensity after nest relief	43.5	-9.8	97.5
			Sex × Calling intensity returning parent	-2.5	-61.7	55.5
			Sex × Calling intensity incubating parent	-0.5	-57.1	56.1
			Sex × Calling intensity during exchange gap	33.1	-24.9	88
			Sex × Calling intensity after nest relief	-78.6	-165.3	5.2
		Random (variance)	Bird ID (intercept)	1%		
			Calling intensity during exchange gap	18%		
			Nest ID (intercept)	31%		
			Residual	50%		

Shown are the posterior estimates (medians) of the effect sizes with the 95% credible intervals (CI) from a posterior distribution of 5,000 simulated values generated by the 'sim' function in R (Gelman and Su 2015). Variance components were estimated by the 'lmer' function in R (Bates *et al.* 2015).

Calling intensities were z-transformed (mean-centred and divided by SD). Note that effects of those continuous predictors that are not specified as random slopes (because of convergence issues) might be inflated, e.g. effect of 'Calling intensity during exchange gap' in (c) disappears when added as random slope. Also note that the estimate of Sex × Calling intensity after nest relief in the (c) is not driven by the other predictors as model containing only this single interaction gives similar results.

N_a = 119 exchange observations of 47 parents from 27 nests.

N_b = 28 calling while arriving observations of 22 parents from 17 nests.

N_c = 92 exchange observations of 42 parents from 25 nests.

Table S7 | 'Please leave display' in relation to sex, day of incubation and current bout length.

Model	Response variable	Effect type	Effect	Estimate	95% CI	
					Lower	Upper
a	'Please leave' (0,1) (binomial scale)	Fixed	Intercept (♀)	-3.8	-5.82	-1.77
			Sex of incubating parent (♂)	1.31	-1.14	3.78
			Current bout (min)	-1.03	-2.42	0.3
			Sex × Current bout	1.53	-0.3	3.41
		Random (variance)	Bird ID (intercept)	99%		
			Current bout	1%		
			Nest ID (intercept)	0%		
b	'Please leave' (0,1) (binomial scale)	Fixed	Intercept (♀)	-6.2	-10.74	-1.7
			Sex of incubating parent (♂)	0.73	-4.84	6.51
			Day of incubation	1.65	0.14	3.14
			Sex × Day of incubation	2.35	-0.77	5.42
		Random	Bird ID (intercept)	93%		
			Day of incubation	7%		

Shown are the posterior estimates (medians) of the effect sizes with the 95% credible intervals (CI) from a posterior distribution of 5,000 simulated values generated by the 'sim' function in R (Gelman and Su 2015). Variance components were estimated by the 'glmer' function in R (Bates *et al.* 2015).

N = 117 exchange observations of 49 birds from 29 nests. 'Current bout' length and 'Day of incubation' were z-transformed (mean-centred and divided by SD). Note that that models without interactions generated similar results and that in (b) model with nest as random intercept did not converge.

Table S8 | Probability of calling while leaving the nest.

Response variable	Effect type	Effect	Estimate	95% CI	
				Lower	Upper
Called when left (0,1) (binomial scale)	Fixed	Intercept (♀)	-2.22	-3.01	-1.42
		Sex of incubating parent (♂)	1.01	0.1	1.93
		Day of incubation (min)	-0.24	-1.02	0.53
		Sex × Day of incubation	0.27	-0.65	1.18
	Random (variance)	Bird ID (intercept)	0%		
		Nest ID (intercept)	100%		

Shown are the posterior estimates (medians) of the effect sizes with the 95% credible intervals (CI) from a posterior distribution of 5,000 simulated values generated by the 'sim' function in R (Gelman and Su 2015). Variance components were estimated by the 'glmer' function in R (Bates *et al.* 2015).

N = 150 exchange observations of 55 birds from 30 nests. 'Day of incubation' was z-transformed (mean-centred and divided by SD). Note that model with 'Day of incubation' as random slope did not converge and model without the interaction generated similar results.