

Figure S1 | Changes in the daily high tide levels at Ceuta, Mexico. Each point depicts one day and its color indicates spring tide cycle # (1 refers to the 1st spring tide cycle of the year). Depicted are only spring tide cycles for the breeding season (i.e. period between first and last initiated clutch in a given year). The tide height data were downloaded from mobilegeographics.com (see Methods for details).




Figure S2 | Spatial variation in Snowy Plover nest locations and flooding across years at Ceuta, Mexico. Each point depicts one nest, its color indicates whether the nest was flooded (red) or not (green). The map was created with 'ggmap' R package (1).




Figure S3 | Changes in the number of initiated Snowy Plover nests across the breeding season. Number of initiated nests over the breeding season (as indicated by spring tide cycle #) and over the spring tide cycle (**a**), or in relation to maximum daily high tide (**b**). Circles represent the mean number of initiated nests within each day of a spring tide cycle (**a**), and for each 10 m tide height interval (**b**). Circle size indicates number of years. The lines and the shaded areas represent model predictions with 95% CI based on a posterior distribution of 5,000 simulated values generated from 'complex Gaussian model' outputs (Table S2 and S3) using the 'sim' function in R (2). $N = 776$ days from ten breeding seasons encompassing 62 spring tide cycles.

Table S1 | Nest initiation according to maximum daily tide height.

					95% CI				
Model	Response variable	Effect type	Effect	Estimate	Lower	Upper			
Gaussian complex	# of initiated nests (count)	Fixed	Intercept	0.952	0.647	1.242			
			Spring tide cycle number	-0.192	-0.325	-0.061			
			Maximum daily tide height	0.033	-0.044	0.113			
			Maximum daily tide height × Spring tide cycle	-0.028	-0.107	0.052			
		Random (variance)	First or second half : Spring tide cycle : Year (intercept)	9 %					
			Spring tide cycle : Year (intercept)	7 %					
			Year (intercept)	12 %					
			Residual	72 %					
			Gaussian simple	# of initiated nests (count)	Fixed	Intercept	0.945	0.65	1.23
Spring tide cycle number	-0.193	-0.321				-0.063			
Maximum daily tide height	0.036	-0.042				0.113			
Random (variance)	First or second half : Spring tide cycle : Year (intercept)	9 %							
	Spring tide cycle : Year (intercept)	7 %							
	Year (intercept)	12 %							
	Residual	72 %							
	Poisson complex	# of initiated nests (count)			Fixed	Intercept	-0.297	-0.586	-0.009
						Spring tide cycle number	-0.237	-0.37	-0.102
Maximum daily tide height			0.029	-0.055		0.115			
Maximum daily tide height × Spring tide cycle			-0.033	-0.121		0.059			
Random (variance)			Observation (intercept)	13 %					
			First or second half : Spring tide cycle : Year (intercept)	30 %					
			Spring tide cycle : Year (intercept)	19 %					
			Year (intercept)	38 %					
			Poisson simple	# of initiated nests (count)	Fixed	Intercept	-0.3	-0.589	-0.032
Spring tide cycle number	-0.237	-0.378				-0.109			
Maximum daily tide height	0.037	-0.044				0.116			
Random (variance)	Observation (intercept)	13 %							
	First or second half : Spring tide cycle : Year (intercept)	30 %							
	Spring tide cycle : Year (intercept)	19 %							
	Year (intercept)	38 %							

The posterior estimates (medians) of the effect sizes with the 95% CIs derived from a posterior distribution of 5,000 simulated values generated by the ‘sim’ function in R. Variance components were estimated by the ‘lmer’ function in R. To account for non-independence of data points ‘Year’, ‘Spring tide cycle number’ within year and indication whether the nest was initiated in the ‘First or Second half’ of the spring tide cycle where fitted as random intercepts. Overdispersion was modelled by adding ‘Observation’ level random intercept. ‘Spring tide cycle number’ is standardized within the year so that first spring tide cycle in the given year corresponds to the cycle when the first nest was initiated. ‘Day of spring tide cycle’ was transformed to radians ($2 \times \text{number of days after the last spring tide} \times \pi / \text{length of the given spring tide cycle} [\sim 14.75])$ and fitted as sine and cosine of radians. Note that despite violating model assumptions our ‘Gaussian model’ fits the data better and unlike our ‘Binomial model’ also accounts for spatial auto-correlation in residuals. $N = 776$ days from 10 years encompassing 62 spring tide cycles. Note that “Maximum daily tide height” was missing for ten days and we imputed the missing values as the mean of maximum tide heights from previous and next day. Also, excluding these 10 days does not change the results.

References

1. Kahle D, Wickham H. Spatial visualization with ggplot2. The R Journal. 2013;5(1):144-61.
2. Gelman A, Su Y. arm: Data analysis using regression and multilevel/hierarchical models [Online]. R package version 1.8-6. 2015.
3. Bates D, Mächler M, Bolker B, Walker S. Fitting Linear Mixed-Effects Models using lme4. 2015. 2015;67(1):48.