

Model-independent determination of the CKM angle γ in $B^\pm \rightarrow (K^+ K^- \pi^+ \pi^-)_D h^\pm$ decays

Martin Tat Guy Wilkinson Sneha Malde

University of Oxford

102nd LHCb week

9th December 2021




Introduction to the CKM angle γ

γ and the unitary triangle

- CP violation in SM is described by the Unitary Triangle



$$\gamma = \arg\left(-\frac{V_{ud}V_{ub}^*}{V_{cd}V_{cb}^*}\right)$$

- Only CKM angle accessible at tree level \Rightarrow
 - Negligible theoretical uncertainties
 - Ideal Standard Model benchmark
 - Compare with indirect measurements



CKMfitter Group (J. Charles et al.), Eur. Phys. J. C41, 1-131 (2005)

Sensitivity through interference



- Superposition of D^0 and \bar{D}^0
- $b \rightarrow u\bar{c}s$ and $b \rightarrow c\bar{u}s$ interference \rightarrow Sensitivity to γ

$$\begin{aligned}\mathcal{A}(B^-) &= \mathcal{A}(D^0) + r_B e^{i(\delta_B - \gamma)} \mathcal{A}(\bar{D}^0) \\ \mathcal{A}(B^+) &= \mathcal{A}(\bar{D}^0) + r_B e^{i(\delta_B + \gamma)} \mathcal{A}(D^0)\end{aligned}$$

Measurement of γ from $B^\pm \rightarrow DK^\pm$, $D \rightarrow K^+K^-\pi^+\pi^-$



- First proposed by J. Rademacker and G. Wilkinson
 - [Physics Letters B **647** \(2007\) 400](#)
 - Amplitude model by FOCUS
 - Expected γ precision from amplitude fit with 1000 candidates: 14°
- CLEO amplitude analysis
 - [Phys. Rev. D **85** \(2012\) 122002](#)
 - Expected γ precision from amplitude fit with 2000 candidates: 11°
- State of the art amplitude analysis by LHCb:
 - [JHEP **02** \(2019\) 126](#)
 - Use to develop efficient binning scheme

The $D \rightarrow K^+ K^- \pi^+ \pi^-$ decay

Binned γ analysis of the
 $D \rightarrow K^+ K^- \pi^+ \pi^-$ mode

Binned measurement of γ

- Final measurement will be model-independent
 - Non-optimal binning reduces statistical sensitivity
 - But no bias is induced in final result
- Need strong phases of D decay \rightarrow Will be measured at BESIII
- Analogous approach to $B^\pm \rightarrow Dh^\pm$, $D \rightarrow K_S^0 h^+ h^-$
 - [JHEP 02 \(2021\) 0169](#)
 - Single most precise measurement: $\gamma = (68.7^{+5.2}_{-5.1})^\circ$



The BPGGSZ method

- $B^\pm \rightarrow Dh^\pm$ amplitude:

$$\begin{aligned}\mathcal{A}(B^-) &= \mathcal{A}(D^0) + r_B e^{i(\delta_B - \gamma)} \mathcal{A}(\bar{D}^0) \\ \mathcal{A}(B^+) &= \mathcal{A}(\bar{D}^0) + r_B e^{i(\delta_B + \gamma)} \mathcal{A}(D^0)\end{aligned}$$

- $\mathcal{A}(D^0)$ and $\mathcal{A}(\bar{D}^0)$ depend on D phase space
- Strong-phase difference of D^0 and \bar{D}^0 decays inaccessible at LHCb
- Model-independent measurement: Integrate over bins of phase space

Event yield in bin i

$$\begin{aligned}N_i^- &= h_{B^-} \left(F_i + (x_-^2 + y_-^2) \bar{F}_i + 2\sqrt{F_i \bar{F}_i} (x_- c_i + y_- s_i) \right) \\ N_{-i}^+ &= h_{B^+} \left(F_i + (x_+^2 + y_+^2) \bar{F}_i + 2\sqrt{F_i \bar{F}_i} (x_+ c_i + y_+ s_i) \right)\end{aligned}$$

The BPGGSZ method

Event yield in bin i

$$N_i^- = h_{B^-} (F_i + (x_-^2 + y_-^2) \bar{F}_i + 2\sqrt{F_i \bar{F}_i} (x_- c_i + y_- s_i))$$

$$N_i^+ = h_{B^+} (F_i + (x_+^2 + y_+^2) \bar{F}_i + 2\sqrt{F_i \bar{F}_i} (x_+ c_i + y_+ s_i))$$

- CP observables:

- $x_{\pm}^{DK} = r_B^{DK} \cos(\delta_B^{DK} \pm \gamma), \quad y_{\pm}^{DK} = r_B^{DK} \sin(\delta_B^{DK} \pm \gamma)$
- $x_{\xi}^{D\pi} = \text{Re}(\xi^{D\pi}), \quad y_{\xi}^{D\pi} = \text{Im}(\xi^{D\pi}) \quad \left(\xi^{D\pi} = \frac{r_B^{D\pi}}{r_B^{DK}} e^{i(\delta_B^{D\pi} - \delta_B^{DK})} \right)$

- Fractional bin yield:

- $F_i = \frac{\int_i d\Phi |\mathcal{A}(D^0)|^2}{\sum_j \int_j d\Phi |\mathcal{A}(D^0)|^2}$
- Floated in the fit, mostly constrained by $B^{\pm} \rightarrow D\pi^{\pm}$

- Amplitude averaged strong phases can be obtained from BESIII:

$$c_i = \frac{\int_i d\Phi |\mathcal{A}(D^0)| |\mathcal{A}(\bar{D}^0)| \cos(\delta_D)}{\sqrt{\int_i d\Phi |\mathcal{A}(D^0)|^2 \int_i d\Phi |\mathcal{A}(\bar{D}^0)|^2}}, \quad s_i = \frac{\int_i d\Phi |\mathcal{A}(D^0)| |\mathcal{A}(\bar{D}^0)| \sin(\delta_D)}{\sqrt{\int_i d\Phi |\mathcal{A}(D^0)|^2 \int_i d\Phi |\mathcal{A}(\bar{D}^0)|^2}}$$

Binning scheme

Binning scheme requirements

A binning scheme must satisfy the following:

- Minimal dilution of strong phases when integrating over bins
- Enhance interference between $B^\pm \rightarrow D^0 h^\pm$ and $B^\pm \rightarrow \bar{D}^0 h^\pm$

How to bin a 5-dimensional phase space?


- Generate C++ code for LHCb amplitude model using AmpGen¹
- For each B^\pm candidate, calculate

$$\frac{\mathcal{A}(D^0)}{\mathcal{A}(\bar{D}^0)} = r_D e^{i\delta_D}$$

- Bin along δ_D and r_D , maximize Q -value to optimize


¹AmpGen by Tim Evans

Binning scheme




γ precision benchmark

- Generate 2000 $B^\pm \rightarrow DK^\pm$ candidates using LHCb model in AmpGen
 - Input values: $\gamma = 75^\circ$, $\delta_B = 130^\circ$, $r_B = 0.1$
- Perform unbinned fit to the same amplitude model using AmpGen



(a) Pull of γ




(b) Fitted γ values


Precision of γ in unbinned fit: 11°

Study of γ precision

- Binned fit setup: Optimized 2×8 bins
- Fit same AmpGen samples, using c_i , s_i and F_i from LHCb model



(a) Pull of γ



(b) Fitted γ values

Precision of γ in binned fit: 12°
Consistent with unbinned fit and Q -value

Global mass fit

Signal parameterisation

- PDF shape parameterization identical to LHCb-ANA-2020-001
- Signal: Gaussian + Modified Cruijff
- Shape fixed from MC, yield and width floated
- Combinatorial background (exponential)
- Partially reconstructed background (dini shapes)

$$f_{\text{MG}}(m|m_B, \sigma, \alpha_L, \alpha_R, \beta) \propto \begin{cases} \exp\left(\frac{-\Delta m^2(1+\beta\Delta m^2)}{2\sigma^2+\alpha_L\Delta m^2}\right), & \Delta m = m - m_B < 0 \\ \exp\left(\frac{-\Delta m^2(1+\beta\Delta m^2)}{2\sigma^2+\alpha_R\Delta m^2}\right), & \Delta m = m - m_B > 0 \end{cases}$$

Global mass fit




Figure 4: Fit of $B^\pm \rightarrow DK^\pm$ channel (left) and $B^\pm \rightarrow D\pi^\pm$ channel (right) using full Run 1 and 2 data

- $B^\pm \rightarrow DK^\pm$ yield: 3306 ± 75
- $B^\pm \rightarrow D\pi^\pm$ yield: $46\,695 \pm 256$


Binned CP fit

Binned CP fit

- Fit for CP observables
- c_i and s_i not available from BESIII yet \implies
 - Calculate using MC integration of LHCb amplitude model for now
- Use 2×8 bins
- PDF shape parameters fixed from global mass fit
- Yield of signal, low mass partially reconstructed background and combinatorial background floated
- Fractional yields F_i floated

$$\mathcal{R}_i = \begin{cases} F_i, & i = -8 \\ F_i / \sum_{j \geq i} F_j, & -8 < i \leq +8 \end{cases}$$

CP observables result: x_-^{DK} and y_-^{DK}







(a) $x_- = r_B \cos(\delta_B - \gamma)$

(b) $y_- = r_B \sin(\delta_B - \gamma)$

Pulls and uncertainties from toy studies, uncertainty from data fit in blue

CP observables result: x_+^{DK} and y_+^{DK}







(a) $x_+ = r_B \cos(\delta_B + \gamma)$

(b) $y_+ = r_B \sin(\delta_B + \gamma)$

Pulls and uncertainties from toy studies, uncertainty from data fit in blue

CP observables result: $x_\xi^{D\pi}$ and $y_\xi^{D\pi}$



(a) $x_\xi = \text{Re}(\xi)$

(b) $y_\xi = \text{Im}(\xi)$

Pulls and uncertainties from toy studies, uncertainty from data fit in blue

Systematic uncertainties

c_i and s_i systematic uncertainty

- c_i and s_i uncertainty will come from BESIII analysis
 - Mostly statistical in origin
- Largest systematic uncertainty
- Use estimated uncertainties corresponding to 20 fb^{-1} at $\psi(3770)$
- Smear c_i and s_i and do many fits to data

Summary of all systematic uncertainties

Uncertainties of CP observables in units of 10^{-2}

Source	x_-^{DK}	y_-^{DK}	x_+^{DK}	y_+^{DK}	$x_\xi^{D\pi}$	$y_\xi^{D\pi}$
Statistical	2.91	3.41	2.51	3.04	4.04	4.89
c_i, s_i	0.66	1.55	0.32	1.31	1.73	1.03
$B^\pm \rightarrow D\mu\nu$ background	0.04	0.03	0.02	0.15	0.30	0.10
$D \rightarrow K(X)l\nu_l$ background	0.15	0.05	0.11	0.03	0.35	0.25
$D \rightarrow K\pi\pi\pi$ background	0.17	0.03	0.04	0.01	0.46	0.18
Λ_b background	0.09	0.11	0.00	0.18	0.16	0.21
Bin dependent mass shape	0.21	0.05	0.17	0.01	0.37	0.11
Fit bias	0.19	0.03	0.16	0.04	0.30	0.16
Fixed yield fractions	0.02	0.03	0.02	0.02	0.01	0.01
Low mass physics effects	0.05	0.09	0.05	0.18	0.41	0.48
Mass shape	0.03	0.03	0.02	0.02	0.04	0.01
PID Efficiency	0.03	0.03	0.02	0.02	0.04	0.01
Total LHCb systematic	0.39	0.17	0.27	0.30	0.92	0.65
Total systematic	0.77	1.55	0.41	1.34	2.01	1.23

Summary of $KK\pi\pi$ analysis

Summary of CP observables

- Measured CP observables:

$$x_-^{DK} = (x.x \pm 2.9 \pm 0.4 \pm 0.7) \times 10^{-2},$$

$$y_-^{DK} = (x.x \pm 3.4 \pm 0.2 \pm 1.6) \times 10^{-2},$$

$$x_+^{DK} = (x.x \pm 2.5 \pm 0.3 \pm 0.3) \times 10^{-2},$$

$$y_+^{DK} = (x.x \pm 3.0 \pm 0.3 \pm 1.3) \times 10^{-2},$$

$$x_\xi^{D\pi} = (x.x \pm 4.0 \pm 0.9 \pm 1.7) \times 10^{-2},$$

$$y_\xi^{D\pi} = (x.x \pm 4.9 \pm 0.7 \pm 1.0) \times 10^{-2},$$

- Note: Currently using c_i and s_i from the LHCb model
- Publication strategy: Publish current results together with binned yields \rightarrow Redo fit to obtain model-independent CP observables once c_i and s_i from BESIII are available
- Will update c_i and s_i with model-dependent uncertainties

Bonus measurement

- The mode $B^\pm \rightarrow Dh^\pm$, $D \rightarrow \pi^+\pi^-\pi^+\pi^-$ very similar
- Run this through same selection (including BDT)
- Quasi-GLW observables provide additional constraints on γ
- Measure observables for both $\pi^+\pi^-\pi^+\pi^-$ and $K^+K^-\pi^+\pi^-$

$$A_h = \frac{\Gamma(B^- \rightarrow Dh^-) - \Gamma(B^+ \rightarrow Dh^+)}{\Gamma(B^- \rightarrow Dh^-) + \Gamma(B^+ \rightarrow Dh^+)},$$

$$R_{\text{CP}} = \frac{R(4h)}{R(K3\pi)},$$

$$R = \frac{\Gamma(B \rightarrow DK)}{\Gamma(B \rightarrow D\pi)}.$$

- $B^\pm \rightarrow Dh^\pm$, $D \rightarrow K\pi\pi\pi$ yields provided by Tim Evans

Global mass fit of $B^\pm \rightarrow Dh^\pm$, $D \rightarrow \pi^+\pi^-\pi^+\pi^-$




Figure 8: Fit of $B^\pm \rightarrow DK^\pm$ channel (left) and $B^\pm \rightarrow D\pi^\pm$ channel (right) using full Run 1 and 2 data

- Total yields are consistent with $KK\pi\pi$

Conclusion

- Binned γ analysis of $B^\pm \rightarrow (K^+ K^- \pi^+ \pi^-)_D h^\pm$ is mostly complete, with promising results
 - Expect total uncertainty around 12° - 14°
- Add quasi-GLW observables of $K^+ K^- \pi^+ \pi^-$ and $\pi^+ \pi^- \pi^+ \pi^-$ to further constrain γ
- Will publish model-dependent measurement for now, update with model-independent results when c_i and s_i from BESIII are available
- Analysis note: [LHCb-ANA-2021-051](#)


Thank you!

Thank you!

Backup


Binning scheme

2x4 binning scheme for $K^+K^-\pi^+\pi^-$ phase space





(a) $Q = 0.85$

2x6 binning scheme for $K^+K^-\pi^+\pi^-$ phase space




(b) $Q = 0.89$

c_i , s_i and F_i



Comparison of binned fit precision with unbinned fit

γ precision vs number of bins



Trigger requirements

Run 1 trigger requirements	(Bu_LOGlobal_TIS or Bu_LOHadronDecision_TOS) and (Bu_Hlt1TrackAllL0Decision_TOS) and (Bu_Hlt2Topo2BodyBBDTDecision_TOS or Bu_Hlt2Topo3BodyBBDTDecision_TOS or Bu_Hlt2Topo4BodyBBDTDecision_TOS or Bu_Hlt2IncPhiDecision_TOS)
Run 2 trigger requirements	(Bu_LOGlobal_TIS or Bu_LOHadronDecision_TOS) and (Bu_Hlt1TrackMVADecision_TOS or Bu_Hlt1TwoTrackMVADecision_TOS) and (Bu_Hlt2Topo2BodyDecision_TOS or Bu_Hlt2Topo3BodyDecision_TOS or Bu_Hlt2Topo4BodyDecision_TOS or Bu_Hlt2IncPhiDecision_TOS)


Rectangular cuts before BDT

Number	Variable description	Cut
1	DTF converged	True
2	Bachelor momentum	$< 100\text{GeV}$
3	Bachelor has RICH	True
4	D invariant mass	$[1839.84, 1889.84]\text{MeV}$
5	B^\pm invariant mass	$[5080, 5800]\text{MeV}$
6	K^\pm daughter PID	> -10
7	π^\pm daughter PID	< 20

Boosted Decision Tree

- BDTG from TMVA Toolkit
- Signal sample: $B^\pm \rightarrow DK^\pm$ and $B^\pm \rightarrow D\pi^\pm$ MC samples
- Background sample: Data sample with $m_{B^\pm}^{\text{DTF}} \in [5800, 7000]\text{MeV}$
- Random, equal sized test and training samples

BDT training results



(a) BDT output

(b) BDT output on a logarithmic scale

Rectangular cuts after BDT

Number	Variable description	Cut
8	K^\pm bachelor PID	> 4
9	π^\pm bachelor PID	< 4
10	Bachelor is muon	False
11	z flight significance	> 2
12	K^\pm PID	> 0
13	K_S^0 mass veto	$[477, 507]\text{MeV}$


BDT training variables

Name	Rank (%)	Description
$\log(D0_RHO_BPV)$	7.7	D radial distance to beamline
$\log(Bu_FDCHI2_OWNPV)$	6.3	B^\pm flight distance χ^2
$\log(Bu_RHO_BPV)$	6.1	B^\pm radial distance to beamline
$\log(Bach_PT)$	6.1	Bachelor transverse momentum
$Bu_PTASY_1.5$	5.3	B^\pm asymmetry parameter
$\log(1-D0_DIRA_BPV)$	5.0	Angle between PV and D
$\log(Bu_IPCHI2_OWNPV)$	4.8	B^\pm impact parameter χ^2
$\log(1-Bu_DIRA_BPV)$	4.7	Angle between PV and B^\pm
$\log(h[1,2]_PT)$	4.4	K^\pm transverse momentum
$Bu_MAXDOCA$	4.4	B^\pm distance of closest approach
$\log(Bach_IPCHI2_OWNPV)$	4.1	Bachelor impact parameter χ^2


BDT training particles

Name	Rank (%)	Description
$\log(\text{Bu_constD0PV_D0_P})$	3.7	D momentum from DTF
$\log(\text{D0_VTXCHI2D0F})$	3.3	$D0$ vertex fit χ^2
$\log(\text{h}[3,4]_{\text{-IPCHI2_OWNPV}})$	3.3	π^\pm impact parameter χ^2
$\log(\text{D0_IPCHI2_OWNPV})$	3.2	D impact parameter χ^2
$\log(\text{h}[3,4]_{\text{-PT}})$	3.2	π^\pm transverse momentum
$\log(\text{Bu_PT})$	2.8	B^\pm transverse momentum
$\log(\text{h}[1,2]_{\text{-P}})$	2.8	K^\pm momentum
$\log(\text{Bach_P})$	2.7	Bachelor momentum
$\log(\text{Bu_constD0PV_P})$	2.6	B^\pm momentum from DTF
$\log(\text{h}[1,2]_{\text{-IPCHI2_OWNPV}})$	2.5	K^\pm impact parameter χ^2
D0_MAXDOCA	2.5	D distance of closest approach
$\log(\text{Bu_VTXCHI2D0F})$	2.0	B^\pm vertex fit χ^2
$\log(\text{h}[3,4]_{\text{-P}})$	1.9	π^\pm momentum

BDT optimization study



(a) Run 1



(b) Run 2

- Run 1: Pick BDT working point at 0.65
- Run 2: Pick BDT working point at 0.75

Partially reconstructed background

- $B^\pm \rightarrow D\pi^\pm$:

- ① $B^\pm \rightarrow (D^{*0} \rightarrow D^0[\pi^0])\pi^\pm$

- ② $B^0 \rightarrow (D^{*\mp} \rightarrow D^0[\pi^\mp])\pi^\pm$

- ③ $B^{\pm(0)} \rightarrow D^0[\pi^{0(\mp)}]\pi^\pm$

- ④ $B^\pm \rightarrow (D^{*0} \rightarrow D^0[\gamma])\pi^\pm$

- $B^\pm \rightarrow DK^\pm$:

- ① $B^\pm \rightarrow (D^{*0} \rightarrow D^0[\pi^0])K^\pm$

- ② $B^0 \rightarrow (D^{*\mp} \rightarrow D^0[\pi^\mp])K^\pm$

- ③ $B^{\pm(0)} \rightarrow D^0[\pi^{0(\mp)}]K^\pm$


- ④ $B^\pm \rightarrow (D^{*0} \rightarrow D^0[\gamma])K^\pm$

- ⑤ $B_s^0 \rightarrow \bar{D}^0[\pi^+]K^-$


- ⑥ Mis-ID from partially reconstructed $B^\pm \rightarrow D\pi^\pm$ channel

Charmless background

- $B \rightarrow KK\pi\pi K$ background in $B \rightarrow DK$ channel
- Flight significance cut at 2
- Fix remaining background with Gaussian shape of lower sideband



(a) No FS cut




(b) FS cut at 2

Figure 13: B invariant mass in lower D sideband

$D \rightarrow K\pi\pi\pi\pi^0$ mis-ID background


- $B^\pm \rightarrow Dh^\pm$, $D \rightarrow K\pi\pi\pi[\pi^0]$
- π^0 not reconstructed \rightarrow Lower D mass
- Single mis-ID: $K\pi\pi\pi \rightarrow KK\pi\pi \rightarrow$ Higher D mass
- Generate RapidSim samples, reweight with PIDCalib2




Conclusion: Fix shape from RapidSim, allow yield to float

$D \rightarrow K\pi\pi\pi$ mis-ID background

- $B^\pm \rightarrow Dh^\pm$, $D \rightarrow K\pi\pi\pi$
- Single mis-ID: $K\pi\pi\pi \rightarrow KK\pi\pi$
- Triple mis-ID: $\pi\pi K\pi \rightarrow KK\pi\pi$
- Use LHCb MC generated with AmpGen, reweight with PIDCalib2




(a) Single mis-ID




(b) Triple mis-ID

Figure 14: D invariant mass

$D \rightarrow K\pi\pi\pi$ mis-ID background



(a) Single mis-ID



(b) Triple mis-ID

Figure 15: B invariant mass


Conclusion: Negligible impact, include in systematics

D semileptonic backgrounds

- $B^\pm \rightarrow Dh^\pm$, $D \rightarrow K(X)l\nu$, $K(X) \rightarrow K\pi\pi$
 - $K_1(1270)$
 - $K_1(1400)$
 - $K^*(1410)$
 - $K^*(1680)$
 - $K_2^*(1430)$
- Single mis-ID: $K\mu\pi\pi \rightarrow KK\pi\pi$
- Double mis-ID: $K\pi\pi\mu \rightarrow KK\pi\pi$
- Generate RapidSim samples, reweight with PIDCalib2


D semileptonic backgrounds

B mass of SL $D \rightarrow K(X)\mu\nu$ backgrounds, $B \rightarrow DK$



(a) $B \rightarrow DK$

B mass of SL $D \rightarrow K(X)\mu\nu$ backgrounds, $B \rightarrow D\pi$



(b) $B \rightarrow D\pi$

Conclusion: Negligible impact, include in systematics

Remaining systematic uncertainties


Different strategies for evaluating systematic uncertainties:

- Generate toy datasets with systematics, fit with default model and take the bias as a systematic:
 - Small backgrounds ($D \rightarrow K(X)l\nu_l$, $D \rightarrow K\pi\pi\pi$, $B \rightarrow Dl\nu_l$, Λ_b)
 - Bin dependent mass shape
 - Low mass physics effects
- Do multiple fits to data while smearing parameters:
 - c_i and s_i
 - Mass shape
 - Fixed yield fractions
 - PID efficiency
- Fit bias: Take bias toys as systematic uncertainty

Efficiency related systematics:


- Difference in $B^\pm \rightarrow DK^\pm$ and $B^\pm \rightarrow D\pi^\pm$ phase space acceptance
- Efficiency correction of c_i and s_i

Efficiency differences between $B^\pm \rightarrow DK^\pm$ and $B^\pm \rightarrow D\pi^\pm$




Conclusion: More or less identical phase space acceptance, no systematic uncertainty considered

Efficiency correction of c_i and s_i



Need to reweight events to account for efficiency differences between AmpGen samples and LHCb MC

Efficiency correction of c_i and s_i



After reweighting, use weights to recalculate c_i and s_i

Conclusion: Efficiency correction of c_i and s_i is an order of magnitude smaller than their uncertainties, no systematic uncertainty considered