

Model-independent measurement of γ using $B^\pm \rightarrow [h^+ h^- \pi^+ \pi^-]_D h^\pm$ decays

Sneha Malde ¹, Jonas Rademacker ², **Martin Tat**^{1,4}, Ben Westhenry ²,
Mark Whitehead ³, Guy Wilkinson ¹

¹University of Oxford, ²University of Bristol, ³University of Glasgow, ⁴Heidelberg University

5th September 2024



UNIVERSITÄT
HEIDELBERG
ZUKUNFT
SEIT 1386




Outline


- 1 Introduction to γ and CP violation
- 2 Strong-phase inputs from BESIII
- 3 Analysis: Global fit
- 4 Analysis: CP fit
- 5 Analysis: Interpretation
- 6 Conclusion and future prospects

Introduction to γ and CP violation

- CPV in SM is described by the Unitary Triangle, with angles α , β , γ
- The angle $\gamma = \arg\left(-\frac{V_{ud}V_{ub}^*}{V_{cd}V_{cb}^*}\right)$ is very important:
 - 1 Negligible theoretical uncertainties: Ideal SM benchmark
 - 2 Accessible at tree level: Indirectly probe New Physics that enter loops
 - 3 Compare with a global CKM fit: Is the Unitary Triangle a triangle?



(a) Tree level: $\gamma = (72.1^{+5.4}_{-5.7})^\circ$




(b) Loop level: $\gamma = (65.5^{+1.1}_{-2.7})^\circ$

CKMfitter Group (J. Charles et al.), Eur. Phys. J. C41, 1-131 (2005), updated results and plots available at:
<http://ckmfitter.in2p3.fr>

Sensitivity through interference

Measure γ through interference effects in $B^\pm \rightarrow DK^\pm$



- Superposition of D^0 and \bar{D}^0
 - Consider D^0/\bar{D}^0 decays to the same final state, such as $D \rightarrow K^+ K^-$
- $b \rightarrow u\bar{c}s$ and $b \rightarrow c\bar{u}s$ interference \rightarrow Sensitivity to γ

$$\mathcal{A}(B^-) = \mathcal{A}_B \left(\mathcal{A}_{D^0} + r_B e^{i(\delta_B - \gamma)} \mathcal{A}_{\bar{D}^0} \right)$$

$$\mathcal{A}(B^+) = \mathcal{A}_B \left(\mathcal{A}_{\bar{D}^0} + r_B e^{i(\delta_B + \gamma)} \mathcal{A}_{D^0} \right)$$

Multi-body charm decays

In this presentation, four-body charm decays are considered:

$$D^0 \rightarrow K^+ K^- \pi^+ \pi^-$$

$$D^0 \rightarrow \pi^+ \pi^- \pi^+ \pi^-$$

Note: Such decays have a five-dimensional phase space!

Degrees of freedom for an N -body decay

$4N$ (momentum components)

– N ($E_i^2 - p_i^2 = m_i^2$)

– 4 (energy-momentum conservation)


– 3 (choice of frame)

= $3N - 7$ degrees of freedom

Previous studies of γ with $B^\pm \rightarrow DK^\pm$, $D \rightarrow K^+K^-\pi^+\pi^-$

- ① First proposed by J. Rademacker and G. Wilkinson:
 - [Physics Letters B **647** \(2007\) 400](#)
 - Amplitude model by FOCUS
 - Expected γ precision from amplitude fit with 1000 candidates: 14°
- ② CLEO amplitude analysis:
 - [Phys. Rev. D **85** \(2012\) 122002](#)
 - Expected γ precision from amplitude fit with 2000 candidates: 11°
- ③ State of the art amplitude analysis by LHCb:
 - [JHEP **02** \(2019\) 126](#)
- ④ Model-dependent measurement by LHCb:
 - [Eur. Phys. J. C **83** 547 \(2023\)](#)
 - Optimised binning scheme using LHCb amplitude model

Inputs to $D^0 \rightarrow K^+ K^- \pi^+ \pi^-$ mode



Previous studies of γ with $B^\pm \rightarrow DK^\pm$, $D \rightarrow \pi^+\pi^-\pi^+\pi^-$

① CLEO amplitude analysis:

- [JHEP 05 \(2017\) 143](#)


② CLEO-c strong-phase measurement:

- [JHEP 01 \(2018\) 144](#)
- Expected γ statistical (systematic) precision with 2×5 bins is 9.7° (7.4°)

③ For this LHCb publication:

- New amplitude model from BESIII [Chin. Phys. C 48 \(2024\) 083001](#)
- \rightarrow Motivation to perform c_i/s_i measurements using a new binning scheme [Phys. Rev. D 110 \(2024\) 112008](#)
- Results presented from both CLEO-c and BESIII binning schemes
- Selection unified between Bristol and Oxford

Inputs to $D^0 \rightarrow \pi^+ \pi^- \pi^+ \pi^-$ mode





Selection of $B^\pm \rightarrow DK^\pm$, $D \rightarrow \pi^+\pi^-\pi^+\pi^-$

Selection of $B^\pm \rightarrow [\pi^+\pi^-\pi^+\pi^-]_D h^\pm$ is mostly similar to that in phase-space integrated measurement [Eur. Phys. J. C **83** 547 \(2023\)](#)

- Minor differences adapted for binned measurement:
- Flight significance cut loosened from 4 to 2
 - Phase-space binned analysis is less sensitive to charmless background
- Hadrons from D^0 : $\text{ProbNN}\pi \cdot (1 - \text{ProbNN}k) > 0.05$
 - Highly efficient at rejecting combinatorial background
 - Phase-space binned analysis benefits from higher purity in low-yield bins

Phase-space integrated CP observables

Phase-space integrated study of γ :
 Charged asymmetries measured for $D \rightarrow K^+ K^- \pi^+ \pi^-$ and
 $D \rightarrow \pi^+ \pi^- \pi^+ \pi^-$ in Eur. Phys. J. C **83** 547 (2023)





$$D \rightarrow K^+ K^- \pi^+ \pi^-$$

• $B^\pm \rightarrow [h^+ h^- \pi^+ \pi^-]_D h^\pm$ asymmetries:

- $D \rightarrow K^+ K^- \pi^+ \pi^-$: $\mathcal{A} = 0.095 \pm 0.023 \pm 0.002$
- $D \rightarrow \pi^+ \pi^- \pi^+ \pi^-$: $\mathcal{A} = 0.061 \pm 0.013 \pm 0.002$

Phase-space integrated CP observables

Phase-space integrated study of γ :
 Charged asymmetries measured for $D \rightarrow K^+ K^- \pi^+ \pi^-$ and
 $D \rightarrow \pi^+ \pi^- \pi^+ \pi^-$ in Eur. Phys. J. C **83** 547 (2023)



$$D \rightarrow \pi^+ \pi^- \pi^+ \pi^-$$

• $B^\pm \rightarrow [h^+ h^- \pi^+ \pi^-]_D h^\pm$ asymmetries:


- $D \rightarrow K^+ K^- \pi^+ \pi^-$: $\mathcal{A} = 0.095 \pm 0.023 \pm 0.002$
- $D \rightarrow \pi^+ \pi^- \pi^+ \pi^-$: $\mathcal{A} = 0.061 \pm 0.013 \pm 0.002$

Multi-body D decays

Main focus of this talk: Discuss phase-space binned analysis of

$$D \rightarrow h^+ h^- \pi^+ \pi^-$$

- Strong-phase difference δ_D is a function of phase space
- Compare yields of B^+ and B^- and determine the asymmetry in local phase space regions, known as phase-space bins



$$\begin{aligned} |\mathcal{A}(B^-)|^2 &\propto |\mathcal{A}_{D^0}(\Phi)|^2 + r_B^2 |\mathcal{A}_{\bar{D}^0}(\Phi)|^2 \\ &\quad + 2r_B |\mathcal{A}_{D^0}(\Phi)| |\mathcal{A}_{\bar{D}^0}(\Phi)| \cos(\delta_B - \gamma + \delta_D) \end{aligned}$$

The BPGGSZ method

Event yield in bin i

$$N_i^- = h_{B^-} (F_i + (x_-^2 + y_-^2) \bar{F}_i + 2\sqrt{F_i \bar{F}_i} (x_- c_i + y_- s_i))$$
$$N_{-i}^+ = h_{B^+} (F_i + (x_+^2 + y_+^2) \bar{F}_i + 2\sqrt{F_i \bar{F}_i} (x_+ c_i + y_+ s_i))$$

- CP observables:

- $x_{\pm}^{DK} = r_B^{DK} \cos(\delta_B^{DK} \pm \gamma), \quad y_{\pm}^{DK} = r_B^{DK} \sin(\delta_B^{DK} \pm \gamma)$
- $x_{\xi}^{D\pi} = \text{Re}(\xi^{D\pi}), \quad y_{\xi}^{D\pi} = \text{Im}(\xi^{D\pi}) \quad \left(\xi^{D\pi} = \frac{r_B^{D\pi}}{r_B^{DK}} e^{i(\delta_B^{D\pi} - \delta_B^{DK})} \right)$

- Fractional bin yield:



- $F_i = \frac{\int_i d\Phi |\mathcal{A}(D^0)|^2}{\sum_j \int_j d\Phi |\mathcal{A}(D^0)|^2}$
- Floated in the fit, mostly constrained by $B^{\pm} \rightarrow D\pi^{\pm}$

- Amplitude-averaged strong phases:

$$c_i = \frac{\int_i d\Phi |\mathcal{A}(D^0)| |\mathcal{A}(\bar{D}^0)| \cos(\delta_D)}{\sqrt{\int_i d\Phi |\mathcal{A}(D^0)|^2 \int_i d\Phi |\mathcal{A}(\bar{D}^0)|^2}} \quad s_i = \frac{\int_i d\Phi |\mathcal{A}(D^0)| |\mathcal{A}(\bar{D}^0)| \sin(\delta_D)}{\sqrt{\int_i d\Phi |\mathcal{A}(D^0)|^2 \int_i d\Phi |\mathcal{A}(\bar{D}^0)|^2}}$$

$D^0 \rightarrow K^+ K^- \pi^+ \pi^-$ binning scheme

- Interpretation of γ from the multi-body charm decays require external inputs of the charm strong-phase differences
- Measure model-independent strong-phases at a charm factory, such as BESIII, using an optimised binning scheme





$D^0 \rightarrow K^+ K^- \pi^+ \pi^-$ binning scheme

Model-dependent measurement with $D \rightarrow K^+ K^- \pi^+ \pi^-$

From the phase-space binned asymmetries, we obtain:

$$\gamma = (116_{-14}^{+12})^\circ$$



[Eur. Phys. J. C 83, 547 \(2023\)](#)

How will this evolve with model-independent BESIII inputs? Will the 3σ tension reduce?


What is new from previous analysis of $K^+K^-\pi^+\pi^-$?

- New binned strong-phase analyses of $D \rightarrow K^+K^-\pi^+\pi^-$ and $D \rightarrow \pi^+\pi^-\pi^+\pi^-$ have recently been made public by BESIII
 - $D^0 \rightarrow KK\pi\pi$: [arXiv:2502.12873](https://arxiv.org/abs/2502.12873)
 - $D^0 \rightarrow \pi\pi\pi\pi$: [Phys. Rev. D **110** \(2024\) 112008](#)
 - For $D \rightarrow \pi^+\pi^-\pi^+\pi^-$, these improve in precision on earlier binned study made with CLEO-c data [JHEP **01** \(2018\) 144](#)
- Make first binned model-independent measurement with $D \rightarrow K^+K^-\pi^+\pi^-$, updating earlier LHCb model-dependent analysis
- Use same strategy for $D \rightarrow \pi^+\pi^-\pi^+\pi^-$ with a joint Oxford-Bristol selection
- After checking for compatibility, perform joint analysis

BESIII preliminary $D^0 \rightarrow K^+ K^- \pi^+ \pi^-$ strong-phase results

First binned strong-phase analysis of $D^0 \rightarrow K^+ K^- \pi^+ \pi^-$, which uses the 2×4 binning scheme with 20 fb^{-1} $\psi(3770)$ data

$$\begin{aligned}c_1 &= -0.22 \pm 0.08 \pm 0.01 \\s_1 &= -0.47 \pm 0.22 \pm 0.04 \\c_2 &= +0.79 \pm 0.04 \pm 0.01 \\s_2 &= -0.17 \pm 0.16 \pm 0.04 \\c_3 &= +0.862 \pm 0.029 \pm 0.008 \\s_3 &= +0.26 \pm 0.14 \pm 0.02 \\c_4 &= -0.39 \pm 0.08 \pm 0.01 \\s_4 &= +0.52 \pm 0.24 \pm 0.04\end{aligned}$$



Measured values (black) are consistent and close to LHCb model predictions (blue), so central values are not expected to change much

BESIII preliminary $D^0 \rightarrow \pi^+\pi^-\pi^+\pi^-$ strong-phase results

Small differences between model prediction and measurement, but data points are generally close to the unit circle

$$c_1 = +0.12 \pm 0.09 \pm 0.02$$

$$s_1 = -0.42 \pm 0.21 \pm 0.04$$

$$c_2 = +0.74 \pm 0.04 \pm 0.02$$

$$s_2 = -0.39 \pm 0.16 \pm 0.06$$

$$s_3 = -0.25 \pm 0.12 \pm 0.03$$


$$c_3 = +0.81 \pm 0.03 \pm 0.01$$

$$c_4 = +0.42 \pm 0.06 \pm 0.02$$

$$s_4 = +0.86 \pm 0.19 \pm 0.07$$



$$c_5 = -0.27 \pm 0.09 \pm 0.03$$

$$s_5 = -0.22 \pm 0.25 \pm 0.08$$





The HyperPlot software is used (binary lookup tree in 5D phase space)

Global fit of $K^+K^-\pi^+\pi^-$ remains as in model-dependent publication:



- $B^\pm \rightarrow [K^+K^-\pi^+\pi^-]_D h^\pm$ signal yield:
 - $B^\pm \rightarrow DK^\pm$: 3280 ± 41
 - $B^\pm \rightarrow D\pi^\pm$: 47610 ± 231

Global fit of $\pi^+\pi^-\pi^+\pi^-$ has a good fit quality:





- $B^\pm \rightarrow [\pi^+\pi^-\pi^+\pi^-]_D h^\pm$ signal yield:
 - $B^\pm \rightarrow DK^\pm$: 9172 ± 110
 - $B^\pm \rightarrow D\pi^\pm$: 132246 ± 394

After global fit, perform a “CP fit” to study CP violation:





- Split candidates by:
 - ① B^+ and B^- charges
 - ② $B^\pm \rightarrow DK^\pm$ and $B^\pm \rightarrow D\pi^\pm$ decays
 - ③ D phase-space bins
- Combinatorial and low-mass backgrounds are floating in each category
- Parameterise signal yields in terms of x_\pm^{DK} , y_\pm^{DK} , $x_\xi^{D\pi}$, $y_\xi^{D\pi}$
- $2N - 1$ floating F_i parameters
- c_i and s_i are Gaussian constrained

Example of bin asymmetry in $D \rightarrow K^+ K^- \pi^+ \pi^-$ bin -3:




CP fit bin asymmetry

Example of bin asymmetry in $D \rightarrow \pi^+ \pi^- \pi^+ \pi^-$ bin +5:




Bin asymmetries

$B^\pm \rightarrow [K^+ K^- \pi^+ \pi^-]_D h^\pm$ bin asymmetries




$B^\pm \rightarrow DK^\pm$




$B^\pm \rightarrow D\pi^\pm$

Bin asymmetries

$$B^\pm \rightarrow [\pi^+ \pi^- \pi^+ \pi^-]_D h^\pm \text{ bin asymmetries}$$




$$B^\pm \rightarrow DK^\pm$$




$$B^\pm \rightarrow D\pi^\pm$$

Likelihood scan of CP observables

x_{\pm}^{DK} agree well between likelihood scan and Hesse approximation



x_{-}^{DK}




x_{+}^{DK}


$$D^0 \rightarrow K^+ K^- \pi^+ \pi^-$$

Likelihood scan of CP observables

y_{\pm}^{DK} diverges from Hesse approximation outside 1σ



y_{-}^{DK}



y_{+}^{DK}

$$D^0 \rightarrow K^+ K^- \pi^+ \pi^-$$

What do the likelihood scans tell us?

- Uncertainties from c_i and s_i are significant, which justifies Gaussian constraining c_i and s_i
- New strategy:
 - ① Produce a likelihood function from CP fit
 - ② Interpret CP observables in terms of γ , etc
 - ③ Must profile all nuisance parameters (F_i , c_i , s_i , backgrounds yields, normalisation constants)
 - ④ Provide direct measurements of γ , δ_B and r_B

Summary of LHCb internal systematic uncertainties

| Source | x_-^{DK} | y_-^{DK} | x_+^{DK} | y_+^{DK} | $x_\xi^{D\pi}$ | $y_\xi^{D\pi}$ |
|--|------------|------------|------------|------------|----------------|----------------|
| Statistical | 2.87 | 3.40 | 2.51 | 3.05 | 4.24 | 5.17 |
| Mass shape | 0.02 | 0.02 | 0.03 | 0.06 | 0.02 | 0.04 |
| Bin-dependent mass shape | 0.11 | 0.05 | 0.10 | 0.19 | 0.68 | 0.16 |
| PID efficiency | 0.02 | 0.02 | 0.03 | 0.06 | 0.02 | 0.04 |
| Low-mass background model | 0.02 | 0.02 | 0.03 | 0.04 | 0.02 | 0.02 |
| Charmless background | 0.14 | 0.15 | 0.12 | 0.14 | 0.01 | 0.02 |
| CP violation in low-mass background | 0.01 | 0.10 | 0.08 | 0.12 | 0.07 | 0.26 |
| Semi-leptonic b -hadron decays | 0.05 | 0.27 | 0.06 | 0.01 | 0.07 | 0.19 |
| Semi-leptonic charm decays | 0.02 | 0.07 | 0.03 | 0.15 | 0.06 | 0.24 |
| $D \rightarrow K^- \pi^+ \pi^- \pi^+$ background | 0.11 | 0.05 | 0.07 | 0.04 | 0.09 | 0.05 |
| $\Lambda_b \rightarrow p D \pi^-$ background | 0.01 | 0.25 | 0.14 | 0.04 | 0.06 | 0.34 |
| $D \rightarrow K^- \pi^+ \pi^- \pi^+ \pi^0$ background | 0.30 | 0.05 | 0.19 | 0.07 | 0.05 | 0.01 |
| Fit bias | 0.06 | 0.05 | 0.13 | 0.02 | 0.06 | 0.13 |
| Total LHCb systematic | 0.37 | 0.43 | 0.34 | 0.32 | 0.70 | 0.57 |

Give systematic uncertainties in terms of CP observables (not γ) since these are more Gaussian and better behaved

From CP fit, we have a (negative log) likelihood function with nuisance parameters n_k :

$$\mathcal{L}(x_-^{DK}, y_-^{DK}, x_+^{DK}, y_+^{DK}, x_\xi^{D\pi}, y_\xi^{D\pi}, \{n_k\})$$

Express in terms of physics parameters:

$$\mathcal{L}(\gamma, \delta_B^{DK}, r_B^{DK}, \delta_B^{D\pi}, r_B^{D\pi}, \{n_k\})$$

In this step, also add a Gaussian smearing term on CP observables to account for internal LHCb systematics (see backup)

Interpretation results

Results from interpretation of $K^+K^-\pi^+\pi^-$, after correcting for biases in central values (not uncertainties):

Model independent

$$\gamma = (121 \pm 16)^\circ$$

$$\delta_B^{DK} = (74 \pm 14)^\circ$$

$$r_B^{DK} = (12.1 \pm 3.0) \times 10^{-2}$$

$$\delta_B^{D\pi} = (243 \pm 116)^\circ$$

$$r_B^{D\pi} = (1 \pm 6) \times 10^{-3}$$

Model dependent

$$\gamma = (116_{-14}^{+12})^\circ$$

$$\delta_B^{DK} = (81_{-13}^{+14})^\circ$$

$$r_B^{DK} = (11.0 \pm 2.0) \times 10^{-2}$$

$$\delta_B^{D\pi} = (298_{-118}^{+62})^\circ$$

$$r_B^{D\pi} = (4_{-4}^{+5}) \times 10^{-3}$$

Central value of γ remains high...

... it seems that the large tension with the LHCb global result

$$\gamma = (64.6 \pm 2.8)^\circ \text{ remains}$$

Interpretation results

Results from interpretation of $h^+h^-\pi^+\pi^-$, after correcting for biases in central values (not uncertainties):

$$K^+K^-\pi^+\pi^-$$

$$\gamma = (121 \pm 16)^\circ$$

$$\delta_B^{DK} = (74 \pm 14)^\circ$$

$$r_B^{DK} = (12.1 \pm 3.0) \times 10^{-2}$$

$$\delta_B^{D\pi} = (243 \pm 116)^\circ$$

$$r_B^{D\pi} = (1 \pm 6) \times 10^{-3}$$

$$\pi^+\pi^-\pi^+\pi^-$$

$$\gamma = (45 \pm 10)^\circ$$

$$\delta_B^{DK} = (115 \pm 9)^\circ$$

$$r_B^{DK} = (9.4 \pm 1.9) \times 10^{-2}$$

$$\delta_B^{D\pi} = (194 \pm 74)^\circ$$

$$r_B^{D\pi} = (0 \pm 4) \times 10^{-3}$$

$\pi^+\pi^-\pi^+\pi^-$ is in much better agreement with LHCb global result, but there is a tension with $K^+K^-\pi^+\pi^-$...

Interpretation results

Results from interpretation of $h^+h^-\pi^+\pi^-$, after correcting for biases in central values (not uncertainties):

$$K^+K^-\pi^+\pi^-$$

$$\pi^+\pi^-\pi^+\pi^-$$

$$\gamma = (121 \pm 16)^\circ$$

$$\gamma = (45 \pm 10)^\circ$$

$$\delta_B^{DK} = (74 \pm 14)^\circ$$

$$\delta_B^{DK} = (115 \pm 9)^\circ$$

$$r_B^{DK} = (12.1 \pm 3.0) \times 10^{-2}$$

$$r_B^{DK} = (9.4 \pm 1.9) \times 10^{-2}$$

$$\delta_B^{D\pi} = (243 \pm 116)^\circ$$

$$\delta_B^{D\pi} = (194 \pm 74)^\circ$$

$$r_B^{D\pi} = (1 \pm 6) \times 10^{-3}$$

$$r_B^{D\pi} = (0 \pm 4) \times 10^{-3}$$

$\pi^+\pi^-\pi^+\pi^-$ is in much better agreement with LHCb global result, but there is a tension with $K^+K^-\pi^+\pi^-$...
...but how Gaussian are these uncertainties?

Interpretation results

We can also compare the statistical sensitivity of $\pi^+\pi^-\pi^+\pi^-$ between the CLEO-c and BESIII binning schemes (keep c_i and s_i fixed)

BESIII

$$\gamma = (47 \pm 10)^\circ$$

$$\delta_B^{DK} = (113 \pm 9)^\circ$$

$$r_B^{DK} = (9.2 \pm 1.6) \times 10^{-2}$$

$$\delta_B^{D\pi} = (208 \pm 58)^\circ$$

$$r_B^{D\pi} = (3.9 \pm 2.7) \times 10^{-3}$$

CLEO-c

$$\gamma = (51 \pm 20)^\circ$$

$$\delta_B^{DK} = (109 \pm 19)^\circ$$

$$r_B^{DK} = (6.5 \pm 1.8) \times 10^{-2}$$

$$\delta_B^{D\pi} = (310 \pm 508)^\circ$$


$$r_B^{D\pi} = (0 \pm 5) \times 10^{-3}$$

Very good agreement!


BESIII binning scheme, which has more bins and values of c_i and s_i further from the origin, performs better

Likelihood scan of interpretation fit

In fact, a likelihood scan shows that $D \rightarrow K^+ K^- \pi^+ \pi^-$ and $D \rightarrow \pi^+ \pi^- \pi^+ \pi^-$ 2σ contours overlap



γ vs δ_B^{DK}




γ vs r_B^{DK}


When all biases, correlations and non-Gaussian uncertainties are accounted for, the tension with the LHCb average has reduced significantly

Likelihood scan of interpretation fit

In fact, a likelihood scan shows that $D \rightarrow K^+ K^- \pi^+ \pi^-$ and $D \rightarrow \pi^+ \pi^- \pi^+ \pi^-$ 2σ contours overlap



γ vs δ_B^{DK}



γ vs r_B^{DK}

However, with all the non-Gaussian behaviour, are we sure these contours cover 68% and 95% ?

Feldman-Cousins method, or Plugin, is a “brute-force” approach to assigning a confidence interval

At each scan point of γ , perform these fits to data:

- 1 Fit with all parameters floating, and save the log-likelihood χ^2
- 2 Fit with γ fixed to scan point, and save χ_{fix}^2
- 3 Calculate $\Delta\chi_{\text{data}}^2 = \chi_{\text{fix}}^2 - \chi^2$

We expect $\Delta\chi_{\text{data}}^2$ to become large as we move away from best-fit value, but without direct knowledge of underlying PDF, we cannot determine any confidence intervals from this

Feldman-Cousins method, or Plugin, is a “brute-force” approach to assigning a confidence interval


At each scan point of γ , perform these fits to toy:

- 1 Fix γ to scan point and generate 1000 toys
- 2 Perform fits to each toy, with γ both floating and fixed
- 3 Calculate $\Delta\chi_{\text{toy}}^2$

At each scan point, the fraction of toys with $\Delta\chi_{\text{toy}}^2 > \Delta\chi_{\text{data}}^2$ is equal to $1 - \text{CL}$, and the exact 68% confidence interval can then be obtained using an interpolation between points

Plugin/Feldman-Cousins method

LHCb average within 2σ of $D \rightarrow K^+ K^- \pi^+ \pi^-$ Plugin result
Combined fit shows good agreement between Plugin and Prob scans




Combined fit result: $\gamma = (53.9^{+9.5}_{-8.9})^\circ$


One of the most precise single measurements of γ !

Combining phase-space binned and integrated results

We can add phase-space integrated observables as a constraint:



γ vs δ_B^{DK}




γ vs r_B^{DK}

The global asymmetries contain useful information!

Combining phase-space binned and integrated results

Run Plugin with phase-space integrated constraints:



Final measurement: $\gamma = (52.6^{+8.5}_{-6.4})^\circ$

Conclusion

- ① Binned model-independent measurement of γ with $B^\pm \rightarrow [h^+ h^- \pi^+ \pi^-]_D h^\pm$ has been performed
 - Result: $\gamma = (53.9_{-8.9}^{+9.5})^\circ$
- ② Can also combine with existing phase-space integrated measurements
 - Result: $\gamma = (52.6_{-6.4}^{+8.5})^\circ$
- ③ 3σ tension in $D \rightarrow K^+ K^- \pi^+ \pi^-$ has reduced


Future prospects

- Statistically limited measurement, but s_i uncertainties are large
- $\pi^+\pi^-\pi^+\pi^-$ inputs will become more precise:
 - Current analysis uses 3 fb^{-1}
 - Future updates will use the full 20 fb^{-1} data set
- Minor improvements from BESIII are expected with $K^+K^-\pi^+\pi^-$:
 - Current analysis already uses 20 fb^{-1}
 - Charm mixing studies can improve s_i precision
- We believe the analysis is ready for approval to go to paper
 - You can find the TWiki [here](#)


Thanks for your attention!

Backup: CP fit toy studies

In toy studies biases in $D\pi$ observables are consistent with model-dependent analysis



$x_{\xi}^{D\pi}$




$y_{\xi}^{D\pi}$


$$D^0 \rightarrow K^+ K^- \pi^+ \pi^-$$

Backup: CP fit toy studies

Minor biases in x_{\pm}^{DK} are seen but can be corrected for...



x_{-}^{DK}




x_{+}^{DK}


$$D^0 \rightarrow K^+ K^- \pi^+ \pi^-$$

Backup: CP fit toy studies

...but y_{\pm}^{DK} pulls are now slightly asymmetric!



y_-^{DK}




y_+^{DK}


$$D^0 \rightarrow K^+ K^- \pi^+ \pi^-$$

Backup: Likelihood scan of CP observables

x_{\pm}^{DK} agree well between likelihood scan and Hesse approximation



x_{-}^{DK}




x_{+}^{DK}


$$D^0 \rightarrow \pi^+ \pi^- \pi^+ \pi^-$$

Backup: Likelihood scan of CP observables

y_{\pm}^{DK} diverges from Hesse approximation outside 1σ



y_{-}^{DK}




y_{+}^{DK}


$$D^0 \rightarrow \pi^+ \pi^- \pi^+ \pi^-$$

Backup: Interpretation toys

We can perform toy studies on the interpretation fit, but we do not expect these to behave very Gaussian...



$K^+K^-\pi^+\pi^-$



$\pi^+\pi^-\pi^+\pi^-$

γ pull distributions

Indeed, small but significant biases are observed!