


# $\gamma$ analysis update in $B^\pm \rightarrow (K^+ K^- \pi^+ \pi^-)_D K^\pm$ decays

Martin Tat

Oxford LHCb

28th June 2021



# Summary of last time

- $\gamma$  from  $B^\pm \rightarrow DK^\pm$ ,  $D \rightarrow K^+K^-\pi^+\pi^-$ , [arXiv:hep-ph/0611272](https://arxiv.org/abs/hep-ph/0611272)
- Model independent measurement with BESIII strong phase input
- Expected precision from signal-only study:  $\Delta\gamma = 12^\circ$
- $B$  candidate selection with BDT
- Initial mass fits

# Summary of analysis procedure

- 1 Perform global fit to fix yields and shapes
- 2 Separate  $B^\pm$  candidates by charge and into bins
- 3 Extract  $x_\pm$  and  $y_\pm$  with a simultaneous fit
- 4 Interpret  $x_\pm$  and  $y_\pm$  in terms of  $\gamma$

## 1 Backgrounds

- Mis-ID from  $D \rightarrow K\pi\pi\pi$
- Charmless backgrounds

## 2 Global mass fit


## 3 Toy studies

- Standard fits with 4 and 8 bins
- Biases in CP observables
- Systematics from  $c_i, s_i$


## 4 Summary and future work

# Mis-ID from $D \rightarrow K\pi\pi\pi$

- Mis-ID of  $D$  daughter  $\pi \rightarrow K$
- Peaks at much higher  $D$  mass
- Much higher branching ratio
- Similar background from  $D \rightarrow \pi\pi\pi\pi$  is negligible



(a)  $D$  invariant mass for  $KK\pi\pi$  and  $K\pi\pi\pi$




(b) Yield of  $KK\pi\pi$  and  $K\pi\pi\pi$  vs PIDK


# Charmless backgrounds

- Background from  $B^\pm \rightarrow KK\pi\pi h^\pm$
- Flight significance (FS) cut at 2
- Look in the lower  $D$  mass sideband  $m(D) = [1770 \text{ MeV}, 1820 \text{ MeV}]$
- Upper sideband contaminated with  $K\pi\pi\pi$
- Train a separate BDT without  $\chi_{\text{D TF}}^2$  to preserve  $D$  sidebands
- Overlap between samples without FS cut: 0.0%
- Overlap between samples with FS cut: 0.0%

# Total charmless background yield




(a) Without flight significance cut




(b) With flight significance cut

**Figure 2:** Charmless background in the  $D$  sideband

# Charmless backgrounds by charge



(a) Without flight significance cut




(b) With flight significance cut


**Figure 3:** Charmless background split by charge



# Charmless backgrounds by bins



(a) Without flight significance cut




(b) With flight significance cut

**Figure 4:** Charmless background split by bins

- Minimize statistical  $\gamma$  error
- Procedure:
  - ① Perform global fit to fix signal and background yields and PDF shapes
  - ② Generate 1000 toy datasets using the global fit parameters
  - ③ Fit for  $\gamma$
  - ④ Extract expected  $\gamma$  precision from  $\gamma$  distribution of toy datasets

# BDT cut optimization




**(a)**  $\gamma$  precision vs BDT cut for 4 bins

**(b)**  $\gamma$  precision vs BDT cut for 8 bins

A large, bold, black letter 'A' is centered on a gray background. The background features a grid of thin gray lines, including a horizontal line, a vertical line, and two diagonal lines forming an 'X' shape. The letter 'A' is composed of thick black strokes.


# Standard toy studies with 4 and 8 bins



(a) 4 bins

(b) 8 bins


**Figure 7:**  $\gamma$  precision in toy studies



(a) 4 bins

(b) 8 bins

**Figure 8:** CP observable that has bias




(a) 4 bins

(b) 8 bins

**Figure 9:** CP observable bias with increased statistics

# Systematics from $c_i$ , $s_i$



**Figure 10:** Systematic uncertainty from  $c_i$  and  $s_i$  with extrapolated BESIII statistics



# Summary and future work

- Summary:

- ① Global mass fit looks promising
- ②  $D \rightarrow K\pi\pi\pi$  and charmless backgrounds under control
- ③ Toy studies show no suspicious behaviour
- ④ Expected  $c_i$  and  $s_i$  systematics are not too large

- Next steps:

- ① Study semileptonic backgrounds in RapidSim
- ② Recalculate PID efficiencies with PIDCalib
- ③ Refit MC signal shapes
- ④ Rerun everything with Run 1 (finally!)