Tracking efficiencies from $J/\psi \to \mu^+\mu^-$

Rowina Caspary[1], Michel De Cian[2], Fabian Glaser[1], Stephanie Hansmann-Menzemer[1], Peilian Li[3], Maurice Morgenthaler[1], **Martin Tat**[1]

Heidelberg University[1], University of Manchester[2], UCAS[3]

5th March 2026

Part I

Part I: Tracking efficiencies with $J/\psi \to \mu^+\mu^-$

Introduction to tracking efficiencies

Strategy for tracking efficiency determination:

- Tag-and-probe method with $J/\psi \to \mu^+\mu^-$
- Combine efficiencies of two methods:
  - VeloMuon: SciFi efficiency
  - Downstream: Velo efficiency
  - Combined: Velo $\times$ SciFi efficiencies
- Cross check with long track efficiency with the MuonUT method
  - Main systematic

Tag-and-probe method



Figure by Rowina

- Fully reconstruct one muon from $J/\psi \to \mu^+\mu^-$
- Partially reconstruct the other muon
- Match hits in specific sub-detector with partially reconstructed track

$$\epsilon_{\text{track}} = \frac{N_{\text{matched}}}{N_{\text{matched}} + N_{\text{failed}}}$$

Overlap criteria in matching

Matching is done by comparing the hits of the probe track with a long track in each subdetector

Method	Velo	UT	SciFi	Muon
VeloMuon	$> 40\%$	–	–	$> 40\%$
Downstream	–	–	$> 40\%$	$> 40\%$
MuonUT	–	0 or $> 40\%$	–	$> 40\%$

Note:

- Matching was originally performed in a trigger line
- We now have functors for direct access to overlap fractions
- Requires rerunning reconstruction in 2024, as long-track information was not persisted

Motivation and strategy

Why do we care?

- Imperfect modelling of track reconstruction efficiency in MC
- Physics analyses that rely on efficiencies from MC must correct for this by reweighting

What do we provide?

1. Tracking efficiency ratio between data/MC of muon long tracks
   - Long track efficiency from product of Velo and SciFi efficiencies
   - 2D p–η binning
   - Dominant systematic uncertainty from MuonUT-Combined difference
2. Hadronic corrections for pions, kaons and protons
   - Mainly due to uncertainty in material budget
   - Using binning in η, with the same bin edges as for muons

Overview of previous work

Long history of previous work

- Method and results carefully documented in Rowina's thesis here
- Latest presentations in RTA-WP4 on
  - muon tracking efficiencies here
  - hadronic corrections here

Tracking efficiencies are determined using TrackCalib2

- GitLab repository here
- Mass fit done on tuples from AP to extract efficiencies
- Note: Analysts should <u>not</u> have to run TrackCalib2 themselves, unless the released numbers are not suitable (need different binning, etc)

Datasets considered

Datasets considered:

- 2024
  - Analysis of blocks 2, 1, 5, 6, 7, 8, *pp* ref mostly done
  - Currently waiting for large MC production to finish
  - Blocks 4 and 3 don't have MC yet
- 2025
  - Analysis of pre-TS data in progress
  - Will request 2025 post-TS MC soon
- 2026
  - Plan to look at soon

Example mass fit

Example of mass fit to extract tracking efficiency



- Efficiency is a free parameter in the fit
- Matched and failed samples simultaneously fitted to ensure correct uncertainties

Efficiency comparisons 2024

Comparison of SciFi tracking efficiencies in 2024 data blocks



Efficiency comparisons 2024

Comparison of Velo tracking efficiencies in 2024 data blocks



Efficiency comparisons 2024

Comparison of long tracking efficiencies in 2024 data blocks



General observations for 2024 results

General observations on tracking efficiencies:

1. All data sets show consistent trends
2. Performance seems to (slowly) drop over time

This is very encouraging! Several issues resolved, including:

1. Unusually high efficiencies in blocks 7/8
2. Missing binds when rerunning reconstruction
3. Large charge asymmetries in MuonUT samples

Efficiency comparisons 2025

Comparison of SciFi tracking efficiencies in 2025, compared with 2024



Efficiency comparisons 2025

Comparison of Velo tracking efficiencies in 2025, compared with 2024



Efficiency comparisons 2025

Comparison of long tracking efficiencies in 2025, compared with 2024



Efficiency comparisons 2025

2025 Data-MC discrepancy





Need to understand why MC underestimates the Velo efficiency in 2025!

Remarks on 2025 results

Remark on phase-space differences between 2024 and 2025:

- Probe muon p_T cut was at 500 MeV in 2024
- Loosened to 100 MeV in 2025
- Therefore cannot compare 2024 and 2025 directly

p_T cut comparison

Comparison of Velo tracking efficiencies with $p_T > 500\,\text{MeV}$ cut



p_T cut comparison

Comparison of SciFi tracking efficiencies with $p_T > 500\,\text{MeV}$ cut



Tracking efficiency data/MC ratio

Data/MC ratio





Data/MC ratio for Combined/MuonUT track efficiencies block 5

Tracking efficiency data/MC ratio

Data/MC ratio





Data/MC ratio for Combined/MuonUT track efficiencies 2025 pre-TS

General observations

General observations on tracking efficiency ratios:

1. Velo efficiencies in 2024 are well described, but underestimated in MC for 2025
2. Combined efficiency is mostly driven by SciFi efficiency
3. Generally there is a 1–2% discrepancy between Combined and MuonUT efficiencies, but effect is larger at low p and low η, so discrepancies are smaller in 2D (next slide)
4. Data–MC differ by almost 5% at low p and low η and needs to be understood

Block 1a vs 1b

Do we need to split Block 1 into Block 1a and Block 1b?

- SciFi T2L0Q0M1H0 FE was excluded on 17th August
  - Block 1a: Fill 9982–10012
  - Block 1b: Fill 10017–10056
- Integrated luminosities are 0.92 fb^{-1} and 0.31 fb^{-1}, respectively
- SciFi effectively has a hole in the 3rd quadrant

Block 1a vs 1b

Block 1a (top) vs 1b (bottom) for data (left) and MC (right)



Block 1a vs 1b

Integrated over all quadrants, for data (left) and MC (right)





Block 1a vs 1b

After initial study of block 1b MC:

- The "hole" in the SciFi is clearly visible when looking at tracks in the 3rd quadrant in data
- MC does not describe drop in efficiency well, will check with Simulation group
- The integrated numbers show reasonable agreement with MC, with a small drop in efficiency for a single η bin
- For now, tracking efficiencies are only provided for block 1a...
- ... but separate MC might be necessary for block 1b

2D binning

- Currently we have 1D bins in p and η with 10 bins each
- Using 10×10 2D bins is impossible
  - Insufficient statistics at high (low) p and low (high) η
  - But most bins <u>do</u> have sufficient statistics
- Strategy: Start with 10×10 bins and merge bins
  - Only consider MC, which is important for constraining the fit shape
  - Only consider VeloMuon yields, as Downstream efficiencies are close to 100% anyway and MuonUT is only used as a cross check
  - Try to merge bins along η at high p and vice versa
  - Aim for roughly $\gtrsim 100$ matched candidates in each bin
- The next plots contain many (tiny) numbers, apologies for this!

2D binning

Final 2D binning in p vs η with 72 bins:



2D binning


2024 Block 5 Data tracking efficiencies
0.69(7)
0.79(4)
0.951(9)
0.970(4)
0.957(2)
0.959(2)
0.951(1)
0.9504(7)
0.98(8)
0.964(6)
0.964(3)
0.958(2)
0.959(1)
0.9681(9)
0.9650(9)
0.9549(9)
0.97(1)
0.972(3)
0.965(2)
0.964(1)
0.9686(7)
0.9635(8)
0.9651(7)
0.9539(7)
0.956(8)
0.966(4)
0.9696(8)
0.9631(8)
0.955(1)
0.9629(9)
0.95(1)
0.962(2)
0.955(1)
0.956(1)
0.9695(9
0.9644(8)
0.957(1)
0.92(1)
0.959(1)
0.955(1)
0.9660(8)
0.956(1)
0.9644(8)
0.9594(9)
0.9618(8)
0.965(1)
0.95(2)
0.948(8)
0.962(1)
0.9593(9)
0.9653(8)
0.961(1)
0.9669(8)
0.944(3)
0.957(2)
0.953(1)
0.9569(8)
0.9589(9)
0.957(1)
0.962(1)
0.936(3)
0.9528(9)
0.949(1)
0.9515(9)
0.9384(8)
0.9457(9
0.9512(9)
0.952(1)
0.869(2)
0.895(1)
0.890(1)
0.907(1)
4.5
4.0
3.5
3.0
2.5
2.0
η
10^1
10^2
p [GeV/c]
1.00
0.98
0.96
0.94
0.92
0.90
0.88

2D binning


2024 Block 5 MC tracking efficiencies
0.9(5)
0.93(1)
0.974(5)
0.963(3)
0.968(2)
0.961(2)
0.963(2)
0.958(1)
0.92(1)
0.979(3)
0.979(2)
0.980(2)
0.976(1)
0.978(1)
0.976(1)
0.974(2)
0.965(5)
0.985(1)
0.980(2)
0.978(2)
0.980(1)
0.981(2)
0.977(1)
0.974(2)
0.975(4)
0.983(1)
0.984(1)
0.979(1)
0.977(1)
0.978(2)
0.965(3)
0.977(2)
0.972(2)
0.972(2)
0.982(2)
0.980(1)
0.975(3)
0.970(4)
0.971(2)
0.980(2)
0.986(1)
0.978(2)
0.982(2)
0.977(2)
0.976(2)
0.985(3)
0.963(3)
0.979(3)
0.976(2)
0.978(2)
0.983(2)
0.978(3)
0.977(2)
0.982(2)
0.981(3)
0.978(2)
0.980(2)
0.978(2)
0.980(3)
0.979(2)
0.963(3)
0.976(2)
0.973(2)
0.974(2)
0.967(2)
0.974(2)
0.978(2)
0.970(2)
0.912(3)
0.936(3)
0.937(3)
0.948(3)
η
4.5
4.0
3.5
3.0
2.5
2.0
10^1
10^2
p [GeV/c]
1.00
0.98
0.96
0.94
0.92
0.90
0.88

2D binning


2024 Block 5 Data/MC tracking efficiency ratios
0.7(4)
0.85(4)
0.98(1)
1.007(5)
0.989(3)
0.998(3)
0.987(2)
0.992(2)
1.06(9)
0.985(7)
0.984(3)
0.978(3)
0.982(2)
0.990(2)
0.989(2)
0.981(2)
1.01(1)
0.988(3)
0.984(3)
0.986(2)
0.988(1)
0.982(2)
0.988(2)
0.979(2)
0.981(9)
0.983(4)
0.985(1)
0.984(2)
0.977(2)
0.984(2)
0.98(1)
0.984(3)
0.982(2)
0.983(2)
0.988(2)
0.984(2)
0.982(3)
0.95(1)
0.987(3)
0.974(2)
0.979(2)
0.978(2)
0.982(2)
0.982(2)
0.986(2)
0.980(3)
0.99(2)
0.968(8)
0.986(3)
0.981(2)
0.982(2)
0.983(3)
0.989(2)
0.961(4)
0.976(4)
0.975(2)
0.977(2)
0.980(2)
0.976(3)
0.982(2)
0.972(4)
0.977(2)
0.975(2)
0.977(2)
0.970(2)
0.971(2)
0.973(2)
0.981(3)
0.953(4)
0.956(3)
0.949(3)
0.957(3)
4.5
4.0
3.5
3.0
2.5
2.0
η
10^1
10^2
p [GeV/c]
1.02
1.01
1.00
0.99
0.98
0.97
0.96
0.95

2D binning


2024 Block 5 Data/MC Combined vs MuonUT ratio difference
-0.3(6)
-0.1(2)
-0.01(7)
0.00(1)
-0.003(9)
0.009(6)
-0.006(4)
-0.002(3)
0.1(2)
-0.05(7)
-0.02(2)
-0.012(7
-0.022(5)
-0.010(4)
-0.006(3)
-0.005(3)
0.03(2)
-0.010(5)
-0.010(4)
-0.018(5
-0.014(3)
-0.005(5)
-0.013(3)
-0.018(4)
0.00(4)
-0.015(8)
-0.016(3)
-0.017(3)
0.004(6)
-0.04(1)
-0.01(3)
-0.021(8)
-0.006(5)
-0.004(4)
-0.003(3
-0.017(6)
0.003(6)
-0.05(2)
-0.013(7)
-0.023(4)
-0.009(4)
-0.012(4
-0.007(3)
-0.013(3)
-0.008(4)
-0.027(4)
0.06(9)
-0.03(1)
-0.014(4)
-0.008(4)
-0.015(3)
-0.017(5
-0.005(3)
0.2(7)
-0.01(1)
-0.026(5)
-0.008(4)
-0.020(2)
-0.022(5
-0.015(4)
0.00(3)
-0.020(5)
0.001(5)
-0.003(5)
-0.014(4)
-0.017(4)
-0.032(5
-0.019(6)
-0.02(1)
-0.000(7)
-0.009(7)
-0.006(6)
4.5
4.0
3.5
3.0
2.5
2.0
η
10^1
10^2
p [GeV/c]
0.03
0.02
0.01
0.00
−0.01
−0.02
−0.03

2D binning

Note: After some feedback, the 2D binning will be slightly modified

- Merge bins 64 and 65
- Add a single high p bin and a low η bin
- These are bins with low statistics, but could be useful for some analyses to have





Combined vs MuonUT systematic

Combined vs MuonUT systematic strategy?

- Clearly depends on kinematics, so we should use 2D binning
- It's a method systematic, should not depend on dataset with same μ
- Our current strategy:
  1. Determine difference in data/MC ratio for all data sets with 2D binning
  2. Do weighted average of the difference for blocks 5/6 and 7/8
  3. For bins with a discrepancy above $1\,\sigma$, assign the difference from zero as a systematic
  4. Use blocks 5/6 as proxy for blocks 1/2 due to bug in MuonUT trigger line (see !4047 and !4050)
  5. Systematic will be small in most bins, but a small number of bins have a 1%–2% systematic, in rare cases $> 3\%$

Release of tracking efficiencies

How should analysts use these results?

- Binning schemes are defined in JSON files
- Data/MC ratio in 2D bins provided for physics analysis, also as JSON
- Version 0 is available as a GitLab project here
- Version 1 will come soon
  1. Waiting for large MC production to finish
  2. Separate μ^+ and μ^- results as well
  3. Updated 2D binning scheme

Part II: Hadronic corrections

Introduction to hadronic corrections

- About 11% of kaons and 14% of pions cannot be reconstructed due to hadronic interactions with the detector material
- Modifies the track reconstruction efficiency of long muon tracks

$$\varepsilon^{reco}_{hadron} = \varepsilon^{reco}_{muon} \times \rho^{had.Int.}_{hadron}$$

- Simulation of hadronic interactions has uncertainties due to the material distribution, which is hard to quantify
- In Run 2: Assumed 10% uncertainty $\rightarrow$ 1.1% and 1.4% uncertainty on the kaon and pion tracking efficiency, respecively
- This work: Measure tracking efficiency correction factors $\rho^{had.Int.,Data}_{hadron} / \rho^{had.Int.,MC}_{hadron}$ from data

Method overview

- Measure the relative efficiency of $D^0 \to K^-\pi^+$ and $D^0 \to K^-\pi^+\pi^+\pi^-$ in data (blocks 1,2,5-8) and simulation (Sim10g)
- The total reconstruction&selection efficiency $\varepsilon^{reco\&sel}$ is factorised as

$$\varepsilon^{reco\&sel} = \varepsilon^{acc}\varepsilon^{reco}\varepsilon^{HLT1}\varepsilon^{HLT2}\varepsilon^{offline} \equiv \varepsilon^{reco}\varepsilon^{nuis}$$

- Can isolate hadronic correction for two pions $\rho^{had.Int.}_{\pi\pi}$

$$\frac{\varepsilon^{reco\&sel}_{K\pi\pi\pi}}{\varepsilon^{reco\&sel}_{K\pi}} = \frac{\varepsilon^{reco}_{\mu\mu\mu\mu}\varepsilon^{nuis}_{K\pi\pi\pi}\rho^{had.Int}_{K\pi\pi\pi}}{\varepsilon^{reco}_{\mu\mu}\varepsilon^{nuis}_{K\pi}\rho^{had.Int}_{K\pi}} = \frac{\varepsilon^{reco}_{\mu\mu\mu\mu}\varepsilon^{nuis}_{K\pi\pi\pi}}{\varepsilon^{reco}_{\mu\mu}\varepsilon^{nuis}_{K\pi}}\rho^{had.Int}_{\pi\pi}$$

- Use this information to infer correction for a single pion track $\rho^{had.Int}_{\pi}$

Kinematic reweighting

- High-p_T π^+ in $D^0 \to K^-\pi^+\pi^+\pi^-$ has most similar kinematics to π^+ in $D^0 \to K^-\pi^+$
- For $K^-\pi^+_{(\text{high}p_T)}$ to cancel, ensure cover same phase-space by reweighting:

1. Reweight $D^0 \to K^-\pi^+\pi^+\pi^-$ MC to sWeighted data in (p, η) for $K^-, \pi^+_{\text{high}p_T}$ and ϕ for each track
2. Reweight sWeighted $D^0 \to K^-\pi^+$ data to sWeighted $D^0 \to K^-\pi^+\pi^+\pi^-$ data in (p, η) for $K^-, \pi^+_{(\text{high}p_T)}$
3. Reweight $D^0 \to K^-\pi^+$ MC to sWeighted & kin. reweighted $D^0 \to K^-\pi^+$ data

Data/MC comparison for $D^0 \to K^-\pi^+\pi^+\pi^-$



Comparison of data and simulation of $D^0 \to K^-\pi^+\pi^+\pi^-$ for K^- (top) and $\pi^+_{\text{high}\ p_T}$ (bottom) in p (left) and η (centre) and ϕ (right).

Data/MC comparison for $D^0 \to K^-\pi^+\pi^+\pi^-$



Comparison of data and simulation of $D^0 \to K^-\pi^+\pi^+\pi^-$ for $\pi^+_{\text{low } p_T}$ (top) and π^- (bottom) in p (left) and η (centre) and ϕ (right).

Data/MC comparison for $D^0 \to K^-\pi^+$



Comparison of data and simulation of $D^0 \to K\pi$ and $D^0 \to K\pi\pi\pi$ for K^- (top) and $\pi^+/\pi^+_{\text{high } p_T}$ (bottom) in p (left) and η (centre) and ϕ (right).

Measure $\rho_{\pi\pi}^{had.Int}$ in data and MC

- Measure $\rho_{\pi\pi}^{had.Int.}$ in data and MC

$$\rho_{\pi\pi}^{had.Int,Data} = \frac{N_{K\pi\pi\pi}^{fit}}{N_{K\pi}^{fit}} \frac{\mathcal{B}(D^0 \to K^-\pi^+)}{\mathcal{B}(D^0 \to K^-\pi^+\pi^+\pi^-)} \frac{\varepsilon_{\mu\mu}^{reco}}{\varepsilon_{\mu\mu\mu\mu}^{reco}} \frac{\varepsilon_{K\pi}^{nuis}}{\varepsilon_{K\pi\pi\pi}^{nuis}}$$

$$\rho_{\pi\pi}^{had.Int,MC} = \frac{N_{K\pi\pi\pi}^{sel}}{N_{K\pi\pi\pi}^{gen}} \frac{N_{K\pi}^{gen}}{N_{K\pi}^{sel}} \frac{\varepsilon_{\mu\mu}^{reco}}{\varepsilon_{\mu\mu\mu\mu}^{reco}} \frac{\varepsilon_{K\pi}^{nuis}}{\varepsilon_{K\pi\pi\pi}^{nuis}}$$

- Calculate data/MC correction factor
$\rho_{\pi\pi}^{had.Int.,Data/MC} = \rho_{\pi\pi}^{had.Int.,Data} / \rho_{\pi\pi}^{had.Int.,MC}$

- Use the following additional inputs:
  - Standard (muon) tracking corrections from TrackCalib2
  - $\mathcal{B}(D^0 \to K^-\pi^+\pi^+\pi^-)/\mathcal{B}(D^0 \to K^-\pi^+) = 2.083 \pm 0.031$ from PDG
  - Assume $\frac{\varepsilon_{K\pi}^{nuis,Data} / \varepsilon_{K\pi\pi\pi}^{nuis,Data}}{\varepsilon_{K\pi}^{nuis,MC} / \varepsilon_{K\pi}^{nuis,MC}} = 1$

Phase-space integrated hadronic correction

- Single fit to $D^0 \to K^-\pi^+$ and $D^0 \to K^-\pi^+\pi^+\pi^-$ data
- Kinematic corrections applied to $D^0 \to K^-\pi^+$ data

$$\rho_{\pi\pi}^{had.Int.,Data/MC} = 100.88 \pm 1.51\%$$

- Correction for single hadron track

$$\rho_{\pi}^{had.Int.,Data/MC} = 100.44 \pm 0.75\%$$

- Uncertainty dominated by $\mathcal{B}(D^0 \to K^-\pi^+\pi^+\pi^-)/\mathcal{B}(D^0 \to K^-\pi^+)$



$D^0 \to K^-\pi^+$ (top) and $D^0 \to K^-\pi^+\pi^+\pi^-$ (bottom), 2024 Block 1

Hadronic correction as function of η

- Check for dependence of the hadronic correction factor on η
- Dependence on $K^-\pi^+_{(\text{high } p_T)}$ cancels in the ratio...
- ... but not $\pi^+_{\text{low } p_T}$ and π^- from $D^0 \to K^-\pi^+\pi^+\pi^-$
- $\to$ Need to bin in $\eta(\pi^+_{\text{low } p_T})$ and $\eta(\pi^-)$
  - Use same bin edges as muon tracking efficiency

Hadronic correction as function of η

- Measure $\rho_{\pi\pi}^{had.Int}$ in bins of (η, η)
- Obtain 1D distribution by assuming factorisation $\rho_{\pi\pi}^{had.Int.} = \rho_{\pi}^{had.Int.} \rho_{\pi}^{had.Int.}$
- In practice: Perform a χ^2-fit to get best $\rho_{\pi}^{had.Int.}(\eta)$ accounting for the uncertainty in each bin



Hadronic correction factor for two pions (left) in 2d bins of η (left) and hadronic correction factor for a single pion (right).

Hadronic correction as function of p or ϕ

- Can also bin in (p, p) or (ϕ, ϕ) and measure the hadronic correction factor as function of p or ϕ
- As expected, no dependence on p
- Drop around $\phi \sim -\pi$ observed and is being investigated



Hadronic correction factor for a single track in bin sof p (left) and ϕ (right).

Cancellation of $K^-\pi^+_{(\text{high } p_T)}$

- So far, assume perfect cancellation between $K^-\pi^+$ from $D^0 \to K^-\pi^+$ and $K^-\pi^+_{\text{high } p_T}$ from $D^0 \to K^-\pi^+\pi^+\pi^-$
- Just observed some non-cancellation for at the edges of p, η and ϕ

$\rightarrow$ Need to revisit the kinematic reweighting

- Structures seen as function η and ϕ might be (partially) due to this



Ratio of data/MC for $D^0 \to K^-\pi^+\pi^+\pi^-$ (red), $D^0 \to K^-\pi^+$ (yellow) and the double-ratio (black) of the π^+ cancelling between the two decays in bins of p (left), η (centre) and ϕ (right).

Summary and next steps

Tracking efficiencies with $J/\psi \to \mu^+\mu^-$

- Muon tracking efficiencies v0 for 2024 can now be used by analysts
- Issues to follow up on:
  - Block 1b MC
  - 2025 Velo efficiencies
- Waiting for large MC production before next release

Hadronic corrections

- Hadronic correction factors using 2024 data and MC have been measured using $D^0 \to K^-\pi^+$ and $D^0 \to K^-\pi^+\pi^+\pi^-$
- Well modelled in MC, and corrections provided in η bins
- Uncertainties are about 1% in most bins of η
- Still need to investigate
  - data/MC agreement in p, η and ϕ
  - cancellation of $K^-\pi^+$, nuisance effects

Thanks for your attention!

Backup

Uncertainties for phase-space integrated result

Source	$\Delta\rho_\pi^{had.Int.,Data/MC}$
$D \to K\pi$ simulation sample	0.04%
$D \to K3\pi$ simulation sample	0.07%
$D \to K\pi$ Data sample	0.01%
$D \to K3\pi$ Data sample	0.01%
Tracking weights	0.12%
$\mathcal{B}(D^0 \to K^-\pi^+\pi^+\pi^-)/\mathcal{B}(D^0 \to K^-\pi^+)$	0.74%
HLT selection	x.xx
Offline selection	x.xx
Total	0.76%

- Dominated by uncertainty on the branching ratio
- Uncertainty due to tracking weights from bootstrapping
- Expect HLT and offline uncertainties to cancel in the ratio