



Six-method fit comparison — NaturalEarth / natural_earth_10m_coastline_08619b5b423b_w000_9b30514c

shared MSE tolerance = $1.0\text{e-}04$


Ours v16 — FAIL
input MSE= $3.23\text{e-}04$ | ref MSE= $3.68\text{e-}04$ | K=56
full method=8.46 ms | network=6.49 ms




Park & Lee 2007 (DOM adaptation) — FAIL
input MSE= $3.57\text{e-}04$ | ref MSE= $4.04\text{e-}04$ | K=56
full method=89.95 ms




Liang et al. 2017 (feature-IKI adaptation) — FAIL
input MSE= $2.80\text{e-}04$ | ref MSE= $3.33\text{e-}04$ | K=56
full method=76.19 ms




Dung & Tjahjowidodo 2017 (serial adaptation) — FAIL
input MSE= $6.14\text{e-}04$ | ref MSE= $6.49\text{e-}04$ | K=34
full method=587.22 ms



Kang 2015 (ADMM adaptation) — FAIL
input MSE= $3.80\text{e-}02$ | ref MSE= $3.92\text{e-}02$ | K=1
full method=14.18 ms



Luo et al. 2022 (l-infinity,1 + DE adaptation) — FAIL
input MSE= $1.20\text{e-}02$ | ref MSE= $1.22\text{e-}02$ | K=6
full method=103.22 ms



Original reference curve Final B-spline Control polygon Control vertices Internal-knot locations 192 input samples