



Six-method fit comparison — NaturalEarth / natural_earth_10m_coastline_0b2f7967fa0b_w000_0b2f7967
shared MSE tolerance = $1.0\text{e-}04$


Ours v16 — PASS
input MSE= $2.39\text{e-}05$ | ref MSE= $2.37\text{e-}05$ | K=15
full method=7.97 ms | network=6.68 ms




Park & Lee 2007 (DOM adaptation) — PASS
input MSE= $7.47\text{e-}05$ | ref MSE= $7.65\text{e-}05$ | K=11
full method=14.35 ms




Liang et al. 2017 (feature-IKI adaptation) — PASS
input MSE= $9.65\text{e-}05$ | ref MSE= $9.81\text{e-}05$ | K=10
full method=10.37 ms



Dung & Tjahjowidodo 2017 (serial adaptation) — FAIL
input MSE= $5.56\text{e-}04$ | ref MSE= $5.65\text{e-}04$ | K=6
full method=90.84 ms



Kang 2015 (ADMM adaptation) — PASS
input MSE= $8.36\text{e-}05$ | ref MSE= $8.57\text{e-}05$ | K=9
full method=304.98 ms



Luo et al. 2022 (l-infinity,1 + DE adaptation) — FAIL
input MSE= $4.38\text{e-}03$ | ref MSE= $4.42\text{e-}03$ | K=3
full method=641.22 ms

