



Six-method fit comparison — NaturalEarth / natural_earth_10m_coastline_e046027da0e0_w000_e046027d

shared MSE tolerance = $1.0\text{e-}04$


Ours v16 — PASS
input MSE= $2.52\text{e-}05$ | ref MSE= $2.46\text{e-}05$ | K=19
full method=8.01 ms | network=6.68 ms




Park & Lee 2007 (DOM adaptation) — PASS
input MSE= $5.56\text{e-}05$ | ref MSE= $5.63\text{e-}05$ | K=12
full method=18.30 ms




Liang et al. 2017 (feature-IKI adaptation) — PASS
input MSE= $7.70\text{e-}05$ | ref MSE= $7.75\text{e-}05$ | K=11
full method=11.31 ms




Dung & Tjahjowidodo 2017 (serial adaptation) — FAIL
input MSE= $5.38\text{e-}04$ | ref MSE= $5.42\text{e-}04$ | K=6
full method=109.03 ms



Kang 2015 (ADMM adaptation) — PASS
input MSE= $5.54\text{e-}05$ | ref MSE= $5.60\text{e-}05$ | K=10
full method=304.20 ms



Luo et al. 2022 (l-infinity,1 + DE adaptation) — FAIL
input MSE= $1.04\text{e-}03$ | ref MSE= $1.04\text{e-}03$ | K=4
full method=648.64 ms



Original reference curve
Final B-spline
Control polygon
Control vertices
Internal-knot locations
192 input samples