



Six-method fit comparison — UJI / uji2_02446_stroke_01

shared MSE tolerance = $1.0\text{e-}04$


Ours v16 — PASS
input MSE= $1.33\text{e-}05$ | ref MSE= $2.08\text{e-}05$ | K=5
full method=7.97 ms | network=6.85 ms




Park & Lee 2007 (DOM adaptation) — PASS
input MSE= $1.60\text{e-}05$ | ref MSE= $3.10\text{e-}05$ | K=0
full method=2.38 ms




Liang et al. 2017 (feature-IKI adaptation) — PASS
input MSE= $5.36\text{e-}06$ | ref MSE= $1.12\text{e-}05$ | K=4
full method=4.35 ms




Dung & Tjahjowidodo 2017 (serial adaptation) — PASS
input MSE= $1.60\text{e-}05$ | ref MSE= $3.10\text{e-}05$ | K=0
full method=2.75 ms



Kang 2015 (ADMM adaptation) — PASS
input MSE= $1.60\text{e-}05$ | ref MSE= $3.10\text{e-}05$ | K=0
full method=4.65 ms



Luo et al. 2022 (l-infinity,1 + DE adaptation) — PASS
input MSE= $3.94\text{e-}06$ | ref MSE= $9.00\text{e-}06$ | K=3
full method=646.70 ms



Legend:
— Original reference curve
— Final B-spline
- - Control polygon
■ Control vertices
◆ Internal-knot locations
• 192 input samples