



# Six-method fit comparison — UJI / uji2\_02751\_stroke\_01

## shared MSE tolerance = 1.0e-04


**Ours v16 — PASS**  
input MSE=2.17e-05 | ref MSE=1.38e-05 | K=5  
full method=7.99 ms | network=6.72 ms




**Park & Lee 2007 (DOM adaptation) — PASS**  
input MSE=8.66e-05 | ref MSE=1.39e-04 | K=2  
full method=4.39 ms




**Liang et al. 2017 (feature-IKI adaptation) — PASS**  
input MSE=5.70e-06 | ref MSE=1.30e-05 | K=4  
full method=4.71 ms




**Dung & Tjahjowidodo 2017 (serial adaptation) — PASS**  
input MSE=4.23e-05 | ref MSE=5.81e-05 | K=2  
full method=32.88 ms



**Kang 2015 (ADMM adaptation) — FAIL**  
input MSE=1.30e-04 | ref MSE=1.99e-04 | K=0  
full method=4.86 ms



**Luo et al. 2022 (l-infinity,1 + DE adaptation) — PASS**  
input MSE=2.59e-05 | ref MSE=4.69e-05 | K=2  
full method=640.50 ms



Legend:  
— Original reference curve  
• 192 input samples  
— Final B-spline  
— Control polygon  
— Control vertices  
— Internal-knot locations