



Six-method fit comparison — USGS / usgs_tnm_contours_large_scale_0a876e98cab7_w000_0a876e98

shared MSE tolerance = $1.0\text{e-}04$


Ours v16 — PASS
input MSE= $2.06\text{e-}05$ | ref MSE= $2.05\text{e-}05$ | K=15
full method=7.75 ms | network=6.49 ms




Park & Lee 2007 (DOM adaptation) — PASS
input MSE= $6.92\text{e-}05$ | ref MSE= $6.95\text{e-}05$ | K=14
full method=17.82 ms




Liang et al. 2017 (feature-IKI adaptation) — PASS
input MSE= $9.60\text{e-}05$ | ref MSE= $9.65\text{e-}05$ | K=10
full method=10.15 ms



Dung & Tjahjowidodo 2017 (serial adaptation) — FAIL
input MSE= $4.52\text{e-}04$ | ref MSE= $4.55\text{e-}04$ | K=5
full method=86.68 ms



Kang 2015 (ADMM adaptation) — PASS
input MSE= $7.12\text{e-}05$ | ref MSE= $7.17\text{e-}05$ | K=11
full method=310.11 ms



Luo et al. 2022 (l-infinity,1 + DE adaptation) — FAIL
input MSE= $3.55\text{e-}04$ | ref MSE= $3.56\text{e-}04$ | K=6
full method=643.93 ms

