



Six-method fit comparison — USGS / usgs_tnm_contours_large_scale_195e157634b4_w000_195e1576

shared MSE tolerance = $1.0\text{e-}04$


Ours v16 — PASS
input MSE= $2.23\text{e-}05$ | ref MSE= $2.22\text{e-}05$ | K=13
full method=7.83 ms | network=6.48 ms




Park & Lee 2007 (DOM adaptation) — PASS
input MSE= $3.94\text{e-}05$ | ref MSE= $3.96\text{e-}05$ | K=18
full method=22.51 ms




Liang et al. 2017 (feature-IKI adaptation) — PASS
input MSE= $6.67\text{e-}05$ | ref MSE= $6.72\text{e-}05$ | K=10
full method=10.05 ms




Dung & Tjahjowidodo 2017 (serial adaptation) — FAIL
input MSE= $8.08\text{e-}04$ | ref MSE= $8.12\text{e-}04$ | K=5
full method=91.94 ms



Kang 2015 (ADMM adaptation) — PASS
input MSE= $6.71\text{e-}05$ | ref MSE= $6.73\text{e-}05$ | K=9
full method=302.81 ms



Luo et al. 2022 (l-infinity,1 + DE adaptation) — FAIL
input MSE= $2.71\text{e-}04$ | ref MSE= $2.72\text{e-}04$ | K=7
full method=644.64 ms



Original reference curve | Final B-spline | Control polygon | Control vertices | Internal-knot locations