



# Six-method fit comparison — USGS / usgs\_tnm\_contours\_large\_scale\_204e341b669c\_w000\_204e341b

shared MSE tolerance =  $1.0\text{e-}04$


**Ours v16 — PASS**  
input MSE= $2.31\text{e-}05$  | ref MSE= $2.28\text{e-}05$  | K=8  
full method=7.83 ms | network=6.67 ms




**Park & Lee 2007 (DOM adaptation) — PASS**  
input MSE= $8.25\text{e-}05$  | ref MSE= $8.28\text{e-}05$  | K=6  
full method=8.54 ms




**Liang et al. 2017 (feature-IKI adaptation) — PASS**  
input MSE= $8.79\text{e-}05$  | ref MSE= $8.82\text{e-}05$  | K=6  
full method=6.01 ms




**Dung & Tjahjowidodo 2017 (serial adaptation) — FAIL**  
input MSE= $5.34\text{e-}04$  | ref MSE= $5.36\text{e-}04$  | K=3  
full method=50.88 ms



**Kang 2015 (ADMM adaptation) — PASS**  
input MSE= $2.10\text{e-}05$  | ref MSE= $2.11\text{e-}05$  | K=7  
full method=302.73 ms



**Luo et al. 2022 (l-infinity,1 + DE adaptation) — PASS**  
input MSE= $4.46\text{e-}06$  | ref MSE= $4.49\text{e-}06$  | K=12  
full method=667.09 ms



Original reference curve  
Final B-spline  
Control polygon  
Control vertices  
Internal-knot locations  
192 input samples