



Six-method fit comparison — USGS / usgs_tnm_contours_large_scale_3c08ba8ede56_w000_3c08ba8e

shared MSE tolerance = $1.0\text{e-}04$


Ours v16 — PASS
input MSE= $6.31\text{e-}05$ | ref MSE= $6.99\text{e-}05$ | K=56
full method=8.47 ms | network=6.45 ms




Park & Lee 2007 (DOM adaptation) — PASS
input MSE= $9.04\text{e-}05$ | ref MSE= $9.64\text{e-}05$ | K=54
full method=85.05 ms




Liang et al. 2017 (feature-IKI adaptation) — PASS
input MSE= $9.96\text{e-}05$ | ref MSE= $1.06\text{e-}04$ | K=48
full method=60.56 ms




Dung & Tjahjowidodo 2017 (serial adaptation) — FAIL
input MSE= $5.05\text{e-}04$ | ref MSE= $5.21\text{e-}04$ | K=26
full method=480.93 ms



Kang 2015 (ADMM adaptation) — FAIL
input MSE= $2.90\text{e-}02$ | ref MSE= $2.91\text{e-}02$ | K=1
full method=69.21 ms



Luo et al. 2022 (l-infinity,1 + DE adaptation) — FAIL
input MSE= $3.68\text{e-}03$ | ref MSE= $3.72\text{e-}03$ | K=6
full method=425.23 ms



— Original reference curve — Final B-spline - - - Control polygon ■ Control vertices ◆ Internal-knot locations
192 input samples