



Six-method fit comparison — USGS / usgs_tnm_contours_large_scale_5f2e03d7c054_w000_5f2e03d7

shared MSE tolerance = $1.0\text{e-}04$


Ours v16 — PASS
input MSE= $2.08\text{e-}05$ | ref MSE= $2.08\text{e-}05$ | K=36
full method=8.27 ms | network=6.54 ms




Park & Lee 2007 (DOM adaptation) — PASS
input MSE= $9.86\text{e-}05$ | ref MSE= $9.92\text{e-}05$ | K=29
full method=38.73 ms




Liang et al. 2017 (feature-IKI adaptation) — PASS
input MSE= $8.96\text{e-}05$ | ref MSE= $9.05\text{e-}05$ | K=21
full method=22.33 ms




Dung & Tjahjowidodo 2017 (serial adaptation) — FAIL
input MSE= $4.95\text{e-}04$ | ref MSE= $4.97\text{e-}04$ | K=10
full method=160.11 ms



Kang 2015 (ADMM adaptation) — FAIL
input MSE= $2.25\text{e-}03$ | ref MSE= $2.27\text{e-}03$ | K=7
full method=231.26 ms



Luo et al. 2022 (l-infinity,1 + DE adaptation) — FAIL
input MSE= $3.35\text{e-}04$ | ref MSE= $3.37\text{e-}04$ | K=12
full method=663.07 ms



Legend:
— Original reference curve
— Final B-spline
- - - Control polygon
■ Control vertices
◆ Internal-knot locations
• 192 input samples