



Six-method fit comparison — USGS / usgs_tnm_contours_large_scale_80e7ca044177_w000_80e7ca04

shared MSE tolerance = $1.0\text{e-}04$


Ours v16 — PASS
input MSE= $8.13\text{e-}05$ | ref MSE= $8.29\text{e-}05$ | K=46
full method=8.25 ms | network=6.47 ms




Park & Lee 2007 (DOM adaptation) — PASS
input MSE= $8.84\text{e-}05$ | ref MSE= $9.21\text{e-}05$ | K=45
full method=67.02 ms




Liang et al. 2017 (feature-IKI adaptation) — PASS
input MSE= $9.69\text{e-}05$ | ref MSE= $1.02\text{e-}04$ | K=39
full method=46.83 ms




Dung & Tjahjowidodo 2017 (serial adaptation) — FAIL
input MSE= $6.26\text{e-}04$ | ref MSE= $6.41\text{e-}04$ | K=19
full method=365.72 ms



Kang 2015 (ADMM adaptation) — FAIL
input MSE= $1.59\text{e-}02$ | ref MSE= $1.60\text{e-}02$ | K=1
full method=69.16 ms



Luo et al. 2022 (l-infinity,1 + DE adaptation) — FAIL
input MSE= $9.20\text{e-}03$ | ref MSE= $9.25\text{e-}03$ | K=4
full method=471.24 ms



Original reference curve Final B-spline Control polygon Control vertices Internal-knot locations
192 input samples