



Six-method fit comparison — USGS / usgs_tnm_contours_large_scale_89dbac21c8c3_w000_89dbac21

shared MSE tolerance = $1.0\text{e-}04$


Ours v16 — PASS
input MSE= $1.27\text{e-}05$ | ref MSE= $1.33\text{e-}05$ | K=45
full method=8.44 ms | network=6.48 ms




Park & Lee 2007 (DOM adaptation) — PASS
input MSE= $9.45\text{e-}05$ | ref MSE= $9.58\text{e-}05$ | K=26
full method=34.09 ms




Liang et al. 2017 (feature-IKI adaptation) — PASS
input MSE= $8.05\text{e-}05$ | ref MSE= $8.22\text{e-}05$ | K=17
full method=17.52 ms




Dung & Tjahjowidodo 2017 (serial adaptation) — FAIL
input MSE= $2.27\text{e-}04$ | ref MSE= $2.29\text{e-}04$ | K=11
full method=204.95 ms



Kang 2015 (ADMM adaptation) — FAIL
input MSE= $2.38\text{e-}04$ | ref MSE= $2.41\text{e-}04$ | K=11
full method=311.63 ms



Luo et al. 2022 (l-infinity,1 + DE adaptation) — FAIL
input MSE= $2.18\text{e-}04$ | ref MSE= $2.20\text{e-}04$ | K=12
full method=662.77 ms



Legend:
— Original reference curve
— Final B-spline
- - Control polygon
■ Control vertices
◆ Internal-knot locations
• 192 input samples