



Six-method fit comparison — USGS / usgs_tnm_contours_large_scale_b9fab0458024_w000_b9fab045

shared MSE tolerance = 1.0e-04


Ours v16 — PASS
input MSE=2.55e-05 | ref MSE=2.52e-05 | K=15
full method=7.73 ms | network=6.45 ms




Park & Lee 2007 (DOM adaptation) — PASS
input MSE=8.02e-05 | ref MSE=8.05e-05 | K=15
full method=18.69 ms




Liang et al. 2017 (feature-IKI adaptation) — PASS
input MSE=8.15e-05 | ref MSE=8.19e-05 | K=9
full method=9.04 ms




Dung & Tjahjowidodo 2017 (serial adaptation) — FAIL
input MSE=3.46e-04 | ref MSE=3.47e-04 | K=5
full method=91.88 ms



Kang 2015 (ADMM adaptation) — PASS
input MSE=4.85e-05 | ref MSE=4.86e-05 | K=9
full method=293.76 ms



Luo et al. 2022 (l-infinity,1 + DE adaptation) — PASS
input MSE=9.65e-05 | ref MSE=9.69e-05 | K=10
full method=652.14 ms



Original reference curve Final B-spline Control polygon Control vertices Internal-knot locations
192 input samples