



Six-method fit comparison — USGS / usgs_tnm_contours_large_scale_f80b5d3e52bd_w000_f80b5d3e

shared MSE tolerance = $1.0\text{e-}04$


Ours v16 — PASS
input MSE= $2.35\text{e-}05$ | ref MSE= $2.33\text{e-}05$ | K=12
full method=7.69 ms | network=6.55 ms




Park & Lee 2007 (DOM adaptation) — PASS
input MSE= $2.74\text{e-}05$ | ref MSE= $2.76\text{e-}05$ | K=14
full method=17.49 ms




Liang et al. 2017 (feature-IKI adaptation) — PASS
input MSE= $8.03\text{e-}05$ | ref MSE= $8.07\text{e-}05$ | K=7
full method=7.02 ms




Dung & Tjahjowidodo 2017 (serial adaptation) — FAIL
input MSE= $5.68\text{e-}04$ | ref MSE= $5.70\text{e-}04$ | K=4
full method=72.33 ms



Kang 2015 (ADMM adaptation) — PASS
input MSE= $9.42\text{e-}05$ | ref MSE= $9.45\text{e-}05$ | K=7
full method=296.42 ms



Luo et al. 2022 (l-infinity,1 + DE adaptation) — FAIL
input MSE= $2.21\text{e-}04$ | ref MSE= $2.22\text{e-}04$ | K=3
full method=633.80 ms



Legend:
Original reference curve (blue line)
Final B-spline (blue line)
Control polygon (orange dashed line)
Control vertices (orange squares)
Internal-knot locations (purple diamonds)
192 input samples (light blue dots)