

# Fundamentals of Artificial Intelligence

## Laboratory

Dr. Mauro Dragoni

Department of Information Engineering and Computer Science  
Academic Year 2022/2023

## Exercise 8.1 - Which of the following are correct?

- a.  $\text{False} \models \text{True}$ . This is true because False has no models and hence entails every sentence and because True is true in all models and hence is entailed by every sentence.
- b.  $\text{True} \models \text{False}$ . This is false.
- c.  $(A \wedge B) \models (A \Leftrightarrow B)$ . This is true because the left-hand side has exactly one model that is one of the two models of the right-hand side.
- d.  $A \Leftrightarrow B \models A \vee B$ . This is false because one of the models of  $A \Leftrightarrow B$  has both A and B false, which does not satisfy  $A \vee B$ .
- e.  $A \Leftrightarrow B \models \neg A \vee B$ . This is true because the RHS is  $A \Rightarrow B$ , one of the conjuncts in the definition of  $A \Leftrightarrow B$ .
- f.  $(A \wedge B) \Rightarrow C \models (A \Rightarrow C) \vee (B \Rightarrow C)$ . This is true because the RHS is false only when both disjuncts are false, i.e., when A and B are true and C is false, in which case the LHS is also false. This may seem counterintuitive, and would not hold if  $\Rightarrow$  is interpreted as “causes.”
- g.  $(A \vee B) \wedge (\neg C \vee \neg D \vee E) \models (A \vee B)$ . This is true; removing a conjunct only allows more models.
- h.  $(A \vee B) \wedge (\neg C \vee \neg D \vee E) \models (A \vee B) \wedge (\neg D \vee E)$ . This is false; removing a disjunct allows fewer models.
- i.  $(A \vee B) \wedge \neg(A \Rightarrow B)$  is satisfiable. This is satisfiable; model has A and  $\neg B$ .
- j.  $(A \Leftrightarrow B) \wedge (\neg A \vee B)$  is satisfiable. This is satisfiable; RHS is entailed by LHS so models are those of  $A \Leftrightarrow B$ .

## Exercise 8.2 - Homework

Using truth tables, verify the following equivalences:

a.  $P \wedge Q \Leftrightarrow Q \wedge P$

b.  $P \vee Q \Leftrightarrow Q \vee P$

c.  $P \wedge (Q \wedge R) \Leftrightarrow (P \wedge Q) \wedge R$

d.  $P \vee (Q \vee R) \Leftrightarrow (P \vee Q) \vee R$

## Exercise 8.3

Consider a vocabulary with only four propositions, A, B, C, and D. How many models are there for the following sentences?

a.  $B \vee C$ .

Sentence is false only if B and C are false, which occurs in 4 cases for A and D, leaving 12.

b.  $\neg A \vee \neg B \vee \neg C \vee \neg D$ .

Sentence is false only if A, B, C, and D are true, which occurs in 1 case, leaving 15.

c.  $(A \Rightarrow B) \wedge A \wedge \neg B \wedge C \wedge D$ .

The last four conjuncts specify a model in which the first conjunct is false, so 0.

## Exercise 8.4

According to some political pundits, a person who is radical (R) is electable (E) if he/she is conservative (C), but otherwise is not electable.

a. Which of the following are correct representations of this assertion?

(i)  $(R \wedge E) \Leftrightarrow C$

(ii)  $R \Rightarrow (E \Leftrightarrow C)$

(iii)  $R \Rightarrow ((C \Rightarrow E) \vee \neg E)$

## Exercise 8.4

According to some political pundits, a person who is radical (R) is electable (E) if he/she is conservative (C), but otherwise is not electable.

a. Which of the following are correct representations of this assertion?

(i)  $(R \wedge E) \Leftrightarrow C$

No.

This sentence asserts, among other things, that all conservatives are radical, which is not what was stated.

## Exercise 8.4

According to some political pundits, a person who is radical (R) is electable (E) if he/she is conservative (C), but otherwise is not electable.

a. Which of the following are correct representations of this assertion?

$$(ii) R \Rightarrow (E \Leftrightarrow C)$$

Yes.

This says that if a person is a radical then they are electable if and only if they are conservative.

## Exercise 8.4

According to some political pundits, a person who is radical (R) is electable (E) if he/she is conservative (C), but otherwise is not electable.

a. Which of the following are correct representations of this assertion?

$$(iii) R \Rightarrow ((C \Rightarrow E) \vee \neg E)$$

No.

This is equivalent to  $\neg R \vee \neg C \vee E \vee \neg E$  which is a tautology, true under any assignment.



## Exercise 8.5

Show using the propositional resolution calculus that:

$$(1) \quad P \vee \neg Q$$

$$(2) \quad \neg P \vee R \vee S$$

$$(3) \quad Q$$

$$(4) \quad \neg P \vee \neg R$$

$$(5) \quad Q \vee P \vee R$$

$$(6) \quad S \vee \neg Q$$

$$(7) \quad R$$

is not satisfiable;

## Exercise 8.5 - Solution

Show using the propositional resolution calculus that:

(1)  $P \vee \neg Q$

(2)  $\neg P \vee R \vee S$

(3)  $Q$

(4)  $\neg P \vee \neg R$

(5)  $Q \vee P \vee R$

(6)  $S \vee \neg Q$

(7)  $R$

(8)  $P$

## Exercise 8.5 - Solution

Show using the propositional resolution calculus that:

(1)  $P \vee \neg Q$

(2)  $\neg P \vee R \vee S$

(3)  $Q$

(4)  $\neg P \vee \neg R$

(5)  $Q \vee P \vee R$

(6)  $S \vee \neg Q$

(7)  $R$

(8)  $P$

(9)  $\neg R$

## Exercise 8.5 - Solution

Show using the propositional resolution calculus that:

(1)  $P \vee \neg Q$

(2)  $\neg P \vee R \vee S$

(3)  $Q$

(4)  $\neg P \vee \neg R$

(5)  $Q \vee P \vee R$

(6)  $S \vee \neg Q$

(7)  $R$

(8)  $P$

(9)  $\neg R$

(10)  $\{\}$

## Exercise 8.6

Show using the propositional resolution calculus that:

$$(1) \quad P \vee Q$$

$$(2) \quad \neg Q \vee P \vee \neg R$$

$$(3) \quad \neg P \vee \neg R$$

$$(4) \quad R$$

is not satisfiable.

## Exercise 8.6

Show using the propositional resolution calculus that:

(1)  $P \vee Q$

(2)  $\neg Q \vee P \vee \neg R$

(3)  $\neg P \vee \neg R$

(4)  $R$

(5)  $P \vee P \vee \neg R$

## Exercise 8.6

Show using the propositional resolution calculus that:

(1)  $P \vee Q$

(2)  $\neg Q \vee P \vee \neg R$

(3)  $\neg P \vee \neg R$

(4)  $R$

(5)  $P \vee P \vee \neg R$

(6)  $P \vee \neg R$

## Exercise 8.6

Show using the propositional resolution calculus that:

(1)  $P \vee Q$

(2)  $\neg Q \vee P \vee \neg R$

(3)  $\neg P \vee \neg R$

(4)  $R$

(5)  $P \vee P \vee \neg R$

(6)  $P \vee \neg R$

(7)  $\neg R$



## Exercise 8.6

Show using the propositional resolution calculus that:

(1)  $P \vee Q$

(2)  $\neg Q \vee P \vee \neg R$

(3)  $\neg P \vee \neg R$

(4)  $R$

(5)  $P \vee P \vee \neg R$

(6)  $P \vee \neg R$

(7)  $\neg R$

(8)  $\{\}$

## Exercise 8.7

Show using the propositional resolution calculus that:

$$(1) \quad G \vee D \vee \neg B$$

$$(2) \quad \neg E \vee C$$

$$(3) \quad G$$

$$(4) \quad F \vee C \vee \neg A$$

$$(5) \quad \neg A$$

$$(6) \quad \neg C \vee B$$

$$(7) \quad \neg G \vee \neg F \vee E$$

$$(8) \quad F$$

$$(9) \quad \neg F \vee \neg E \vee D$$

$$(10) \quad \neg F \vee \neg B \vee A$$

is not satisfiable;

## Exercise 8.7

(1)  $G \vee D \vee \neg B$

(2)  $\neg E \vee C$

(3)  $G$

(4)  $F \vee C \vee \neg A$

(5)  $\neg A$

(6)  $\neg C \vee B$


(7)  $\neg G \vee \neg F \vee E$

(8)  $F$

(9)  $\neg F \vee \neg E \vee D$

(10)  $\neg F \vee \neg B \vee A$

Resolution: deduction of a new clause from a pair of clauses with EXACTLY ONE incompatible variable (resolvent)



(11)  $[(7), (3)] \Rightarrow (\neg F \vee E)$

(12)  $[(8), (11)] \Rightarrow (E)$

(13)  $[(12), (2)] \Rightarrow (C)$

(14)  $[(13), (6)] \Rightarrow (B)$

(15)  $[(8), (10)] \Rightarrow (\neg B \vee A)$

(16)  $[(14), (15)] \Rightarrow (A)$

(17)  $[(5), (16)] \Rightarrow \{\}$

## Exercise 8.7

(1)  $G \vee D \vee \neg B$

(2)  $\neg E \vee C$

(3)  $G$

(4)  $F \vee C \vee \neg A$

(5)  $\neg A$

(6)  $\neg C \vee B$


(7)  $\neg G \vee \neg F \vee E$

(8)  $F$

(9)  $\neg F \vee \neg E \vee D$

(10)  $\neg F \vee \neg B \vee A$

Resolution: deduction of a new clause from a pair of clauses with EXACTLY ONE incompatible variable (resolvent)



(11)  $[(7), (3)] \Rightarrow (\neg F \vee E)$

(12)  $[(8), (11)] \Rightarrow (E)$

(13)  $[(12), (2)] \Rightarrow (C)$

(14)  $[(13), (6)] \Rightarrow (B)$

(15)  $[(8), (10)] \Rightarrow (\neg B \vee A)$

(16)  $[(14), (15)] \Rightarrow (A)$

(17)  $[(5), (16)] \Rightarrow \{\}$

## Exercise 8.12

Show using the propositional resolution calculus that:

(1)  $\neg C \vee \neg D \vee E$

(2)  $F$

(3)  $\neg D \vee B$

(4)  $\neg B \vee A$

(5)  $F \vee E \vee \neg A$

(6)  $C \vee B \vee \neg G$

(7)  $\neg C \vee \neg A \vee G$

(8)  $\neg F \vee \neg C \vee D$

(9)  $C$

(10)  $\neg G$

is not satisfiable;

## Exercise 8.12

(1)  $\neg C \vee \neg D \vee E$

(2)  $F$

(3)  $\neg D \vee B$

(4)  $\neg B \vee A$

(5)  $F \vee E \vee \neg A$

(6)  $C \vee B \vee \neg G$

(7)  $\neg C \vee \neg A \vee G$

(8)  $\neg F \vee \neg C \vee D$

(9)  $C$

(10)  $\neg G$

(11)  $[(8), (2)] \Rightarrow (\neg C \vee D)$

(12)  $[(9), (11)] \Rightarrow (D)$

(13)  $[(12), (3)] \Rightarrow (B)$

(14)  $[(13), (4)] \Rightarrow (A)$

(15)  $[(9), (7)] \Rightarrow (\neg A \vee G)$

(16)  $[(14), (15)] \Rightarrow (G)$

(17)  $[(10), (16)] \Rightarrow \{\}$

## Exercise 8.8

Using DPLL algorithm to find a model for the following proposition:

- $(A \vee B) \wedge (C \vee D) \wedge \neg B$

## Exercise 8.8

Using DPLL algorithm to find a model for the following proposition:

- $(A \vee B) \wedge (C \vee D) \wedge \neg B$
- $(A \vee \textcolor{red}{B}) \wedge (C \vee D) \wedge \textcolor{green}{\neg B}$

Propagate:  $B = \text{False}$



## Exercise 8.8

Using DPLL algorithm to find a model for the following proposition:

- $(A \vee B) \wedge (C \vee D) \wedge \neg B$
- $(A \vee \textcolor{red}{B}) \wedge (C \vee D) \wedge \textcolor{green}{\neg B}$
- $(\textcolor{green}{A} \vee \textcolor{red}{B}) \wedge (C \vee D) \wedge \textcolor{green}{\neg B}$

Propagate:  $B = \text{False}$

Propagate:  $A = \text{True}$

## Exercise 8.8

Using DPLL algorithm to find a model for the following proposition:

- $(A \vee B) \wedge (C \vee D) \wedge \neg B$
- $(A \vee \textcolor{red}{B}) \wedge (C \vee D) \wedge \neg \textcolor{green}{B}$
- $(\textcolor{green}{A} \vee \textcolor{red}{B}) \wedge (C \vee D) \wedge \neg \textcolor{green}{B}$
- $(\textcolor{green}{A} \vee \textcolor{red}{B}) \wedge (\textcolor{green}{C} \vee D) \wedge \neg \textcolor{green}{B}$

Propagate:  $B = \text{False}$

Propagate:  $A = \text{True}$

Decide:  $C = \text{True}$

Interpretation:  $A=\text{True}; B=\text{False}; C=\text{True}$

## Exercise 8.9

Using DPLL algorithm to find a model for the following proposition:

- $(\neg A \vee B) \wedge (\neg C \vee \neg B) \wedge (C \vee \neg B) \wedge (A \vee D)$

## Exercise 8.9

Using DPLL algorithm to find a model for the following proposition:

- $(\neg A \vee B) \wedge (\neg C \vee \neg B) \wedge (C \vee \neg B) \wedge (A \vee D)$
- $(\neg \textcolor{red}{A} \vee B) \wedge (\neg C \vee \neg B) \wedge (C \vee \neg B) \wedge (\textcolor{green}{A} \vee D)$

Decide:  $A = \text{True}$

## Exercise 8.9

Using DPLL algorithm to find a model for the following proposition:

- $(\neg A \vee B) \wedge (\neg C \vee \neg B) \wedge (C \vee \neg B) \wedge (A \vee D)$
- $(\neg \textcolor{red}{A} \vee \textcolor{green}{B}) \wedge (\neg C \vee \neg \textcolor{red}{B}) \wedge (C \vee \neg \textcolor{red}{B}) \wedge (\textcolor{green}{A} \vee D)$

Decide:  $A = \text{True}$

Propagate:  $B = \text{True}$

## Exercise 8.9

Using DPLL algorithm to find a model for the following proposition:

- $(\neg A \vee B) \wedge (\neg C \vee \neg B) \wedge (C \vee \neg B) \wedge (A \vee D)$
- $(\neg \textcolor{red}{A} \vee \textcolor{green}{B}) \wedge (\neg C \vee \neg \textcolor{red}{B}) \wedge (C \vee \neg \textcolor{red}{B}) \wedge (\textcolor{green}{A} \vee D)$
- $(\neg \textcolor{red}{A} \vee \textcolor{green}{B}) \wedge (\neg \textcolor{green}{C} \vee \neg \textcolor{red}{B}) \wedge (\textcolor{red}{C} \vee \neg \textcolor{red}{B}) \wedge (\textcolor{green}{A} \vee D)$

Decide:  $A = \text{True}$

Propagate:  $B = \text{True}$

Propagate:  $C = \text{False}$

## Exercise 8.9

Using DPLL algorithm to find a model for the following proposition:

- $(\neg A \vee B) \wedge (\neg C \vee \neg B) \wedge (C \vee \neg B) \wedge (A \vee D)$
- $(\neg \textcolor{red}{A} \vee \textcolor{green}{B}) \wedge (\neg C \vee \neg \textcolor{red}{B}) \wedge (C \vee \neg \textcolor{red}{B}) \wedge (\textcolor{green}{A} \vee D)$
- $(\neg \textcolor{red}{A} \vee \textcolor{green}{B}) \wedge (\neg \textcolor{green}{C} \vee \neg \textcolor{red}{B}) \wedge (\textcolor{red}{C} \vee \neg \textcolor{red}{B}) \wedge (\textcolor{green}{A} \vee D)$

Decide:  $A = \text{True}$

Propagate:  $B = \text{True}$

Propagate:  $C = \text{False}$

## Exercise 8.9

Using DPLL algorithm to find a model for the following proposition:

- $(\neg A \vee B) \wedge (\neg C \vee \neg B) \wedge (C \vee \neg B) \wedge (A \vee D)$
- $(\neg \textcolor{green}{A} \vee B) \wedge (\neg C \vee \neg B) \wedge (C \vee \neg B) \wedge (\textcolor{red}{A} \vee D)$

Decide:  $A = \text{False}$



## Exercise 8.9

Using DPLL algorithm to find a model for the following proposition:

- $(\neg A \vee B) \wedge (\neg C \vee \neg B) \wedge (C \vee \neg B) \wedge (A \vee D)$
- $(\neg \textcolor{green}{A} \vee B) \wedge (\neg C \vee \neg B) \wedge (C \vee \neg B) \wedge (\textcolor{red}{A} \vee \textcolor{green}{D})$

Decide:  $A = \text{False}$

Propagate:  $D = \text{True}$

## Exercise 8.9


Using DPLL algorithm to find a model for the following proposition:

- $(\neg A \vee B) \wedge (\neg C \vee \neg B) \wedge (C \vee \neg B) \wedge (A \vee D)$
- $(\neg \textcolor{green}{A} \vee \textcolor{red}{B}) \wedge (\neg C \vee \neg \textcolor{green}{B}) \wedge (C \vee \neg \textcolor{green}{B}) \wedge (\textcolor{red}{A} \vee \textcolor{green}{D})$

Decide:  $A = \text{False}$

Propagate:  $D = \text{True}$

Decide:  $B = \text{False}$



Interpretation:  $A=\text{False}; B=\text{False}; D=\text{True}$

## Exercise 8.10

Consider the following CNF formula in PL:

$$\begin{aligned} & (\neg L) \wedge \\ & (M \vee N \vee L) \wedge \\ & (\neg N \vee I) \wedge \\ & (\neg H \vee \neg N \vee G) \wedge \\ & (\neg H \vee \neg G \vee \neg N) \wedge \\ & (F \vee \neg N \vee H) \wedge \\ & (H \vee \neg F \vee \neg N) \wedge \\ & (N \vee E \vee L) \wedge \\ & (\neg E \vee \neg D \vee N) \wedge \\ & (D \vee C \vee B) \wedge \\ & (\neg C \vee L \vee \neg B) \wedge \\ & (N \vee \neg C \vee B) \wedge \\ & (C \vee A \vee \neg E) \wedge \\ & (D \vee C \vee \neg A) \end{aligned}$$

Draw the search tree obtained by applying to the above formula the DPLL algorithm without the pure-symbol rule. Variables should be chosen according to **reverse alphabetic order**, and assigned **true** first.

## Exercise 8.10

$\neg L$

$M \vee N \vee L$

$\neg N \vee I$

$\neg H \vee \neg N \vee G$

$\neg H \vee \neg G \vee \neg N$

$F \vee \neg N \vee H$

$H \vee \neg F \vee \neg N$

$N \vee E \vee L$

$\neg E \vee \neg D \vee N$

$D \vee C \vee B$

$\neg C \vee L \vee \neg B$


$N \vee \neg C \vee B$

$C \vee A \vee \neg E$

$D \vee C \vee \neg A$

## Exercise 8.10

$(\neg L \vee I) \wedge$   
 $(M \vee N \vee L) \wedge$   
 $(\neg N \vee I) \wedge$   
 $(\neg H \vee \neg N \vee G) \wedge$   
 $(\neg H \vee \neg G \vee \neg N) \wedge$   
 $(F \vee \neg N \vee H) \wedge$   
 $(H \vee \neg F \vee \neg N) \wedge$   
 $(N \vee E \vee L) \wedge$   
 $(\neg E \vee \neg D \vee N) \wedge$   
 $(D \vee C \vee B) \wedge$   
 $(\neg C \vee L \vee \neg B) \wedge$   
 $(N \vee \neg C \vee B) \wedge$   
 $(C \vee A \vee \neg E) \wedge$   
 $(D \vee C \vee \neg A)$



## Exercise 8.13

Consider the following CNF formula in PL:

$$\begin{aligned} & (\neg C \vee B \vee A \vee C) \wedge \\ & (\neg A \vee D \vee E) \wedge \\ & (\neg E \vee \neg A \vee F) \wedge \\ & (\neg E \vee \neg F \vee \neg A) \wedge \\ & (G \vee \neg A \vee E) \wedge \\ & (E \vee \neg G \vee \neg A) \wedge \\ & (A \vee H \vee C) \wedge \\ & (\neg H \vee \neg I \vee A) \wedge \\ & (I \vee L \vee M) \wedge \\ & (\neg L \vee C \vee \neg M) \wedge \\ & (A \vee \neg L \vee M) \wedge \\ & (L \vee N \vee \neg H) \wedge \\ & (I \vee L \vee \neg N) \end{aligned}$$

Draw the search tree obtained by applying to the above formula the DPLL algorithm without the pure-symbol rule. Variables should be chosen according to **alphabetic order**, and assigned **true** first.

## Exercise 8.13

$\neg C$

B V A V C

$\neg A$  V D

$\neg E$  V  $\neg A$  V F

$\neg E$  V  $\neg F$  V  $\neg A$

G V  $\neg A$  V E

E V  $\neg G$  V  $\neg A$

A V H V C

$\neg H$  V  $\neg I$  V A

I V L V M

$\neg L$  V C V  $\neg M$


A V  $\neg L$  V M

L V N V  $\neg H$

I V L V  $\neg N$

# Exercise 8.13

$\neg C$   
 $B \vee A \vee C$   
 $\neg A \vee D$   
 $\neg E \vee \neg A \vee F$   
 $\neg E \vee \neg F \vee \neg A$   
 $G \vee \neg A \vee E$   
 $E \vee \neg G \vee \neg A$   
 $A \vee H \vee C$   
 $\neg H \vee \neg I \vee A$   
 $I \vee L \vee M$   
 $\neg L \vee C \vee \neg M$   
 $A \vee \neg L \vee M$   
 $L \vee N \vee \neg H$   
 $I \vee L \vee \neg N$





## Exercise 8.11 - Homework

Prove each of the following assertions:

- a.  $\alpha$  is valid if and only if  $\text{True} \models \alpha$ . Forward: If  $\alpha$  is valid it is true in all models, hence it is true in all models of  $\text{True}$ . Backward: if  $\text{True} \models \alpha$  then  $\alpha$  must be true in all models of  $\text{True}$ , i.e., in all models, hence  $\alpha$  must be valid.
- b. For any  $\alpha$ ,  $\text{False} \models \alpha$ .  $\text{False}$  doesn't hold in any model, so  $\alpha$  trivially holds in every model of  $\text{False}$ .
- c.  $\alpha \models \beta$  if and only if the sentence  $(\alpha \Rightarrow \beta)$  is valid. Both sides are equivalent to the assertion that there is no model in which  $\alpha$  is true and  $\beta$  is false, i.e., no model in which  $\alpha \Rightarrow \beta$  is false.
- d.  $\alpha \equiv \beta$  if and only if the sentence  $(\alpha \Leftrightarrow \beta)$  is valid. Both sides are equivalent to the assertion that  $\alpha$  and  $\beta$  have the same truth value in every model.
- a.  $\alpha \models \beta$  if and only if the sentence  $(\alpha \wedge \neg\beta)$  is unsatisfiable. As in c, both sides are equivalent to the assertion that there is no model in which  $\alpha$  is true and  $\beta$  is false.