

VNIVERSITAT Đ VALÈNCIA

The discovery of the Higgs boson in 2012 opened a new field for exploration in the realm of particle physics. To better understand the Standard Model of particle physics, it is of prominent interest to study the Yukawa coupling of the Higgs boson to the top quark y_t .

The only way of directly measuring both the magnitude and sign of y_t at LHC is via the production of a Higgs boson with a single top quark. In this work, a search for the tHq production is presented in a final state with two

light-flavoured-charged leptons (electrons or muons) and one hadronically-decaying tau lepton (named $2\ell + \tau_{had}$ channel). This analysis uses an integrated luminosity of 140fb^{-1} of proton-proton collision data at a centre-of-mass energy of 13 TeV collected by the ATLAS detector during LHC Run 2. This search for the tHq production in the $2\ell + \tau_{had}$ channel is an important contribution to the exploration of the Higgs-top quark Yukawa coupling and a step forward in

Search for the Higgs boson produced in association
with a top quark using τ leptons with ATLAS

Pablo Martínez Agulló

Doctoral
Thesis

2024



VNIVERSITAT Đ VALÈNCIA

Search for the Higgs boson produced in association with a top quark using τ leptons with ATLAS

Doctoral thesis
Pablo Martínez Agulló

València, 2024

Facultat de Física
Departament de Física Atòmica, Molecular i Nuclear