

- Q1** During development of embryo sac in angiosperm plant, polar nuclei located -
 (A) In Primary endosperm cell
 (B) In large central cell above the egg apparatus
 (C) Below the egg apparatus
 (D) both A and B

- Q2** Select the mismatched pair.
 (A) Microsporangium- Pollen sac
 (B) Megasporangium - Ovule
 (C) Pollen grain - Integument
 (D) Embryo sac- Female gametophyte

- Q3** Read the following statements and mark the right option
 A. In angiosperm, each ovule always have only two protective layers known as integument
 B. The synergids have special cellular thickenings at the micropylar tip called filiform apparatus
 C. Nucleus of functional megaspore divide mitotically
 (A) Only B is correct
 (B) Only B and C are correct
 (C) Only A and C are correct
 (D) All A, B and C are correct

- Q4** From among the sets of terms given below, identify those that are associated with the gynoecium.
 (A) Stigma, ovule, embryo sac, placenta
 (B) Thalamus, pistil, style, ovule
 (C) Ovule, ovary, embryo sac, tapetum
 (D) Ovule, stamen, ovary, embryo sac


- Q5** Match the following columns.

Column-I		Column-II	
A.	Chalaza	p.	Integument is absent
B.	Nucellus	q.	Junction between ovule and funicle
C.	Micropyle	r.	Basal part of ovule
D.	Hilum	s.	Parenchymatous mass

Choose the correct answer from the options given below.

- (A) A-(r), B-(q), C-(s), D-(p)
 (B) A-(p), B-(r), C-(s), D-(q)
 (C) A-(r), B-(s), C-(p), D-(q)
 (D) A-(q), B-(r), C-(p), D-(s)

- Q6** In the following figure, label the parts A, B, C, D, E



- (A) Hilum, Funicle, inner integument, nucellus, micropyle
 (B) Funicle, micropyle, inner integument, nucellus, chalazal pole
 (C) Hilum, Funicle integument, Embryo sac, chalazal pole
 (D) Funicle, Hilum, integument, Embryo sac, Micropylar pole

- Q7** Which part of ovule stores abundant food reserves?
 (A) Chalaza (B) Integument
 (C) Funicle (D) Nucellus
- Q8** Nucellus is
 (A) Poorly developed in all flowering plants

- (B) Parenchymatous mass of tissue
(C) Basal part of ovule
(D) Located inside the female gametophyte

Q9 An ovule generally has embryo sac formed from a megaspore through division

- (A) Single, equational
(B) Single, reductional
(C) Four, meiotic
(D) Four, mitotic

Q10 A typical angiosperm embryo sac at maturity is:

- (A) 7-nucleate and 8-celled
(B) 7-nucleate and 7-celled
(C) 8-nucleate and 8-celled
(D) 8-nucleate and 7-celled

Q11 Largest cell of the ovule is

- (A) Megaspore mother cell
(B) Antipodal cell
(C) Central cell
(D) None of these

Q12 Match column I with column II:

	Column I		Column II
A.	Egg apparatus	I.	Micropylar end
B.	Chalazal end	II.	Antipodals
C.	Filiform apparatus	III.	Synergids & Egg cell
D.	MMC differentiated	IV.	Synergids

Choose the correct option from the codes given below

- (A) A-III, B-II, C-IV, D-I
(B) A-II, B-III, C-IV, D-I
(C) A-I, B-III, C-II, D-IV
(D) A-IV, B-I, C-II, D-III

Q13 Which of the following is a direct structure formed by a functional megaspore?

- (A) Ovule (B) Endosperm
(C) Embryo (D) Embryo sac

Q14 Monosporic development in angiosperms implies

- (A) Meiosis in megaspore mother cell
(B) Out of four microspores only one microspore is functional

- (C) Out of four megaspores only one megaspore is functional and undergo meiosis
(D) Out of four megaspores only one megaspore is functional and undergo mitosis.


Q15 Identify the incorrect statement related to Pollination:

- (A) Moths and butterflies are the most dominant pollinating agents among insects
(B) Pollination by water is quite rare in flowering plants
(C) Pollination by wind is more common amongst abiotic pollination
(D) Flowers produce foul odours to attract flies and beetles to get pollinated

Q16 The number of cells present in the egg apparatus in angiosperms is

- (A) Two (B) Five
(C) Three (D) Four

Q17 Given below is the diagrammatic view of an embryo sac. Identify the labelled parts A, B, C, D, and select the right option.




	Part-A	Part-B	Part-C	Part-D
(1)	Synergid	Central cell	Filiform apparatus	Egg nucleus
(2)	Central cell	Synergid	Egg nucleus	Filiform apparatus
(3)	Central cell	Egg nucleus	Synergid	Filiform apparatus
(4)	PEC	Egg nucleus	Synergid	Pollen tube

- (A) (1) (B) (2)
(C) (3) (D) (4)

Q18



In the given diagram, label the parts A, B, C, D, E respectively.



(A) Synergids, Central cell, Egg cell, Antipodals, Polar nuclei

(B)

Antipodals, Polar nuclei, Egg cell, Synergids, Filiform apparatus

(C) Polar nuclei, Antipodals, Egg cell, Synergids, Filiform apparatus

(D) Antipodals, Central cell, Egg cell, Synergids, Filiform apparatus

Q19 Functional megaspore in an angiosperm develops into:

(A) Ovule

(B) Endosperm

(C) Embryo sac

(D) Embryo



Answer Key

Q1 (C)

Q2 (C)

Q3 (B)

Q4 (A)

Q5 (C)

Q6 (B)

Q7 (D)

Q8 (B)

Q9 (B)

Q10 (D)

Q11 (C)

Q12 (A)

Q13 (D)

Q14 (D)

Q15 (A)

Q16 (C)

Q17 (C)

Q18 (B)

Q19 (C)



[Master NCERT with PW Books APP](#)