

Q1 Which of the following is **not** paired structure?

- (A) Seminal vesicle
- (B) Ejaculatory duct
- (C) Bulbourethral gland
- (D) Prostate Gland

Q2 Ejaculatory duct opens into:

- (A) ureter.
- (B) urethra.
- (C) urinary bladder.
- (D) testis.

Q3 Match the **List-I** with **List-II**:

List-I		List-II	
A.	Leydig cells	P.	Testosterone
B.	Prostate gland	Q.	Acidic fluid
C.	Bulbourethral glands	R.	Mucus
D.	Seminal vesicles	S.	Fructose, prostaglandins

Choose the **correct** answer from the options given below:

- (A) A-(P); B-(Q); C-(R); D-(S)
- (B) A-(S); B-(Q); C-(R); D-(P)
- (C) A-(R); B-(P); C-(Q); D-(S)
- (D) A-(Q); B-(S); C-(P); D-(R)

Q4 Human male urethra acts as passage for:

- (A) Urine
- (B) Sperms only
- (C) Semen
- (D) Both (A) & (C)

Q5 Which of the following depicts the **correct** pathway of transport of sperms?

- (A) Rete testis → Epididymis → Efferent ductules → Vas deferens
- (B) Rete testis → Vas deferens → Efferent ductules → Epididymis
- (C)

Efferent ductules → Rete testis → Vas deferens → Epididymis

- (D) Rete testis → Efferent ductules → Epididymis → Vas deferens

Q6 Testes descend into scrotum in mammals for

- (A) Spermatogenesis
- (B) Fertilisation
- (C) Development of sex organs
- (D) Development of visceral organs

Q7 Choose the incorrect statement w.r.t. primary sex organ in males.

- (A) They have immunocompetent cells in their interstitium.
- (B) They bear connection with abdomen.
- (C) They are lined by meiotically dividing sertoli cells and germ cells.
- (D) They have leydig cells that secrete androgens.

Q8 Read the following statements and find out the **incorrect** statements.

1. Each testicular lobule contains one to three highly coiled seminiferous tubules in which sperm are produced.
2. Each seminiferous tubule is lined on its inside by two types of cells called Leydig cells and Sertoli cells.
3. The region outside the seminiferous tubules called interstitial space, contains small blood vessels and male germ cells (spermatogonia) which lead to sperm formation.
4. In testis immunologically competent cells are also present.
5. The seminiferous tubules of the testis open into the rete testis through vasa efferentia.



(A) 2 and 3

(B) 2 and 4

(C) 4 and 5

(D) 2, 3 and 5



Answer Key

Q1 (D)

Q2 (B)

Q3 (A)

Q4 (D)

Q5 (D)

Q6 (A)

Q7 (C)

Q8 (D)



[Master NCERT with PW Books APP](#)

