



NAMIBIA UNIVERSITY
OF SCIENCE AND TECHNOLOGY

**PAPER REVIEW OF Game Theory, Evolutionary Stable Strategies and
the Evolution of Biological Interactions Cowden (2012)**

by

MARIA SHIKWAMBI

222064374

BACHELOR OF ARTIFICIAL INTELLIGENCE HONOURS

FACULTY OF COMPUTING AND INFORMATICS

NAMIBIA UNIVERSITY OF SCIENCE AND TECHNOLOGY

COMPUTATIONAL GAME THEORY

1. Abstract

Evolutionary game theory is presented in Cowden's 2012 educational essay as a framework for comprehending the variety of biological relationships, from cooperation and altruism to competitiveness. The study describes how the Hawk-Dove game, the Iterated Prisoner's Dilemma, and the Evolutionarily Stable Strategy (ESS) idea can be used to forecast which behaviours natural selection would favour. The paper lacks mathematical rigour, ignores spatial and stochastic dynamics, and ignores current discussions in the subject, yet being effective as an approachable synthesis for students. After analysing the paper's limits, contributions, and related work, this review comes to the conclusion that while it is a useful starting point, further research is necessary.

2. Introduction

A key issue in evolutionary biology is: if natural selection promotes "survival of the fittest," why do mutualism, cooperation, and altruism continue to exist? In order to address this, Cowden presents evolutionary game theory, a mathematical framework in which payoffs stand in for fitness and strategies are inherited phenotypes. The paper, which was published in Nature Education Knowledge, compresses decades of theoretical research into a narrative that is understandable to biology teachers and students.

3. Summary

3.1 Problem Being Addressed

"If natural selection is survival of the fittest, why isn't everything a competition?" is the main question posed at the beginning of the paper. The goal of evolutionary ecologists is to comprehend the complex behavioural interactions that occur between species as they compete for resources. In addition to instances of exploitation and altruism, these relationships might be competitive or cooperative. Explaining how cooperation and altruism can withstand when they appear to be at odds with the individualistic, survival-of-the-fittest process of evolution is difficult.

3.2 Main Contribution

Evolutionary game theory is presented in this text as a mathematical framework for comprehending biological interactions. Within this framework:

Strategies are hereditary traits, such as cooperative or aggressive behaviours.
Darwinian fitness (reproductive success) is represented by payoffs.

The strategies taken by other members of the population determine success.
The main idea is the Evolutionarily Stable Strategy (ESS), which is a strategy that cannot be overrun or replaced by any other strategy through natural selection if it is embraced by every member of a population. ESSs may occur for collaboration, competitiveness, and even altruism, as the paper illustrates.

3.3 Theoretical Results

3.3.1 Hawk-Dove Game Competitive Interactions:

Hawk is the ESS when the advantage of the resource (B) outweighs the expense of fighting (C).

A hybrid strategy develops when $B < C$ because neither pure Hawk nor pure Dove is stable.

This explains why behavioural polymorphism and ritualised fighting are prevalent in nature.

3.3.2 Collaborative Exchanges (Iterated Prisoner's Dilemma):

Defection is the sole ESS in one-shot interactions.

The Tit-for-Tat approach turns cooperation into an ESS when interactions are repeated across evolutionary time.

When the probability of future meetings is high enough, cooperation develops.

3.3.3 Complex interactions

Mixed strategies that depend on ownership status, resource quality, or quantity evolve. Conventional tactics can resolve asymmetric competitions (owner vs. intruder) without costly conflict.

Reward and punishment modifiers that produce positive feedback loops can stabilise mutualisms between animals.

3.3.4 Altruism

Reciprocal altruism in repeated interactions can lead to the evolution of altruism, which is the reduction of one's own immediate fitness for the benefit of another.

Altruism is sustained by both direct reciprocity (repeated contacts between the same couple) and indirect reciprocity (reputation-based, with various partners).

Through kin selection, inclusive fitness (Hamilton's Rule) offers a different explanation for altruism.

4. The related work

Stable Evolutionary Strategies. Maynard Smith's (1979) formalisation of the Evolutionarily Stable Strategy (ESS) serves as the foundation for Cowden's paper. By replacing natural selection based on inherited strategies for rational decision-makers, Maynard Smith showed how game theory may be applied to evolutionary biology. An ESS is a strategy that is resistive to uncommon competitors when it is prevalent in a population. This idea gives Cowden the rigorous standard by which she determines which behaviours will be favoured by natural selection.

Collaboration through reciprocity. The Iterated Prisoner's Dilemma and Tit-for-Tat mechanisms for collaboration that Cowden proposes are derived from the key research by Axelrod and Hamilton (1981). They demonstrated using computer tournaments that even in situations where defection would predominate in one-shot contacts, straightforward reciprocal techniques maintain cooperation in repeated interactions. In order to explain how cooperation and altruism can be evolutionarily stable, Cowden combined Trivers' (1971) three requirements for reciprocal altruism repeated contacts, benefits exceeding costs, and observable cheating with Axelrod and Hamilton's formal framework

Dynamics in mathematics. Replicator dynamics were formalised by Taylor and Jonker (1978), who linked ESS to population change over time. These equations support the validity of ESS predicts in changing populations, which makes Cowden's omission of them notable.

Cooperation and Spatial Structure. Spatial organization significantly modifies evolutionary outcomes, as shown by Nowak and May (1992). Individuals only interact with their neighbours in their spatial Prisoner's Dilemma, creating dynamic clusters where cooperators and defectors coexist. This demonstrated that, without kin selection or reciprocity extensions, Cowden ignores it but population structure alone can maintain cooperation.

5. The limitations of the paper

Readers have to refer to technical sources because formal explanations and replicator equations are avoided.

Genetic drift and finite population effects are not discussed.

This paper ignores group selection because of concentrating on individual strategies.

There is Little interaction with the literature on experimental evolution.

It predates advances in microbiome applications and network game theory and disregards continuing discussions (such as the inclusive fitness controversy) and presents the framework as settled.

6. Conclusion

Evolutionary game theory is effectively introduced by Cowden (2012) as a unifying framework for biological interactions. Although its teaching goal requires simplification, its basic understanding and approachable presentation make it beneficial for students. It should be accompanied by advanced texts for readers who are interested in the latest developments or mathematical rigour. The paper is a crucial first step in the development of contemporary evolutionary game theory.

References

1. Cowden, C. (2012). Game theory, evolutionary stable strategies and the evolution of biological interactions. *Nature Education Knowledge*, 3(6).
2. Maynard Smith, J. (1979). Game theory and the evolution of behavior. *Proceedings of the Royal Society of London. Series B. Biological Sciences*, 205(1161), 475-488.
3. Axelrod, R., & Hamilton, W. D. (1981). The evolution of cooperation. *Science*, 211(4489), 1390-1396.
4. Taylor, P. D., & Jonker, L. B. (1978). Evolutionary stable strategies and game dynamics. *Mathematical Biosciences*, 40(1-2), 145-156.
5. Nowak, M. A., & May, R. M. (1992). Evolutionary games and spatial chaos. *Nature*, 359(6398), 826-829.
6. Trivers, R. L. (1971). The evolution of reciprocal altruism. *The Quarterly Review of Biology*, 46(1), 35-57.