1. The probability of a leap year selected at random contain 53 Sunday is:					
(a) 53/366 (b) 1/7 (c) 2/7 (d) 53/365					
2. A bag contains 3 red and 2 blue marbles. A marble is drawn at					
random. The probability of drawing a black ball is :					
(a) 3/5 (b) 2/5 (c) 0/5 (d) 1/5					
3. The probability that it will rain tomorrow is 0.85. What is the					
probability that it will not rain tomorrow					
(a) 0.25 (b) 0.145 (c) 3/20 (d) none of these					
4. What is the probability that a number selected from the numbers					
(1, 2, 3,, 15) is a multiple of 4?					
(a) 1/5 (b) 4/5 (c) 2/15 (d) 1/3					
5. What are the total outcomes when we throw three coins?					
(a) 4 (b) 5 (c) 8 (d) 7					
6. The probability that a prime number selected at random from the					
numbers (1,2,3,35) is :					
(a) 12/35 (b) 11/35 (c) 13/35 (d) none of these					
7. The sum of the probability of an event and non event is :					
(a) 2 (b) 1 (c) 0 (d) none of these.					
8. The following probabilities are given; choose the correct answer					
for that which is not possible.					
(a) 0.15 (b) 2/7 (c) 7/5 (d) none of these.					
9. If three coins are tossed simultaneously, than the probability of					
getting at least two heads, is:					
(a) $1/4$ (b) $3/8$ (c) $\frac{1}{2}$ (d) $1/8$					
10. A letter is chosen at random from the letters of the word					
♦ ASSASSINATION ♦ . The probability that the letter chosen has:					
(a) 6/13 (b) 7/13 (c) 1 (d) none of these.					
(a) 6/10 (b) 1/10 (c) 1 (d) Hone of these.					
11. A dice is thrown. Find the probability of getting an even number.					
(A) 2/3 (B) 1 (C) 5/6 (D) 1/2					
(1) 2/3 (3) 1/2					
12. Two coins are thrown at the same time. Find the probability of getting both heads.					
(A) 3/4 (B) 1/4 (C) 1/2 (D) 0					
13. Two dice are thrown simultaneously. The probability of getting a					

sum of 9 is:

(A) 1/10	(B) 3/10	(C) 1/9	(D) 4/	/9	
14. 100 cards are numbered from 1 to 100. Find the probability of getting a prime number.					
(A) 3/4		(C) 1/4	(D)	29/100	
•	a blue ball is do n a bag is:			f the probability en the number of	
16. A box of 600 bulbs contains 12 defective bulbs. One bulb is taken out at random from this box. Then the probability that it is non-defective bulb is: (A) 143/150 (B) 147/150 (C) 1/25 (D) 1/50					
mixed thoro	narked with nun oughly. One card lity that the nun (B) 1/10	d is drawn fro nber on card i	m this box r s a perfect s	andomly, then square.	
18. What is (A) 1/7	the probability (B) 53/366	•	•		
 19. A card is drawn from a well shuffled deck of 52 cards. Find the probability of getting a king of red suit. (A) 1/26 (B) 3/26 (C) 7/52 (D) 1/13 					
20. A game of chance consists of spinning an arrow which is equally likely to come to rest pointing to one of the number 1,2,312 ,then the probability that it will point to an odd number is: (A) 1/6 (B) 1/12 (C) 7/12 (D) 5/12 1/2					
21. A game consists of tossing a one rupee coin 3 times and noting its outcome each time. Aryan wins if all the tosses give the same result i.e. three heads or three tails and loses otherwise. Then the probability that Aryan will lose the game. (A) 3/4 (B) 1/2 (C) 1 (D) 1/4					

•	ajal are friends. is the same birt	•	both will hav	e the
-	(B) 31/365	•	(D) 1	/133225
2. Then the pro	x is chosen at randomates x is chosen at random x is chosen x is	< 2 is?	numbers -2,	-1, 0 , 1,
a marble is dra red is 2/3, ther	ins 24 marbles. awn at random from the number of (C) 8 (D) 7	om the jar, the white marbles i	probability th	
Then the proba	is selected at rai ability that it is a 4/25 (C) 1/25	multiple of 3 a		umbers.
with n dots sho showing 4 dots	_	ortional to n. Tl	ne probability	
a) $\frac{1}{7}$	b) $\frac{5}{42}$	c) $\frac{1}{21}$	d) $\frac{4}{21}$	
	ed by batsman i	_	tches are 50,	70, 82,
a) 25.79	e standard devia b) 25.49		d) 25.69	9
consecutive da	an and mode of t ays 15, 11, 9, 5,	18, 4, 18, 13, 17		The median is the middle value when a data is ordered Mode is the number that occurs most often
a) 13, 15	b) 13, 18	C) 18, 15		d) 13, 16
29. A coin is to 3 cases is	ossed up 4 time	s. The probabili	ty that tails t	u <mark>rn up in</mark> THTT , HTTT , TTHT , TTTH
a) $\frac{1}{2}$	b) $\frac{1}{3}$	c) $\frac{1}{4}$		d) $\frac{1}{6}$
	te between 0 an	d 3. The value (c) 27	of E(X²) is d) 9	·
	n variables X and .et Z= 5X-2Y. Th			.5
2			25 Va	ar(x) - 4Var(y)

25 * 0.2 + 4 * 0.5

3

a)	3

32.Out of the following values, which one is not possible in probability?

a)
$$P(x) = 1$$

b)
$$\sum x P(x) = 3$$

c)
$$P(x) = 0.5$$

d)
$$P(x) = -0.5$$

33.If E(x) = 2 and E(z) = 4, then E(z - x) = ?

d) Insufficient data

34. The covariance of two independent random variable is

$$c) - 1$$

d) Undefined

35.If $\Sigma P(x) = k^2 - 8$ then, the value of k is?

d) Insufficient data

36.If P(x) = 0.5 and x = 4, then E(x) = ?

d) 2

37.In a discrete probability distribution, the sum of all probabilities is always?

d) Undefined

38.If the probability of hitting the target is 0.4, find mean and variance.

$$M = 0.4$$

V = 0.4(1-0.4)

d) 0.6, 0.16

39. If the probability that a bomb dropped from a place will strike the target is 60% and if 10 bombs are dropped, find mean and variance?

M = 10 * 0.6 V = 10 * 0.6 * (1-0.6)

8 * 1/2 = 4

40. Find the mean of tossing 8 coins.

41. What is the mean and variance for standard normal distribution?

		(3.1) b) Mean is 1 and varion (3.5) d) Mean is (3.5) and	
42.Variance a) E(X)		ariable X is given by c) E(X2) - (E(X))2	
43.Mean of a) E(X)		ble X is given by c) E(X2) - (E(X))2	 d) (E(X))2
44.Mean of a) 0	a constant 'a' is b) a	c) a/2	d) 1
45. Variance	of a constant 'a	a' is	

46. Find the mean and variance of X?

b) a

×	0	1	2	3	4	Mean = 0*1/9 + 4*1/9 = 2
f(x)	1/9	2/9	3/9	2/9	1/9	variance = [(0)/ [(2)^2 *3/9] + [(= 16/3

c) a/2

Mean = 0*1/9 + 1*2/9 + 2*3/9 + 3*2/9 + 4*1/9 = 2

variance = $[(0)^2 *1/9] + [(1)^2 *2/9] + [(2)^2 *3/9] + [(3)^2 *2/9] + [(4)^2 *1/9] = 16/3$

variance = $16/3 - (2)^2 = 4/3$

a) 2, 4/3

a) 0


b) 3, 4/3

c) 2, 2/3

d) 3, 2/3

d) 1

47. Find the expectation of a random variable X?



48. In a Binomial Distribution, if p, q and n are probability of success, failure and number of trials respectively then variance is given by

b) npq

c) np2q

d) npq2

- 49. If 'X' is a random variable, taking values 'x', probability of success and failure being 'p' and 'q' respectively and 'n' trials being conducted, then what is the probability that 'X' takes values 'x'? Use **Binomial Distribution.**
- a) P(X = x) = nCx px qx
- b) P(X = x) = nCx px q(n-x)
- c) P(X = x) = xCn qx p(n-x)
- d) P(x = x) = xCn pn qx
- 50. If 'p', 'q' and 'n' are probability pf success, failure and number of trials respectively in a Binomial Distribution, what is its Standard **Deviation?**

- a) \sqrt{np} b) \sqrt{pq} c) (np)2
- d) \sqrt{npq}