

2026/2/25 RFCMP

moonshine現象の紹介

(Introduction to moonshine phenomena)



RFCMP 2026 Mini Workshop

moonshine 現象の紹介 岡田 昌樹

1. Preliminaries from group theory

1.1. Finite simple groups

1.2. Group extension

2. Monstrous moonshine

3. Conway (super)moonshine

3.1. Duncan's module

3.2. Connection to the SST conjecture

4. Mathieu moonshine

<https://masakiokada0101.github.io/talks.html> 補足資料

1. Preliminaries from group theory


1.1. Finite simple groups Jordan-Hölder theorem

no normal subgrps. other than G and $\{1\}$

Thm (Classification of fin. simple grps.)

A fin. simple grp. is one of the followings.

- cyclic grp. of prime order $\mathbb{Z}_p$ (p : prime)
- alternating grp. A_n ($n \geq 5$)
- simple grps. of Lie type
- 26 sporadic grps.



6 pariahs (not subgrp. quotient of subgrp. of M)

○ ○
○ ○
○ ○

Three important fin. simple grps.:

	order
the Mathieu grp. $M_{24} = \text{Aut}(G_{24}) = ?$	2.45×10^8
the Conway grp. $Co_1 = \text{Aut}(L_{24})/\mathbb{Z}_2 = \text{Aut}_{r=1}(V^{\natural})$	4.16×10^{18}
the monster grp. $M = \text{Aut}(B^{\natural}) = \text{Aut}(V^{\natural})$ $\parallel$ Griess alg. $V^{\natural}/wt=2$	8.08×10^{53}

• Conway grps.

Def

A lattice $(L, \langle -, - \rangle)$ of $\overset{L}{\text{dim. rk.}} n$ is a free ab. grp. $L \subset \mathbb{R}^n$ of rk. n whose basis spans $\mathbb{R}^n$ with a non-dgn. sym. bilin form $\langle -, - \rangle : L \times L \rightarrow \mathbb{R}$.

A 24-dim. pos-def. even self-dual lattice is called a Niemeier lattice.

The Leech lattice Λ_{24} is the Niemeier lattice without vectors with $\|k\|^2 = 2$.
(unique up to isom.)

$$C_{0,0} := \text{Aut}(\Lambda_{24})$$

$$\text{Center}(C_{0,0}) = \{\pm 1\} \cong \mathbb{Z}_2$$

$$C_{0,1} := C_{0,0} / \mathbb{Z}_2$$

simple

• master grp. M

Fact $M = \langle 2^{\overset{\sim}{1+24}} C_{0,1}, \sigma \rangle, \quad \sigma^2 = 1.$
group extension

Def

G is an extraspecial p -group p^{1+n} (p : prime)

$$\Leftrightarrow \begin{cases} \cdot \text{Center}(G) \cong \mathbb{Z}_p \\ \cdot G / \text{Center}(G) \cong (\mathbb{Z}_p)^n \end{cases}$$

ex. gamma matrix algebra

$$\langle \underbrace{\gamma_1, \dots, \gamma_{2m}}_V \mid \{\gamma_i, \gamma_j\} = -2\delta_{ij} \rangle$$

$$\{\pm 1, \pm \gamma_i, \pm \gamma_i \gamma_j, \dots, \pm \gamma_1 \dots \gamma_{2m}\} \cong 2^{1+2m}$$

$$\text{center} = \{\pm 1\}$$

If the number of γ 's is odd $\langle \gamma_1, \dots, \gamma_{2m+1} \rangle$, then $\pm \gamma_1 \dots \gamma_{2m+1} \in \text{center}.$

1.2. Group extension

how to combine two groups N and G .
generalization of $N \times G$, $N \rtimes G$.

Def

G, N : grps.

An extension of G by N is a short exact seq. of grp. homs.

$$1 \rightarrow N \hookrightarrow \hat{G} \xrightarrow[\text{section } S]{\pi} G \rightarrow 1.$$

($\hat{G}$ is also called an extension. Any extension is denoted by $N.G$.)

• $N \triangleleft \hat{G}$

• $\hat{G}/N \cong G$

$\pi \circ s = \text{id}_G$ not necessarily a grp. hom.

Take a section $S : G \rightarrow \hat{G}$. $N \times G \rightarrow \hat{G}$ is a bijection.

$$\begin{matrix} \downarrow & & \downarrow \\ (n, g) & \mapsto & n \cdot s(g) \end{matrix}$$

• If we can take a grp-hom. section S (If the short ex. seq. splits), then the multiplication law of $\hat{G}$ is $\pi(s(g)n's(g')^{-1}) = g \cdot 1 \cdot g'^{-1} = 1 \therefore s(g)n's(g')^{-1} \in N$

$$\begin{aligned} (n \cdot s(g)) \cdot (n' \cdot s(g')) &= (n \cdot s(g)n's(g')^{-1}) \cdot s(gg') \\ &=: (n \cdot \varphi_g(n')) \cdot s(gg') \end{aligned}$$

where $\varphi_g : N \rightarrow N$ defines a G -action $G \curvearrowright N$.

$\therefore \hat{G} \cong N \rtimes_{\varphi} G$. $N:G$

In particular, if φ is a trivial action $\varphi_g = \text{id}_N$, then $\hat{G} \cong N \times G$.

• Suppose S is not a grp. hom.

We can define $\Sigma: G \times G \rightarrow N$ s.t.

$$S(g) S(g') = \Sigma(g, g') S(gg') \quad \pi(\underbrace{S(g) S(g') S(gg')^{-1}}_N) = 1$$

Let N be abelian. Then $\varphi_g: N \rightarrow N$ defines a G -action $G \curvearrowright N$.

$$\begin{aligned} \varphi_g: N &\rightarrow N \\ n &\mapsto S(g)nS(g)^{-1} \end{aligned} \quad \begin{aligned} &S(g) S(g') n S(g')^{-1} S(g)^{-1} \\ &= S(gg') n S(gg')^{-1} \\ &\uparrow \\ &S(gg')^{-1} S(g') S(g) \in N : \text{abelian} \end{aligned}$$

The multi. law of $\hat{G}$ is

$$(n \cdot S(g)) \cdot (n' \cdot S(g')) = (n \cdot \varphi_g(n') \cdot \Sigma(g, g')) \cdot S(gg')$$

Σ is a 2-cocycle:

$$\Sigma(g, g') \cdot \Sigma(gg', g'') = \varphi_g(\Sigma(g', g'')) \Sigma(g, g'g'') \quad \leftarrow \text{associativity of } \hat{G}$$

Prop

G : a grp.

N : a G -module (an ab. grp. with G -action)

$$\left\{ \begin{array}{l} \text{extensions of } G \text{ by } N \\ \text{compatible with the } G\text{-action} \end{array} \right\} / \text{equiv.} \xrightarrow{1:1} H^2(G; N)$$

$$\begin{aligned} &\left(\begin{array}{l} \text{A set } N \times G \text{ with multi. law} \\ (n, g) \cdot (n', g') = (n \cdot \varphi_g(n') \cdot \Sigma(g, g'), gg') \end{array} \right) \longleftarrow [\Sigma] \\ &N \times G \longleftarrow 0 \end{aligned}$$

* Even if $\hat{G} \cong \hat{G}'$, they can be inequiv. as grp. extensions.

$$1 \rightarrow N \rightarrow \hat{G} \rightarrow G \rightarrow 1$$

In particular, if $N \subset \text{Center}(\hat{G})$, then φ is a trivial action.

$$(n \cdot S(g)) \cdot (n' \cdot S(g')) = (n \cdot n' \cdot \Sigma(g, g')) \cdot S(gg')$$

$\hat{G}$ is called a central extension.

Examples

· Fact $H^2(C_0, \overset{\text{trivial action}}{\mathbb{Z}_2}) \cong \mathbb{Z}_2$

$\leadsto$ There are only two central extensions: $\overset{\text{split}}{\mathbb{Z}_2 \times C_0}$ and $\overset{\text{non-split}}{C_{00}}$

· carry (繰り上がり) central extension, non-split

$$\underset{\substack{\uparrow \\ 0-99}}{\mathbb{Z}_{100}} \cong \underset{\substack{\uparrow \\ +99\text{位}}}{\mathbb{Z}_{10}} \cdot \underset{\substack{\uparrow \\ -99\text{位}}}{\mathbb{Z}_{10}}$$

$$27+38=65$$

$$(2, 7) + (3, 8) = (2+3+\underbrace{2(7 \cdot 8)}_{=1 \text{ (carry)}}, 7+8) = (6, 5)$$

2. Monstrous moonshine

modular j -function · modular inv.
(a mod. func. of weight 0)

$$j_1(\tau) = q^{-1} + 196884q + 21493760q^2 + \dots \quad (q = e^{2\pi i \tau})$$

$$j(\tau) = q^{-1} + 744 + \dots$$

$$\underbrace{1 + 196883}_{1 + 196883} \quad \underbrace{1 + 196883 + 21296876}_{1 + 196883 + 21296876}$$

McKay (1978), Thompson (1979)

dim. of irrep. of M

$$d_1 = 1, d_2 = 196883, d_3 = 21296876, \dots \quad \text{"monstrous moonshine"} \quad \text{Conway Norton}$$

Moreover, if we decompose $j_1(\tau)$ into the irr. characters

$$ch_h(\tau) := \text{Tr}_{V_h} q^{L_0 - \frac{c}{24}} \quad (V_h: \text{the irr. highest-weight rep. of weight } h)$$

of the Virasoro alg. of $c=24$,

$$j_1(\tau) = d_1 ch_0 + d_2 ch_2 + d_3 ch_3 + d_4 ch_4$$

$$+ d_6 ch_5 + (d_5 + d_7) ch_6 + \dots$$

very simple!

This observation was explained by a CFT_{VOA} with M symmetry.

idea A chiral CFT_{VOA} V is a rep. sp. of the Vir. alg.:


$$V = V_{h_1} \oplus \dots \oplus V_{h_1} \oplus V_{h_2} \oplus \dots \oplus V_{h_2} \oplus \dots$$

If V has the symmetry M ($\exists$ action $M \curvearrowright V$ which commutes with the Vir. action)
($\text{Aut}(V) \supset M$),

$$V = W_1 \otimes V_{h_1} \oplus W_2 \otimes V_{h_2} \oplus \dots$$

Its character is ← rep. sp. of M

$$\text{Tr}_V q^{L_0 - \frac{c}{24}} = \sum_i (\dim W_i) ch_{h_i}(\tau)$$



Many other relations

fin. grp.

mod. func. /
weak Jacobi form

are observed.

• lattice VOA

pos-def. even self-dual lattice L of rk. n

$\leadsto$ bosonic mod-inv. VOA V_L of $C = n$

As a vec. sp., $V_L = \text{Span}_{\mathbb{C}} \{ \underbrace{\alpha_{-m_1}^{i_1} \cdots \alpha_{-m_\ell}^{i_\ell}}_{\substack{\text{weight} \\ (L_0\text{-eigenvalue})}} |k\rangle \mid \substack{1 \leq i_j \leq n, \ell = 0, 1, \dots, \\ m_j \in \mathbb{Z}_{>0}, k \in L} \}$

creation operator $[\alpha_m^i, \alpha_n^j] = m \delta_{ij} \delta_{m+n,0}$

$m_1 + \cdots + m_\ell + \frac{1}{2}|k|^2$

Its character is

$$\begin{aligned} \text{Tr}_{V_L} q^{L_0 - \frac{c}{24}} &= \frac{\Theta_L(\tau)}{\eta(\tau)^n} \\ &\stackrel{n=24}{=} j_1(\tau) + 24 + \# \{ k \in L \mid |k|^2 = 2 \} \end{aligned}$$

~~$\alpha_{-1,1}^{i_1} |k\rangle$~~

Orbifold by $\mathbb{Z}_2$ ($X^i \mapsto -X^i$)

$L = \Lambda_{24}$

$L_0 - \frac{c}{24} = 0$

So, we consider $V^h := V_{\Lambda_{24}} / \mathbb{Z}_2$.

monster VOA (moonshine module)

How $\text{Aut}(V^h) = M$?

$$\Lambda_{24}$$

cocycle factor $\left\{ \begin{array}{l} \varepsilon: L \times L \rightarrow \mathbb{Z}_2 \end{array} \right\}$

central extension $1 \rightarrow \mathbb{Z}_2 \rightarrow \hat{\Lambda}_{24} \rightarrow \Lambda_{24} \rightarrow 0$

add cr.-an. op. $\downarrow$

$$V_{\Lambda_{24}}$$

$\mathbb{Z}_2$ -orbifold $\left\{ \begin{array}{l} \text{add } \mathbb{Z}_2\text{-twisted sector } \downarrow \\ V_{\Lambda_{24}} \oplus (V_{\Lambda_{24}})_{\text{tw}} \\ \text{project onto } \mathbb{Z}_2\text{-inv. states } \downarrow \end{array} \right.$

$$V^h$$

$$\begin{array}{ccc} (V_L)^{\mathbb{Z}_2\text{-inv.}} \oplus (V_L)^{\mathbb{Z}_2\text{-inv.}}_{\text{tw}} & & \\ \parallel & & \parallel \\ \textcircled{C} V_0 & & T_0 \textcircled{C} \\ \oplus & & \oplus \\ V_3 & \longleftrightarrow & T_3 \\ & \sigma & \text{triality operator} \end{array}$$

$$\text{Aut}(\Lambda_{24}) = C_0$$

$\text{Hom}(L, \mathbb{Z}_2)$
S11
 $(\mathbb{Z}_2)^{24}$

$$\text{Aut}(\hat{\Lambda}_{24}) = 2^{24} \cdot C_0$$

$\uparrow$
non-split

$$\text{Aut}(V_{\Lambda_{24}}) = 2^{24} \cdot C_0$$

$$\hookrightarrow 2^{1+24} \cdot C_0$$

$$\hookrightarrow 2^{1+24} \cdot C_0$$

$\uparrow$
This action does not mix
the untw. sector and the tw. sector.

Thm (Frenkel, Lepowsky, Meurman) "Vertex operator algebras and the Moonshine" (1988)

$$\text{Aut}(V^h) = \langle 2^{1+24} \cdot C_0, \sigma \rangle \cong M.$$

3. Conway (super)moonshine

3.1. Duncan's module

We will construct an $N=1$ VOSA V^{fg} of $C=12$ and $V^{\text{fg}}|_{\text{weight } \frac{1}{2}} = 0$

Thm (Duncan, 2007)

"Conway supermoonshine module"
"Duncan's module"

$$\text{Aut}_{N=1}(V^{\text{fg}}) \cong C_0,$$

Thm (Duncan, Mack-Crane, 2014)

The uniqueness of V^{fg} is still a conj.
($C=24, V^{\text{fg}}|_{\frac{1}{2}}=0$) (FLM uniqueness conj.)

Such a "VOSA" of $C=12$ without weight- $\frac{1}{2}$ states is unique

a self-dual C_2 -cofinite
rational VOSA of CFT type

up to isom.

• n real chiral free fermions

NS sector (Clifford module VOSA)

As a vec. sp.,

$$F_{NS} = \text{Span}_{\mathbb{C}} \left\{ \underbrace{\psi_{-r_1}^{i_1} \dots \psi_{-r_\ell}^{i_\ell}}_{\text{weight } r_1 + \dots + r_\ell} | 0 \rangle \mid \begin{array}{l} 1 \leq i_j \leq n, \ell = 0, 1, 2, \dots \\ r_j \in (\mathbb{Z} + \frac{1}{2})_{>0}, \{\psi_r^i, \psi_s^j\} = -2\delta^{ij} \delta_{r+s,0} \\ \{\psi_0^i, \psi_0^j\} = 0 \end{array} \right\}.$$

R sector (canonically-twisted module)

$$\Psi_0^i := \frac{1}{\sqrt{2}} (\psi_0^{2i-1} + \sqrt{-1} \psi_0^{2i}), \quad \bar{\Psi}_0^i := \frac{1}{\sqrt{2}} (\psi_0^{2i-1} - \sqrt{-1} \psi_0^{2i}) \quad \leftarrow \text{choice of the polarization}$$

As a vec. sp.,

$$F_R = \text{Span}_{\mathbb{C}} \left\{ \underbrace{\psi_{-r_1}^{i_1} \dots \psi_{-r_\ell}^{i_\ell} \Psi_0^{i'_1} \dots \bar{\Psi}_0^{i'_\ell}}_{\text{weight } r_1 + \dots + r_\ell + \frac{n}{16}} | 0 \rangle_R \mid \begin{array}{l} 1 \leq i_j, i'_j \leq n, \ell, \ell' = 0, 1, 2, \dots \\ r_j \in \mathbb{Z}_{>0}, \{\psi_r^i, \psi_s^j\} = -2\delta^{ij} \delta_{r+s,0} \end{array} \right\}.$$

• Spin(n) action

$\psi^1, \dots, \psi^n$: an orthonormal $\mathbb{C}$ -basis of $V \cong \mathbb{C}^n$

$$\begin{aligned} (\psi^i + \psi^j) \otimes (\psi^i + \psi^j) &= -(\psi^i)^2 - (\psi^j)^2 - 2\langle \psi^i, \psi^j \rangle \\ \psi^i \otimes \psi^i + \psi^j \otimes \psi^j &= -2\delta^{ij} \end{aligned}$$

$$\text{Cliff}(V) := \bigoplus_{i=0}^{\infty} V^{\otimes i} / v \otimes v \sim -|v|^2 \quad \{\psi^i, \psi^j\} = -2\delta^{ij}$$

$$\text{Spin}(n) \subset \text{Cliff}(V)$$

$$\begin{array}{ccccccc} 1 & \rightarrow & \mathbb{Z}_2 & \rightarrow & \text{Spin}(n) & \twoheadrightarrow & \text{SO}(n) \rightarrow 1 \\ & & & & \downarrow & & \downarrow \\ & & & & \pm \hat{M} & \mapsto & M \end{array} \quad \text{a double cover of } \text{SO}(n)$$

$$\hat{M}(\psi^1 \dots \psi^n) \begin{pmatrix} c_1 \\ \vdots \\ c_n \end{pmatrix} \hat{M}^{-1} = (\psi^1 \dots \psi^n) M \begin{pmatrix} c_1 \\ \vdots \\ c_n \end{pmatrix}$$

$\chi \in \text{Spin}(n)$ acts on

$$F_{NS} \text{ as } \psi_{-r_1}^{i_1} \dots \psi_{-r_k}^{i_k} |0\rangle \mapsto (\chi \psi_{-r_1}^{i_1} \chi^{-1})_{-r_1} \dots (\chi \psi_{-r_k}^{i_k} \chi^{-1})_{-r_k} |0\rangle \quad \text{SO}(n)\text{-action}$$

$$F_R \text{ as } \psi_{-r_1}^{i_1} \dots \psi_{-r_k}^{i_k} \mathbb{I}_0^{i_1} \dots \mathbb{I}_0^{i_k} |0\rangle_R \mapsto (\chi \psi_{-r_1}^{i_1} \chi^{-1})_{-r_1} \dots (\chi \mathbb{I}_0^{i_k} \chi^{-1})_0 \chi |0\rangle_R$$


$\uparrow$
 $\text{Cliff}(V)$
 $\uparrow$
 $\langle \psi_{-r_1}^{i_1}, \dots, \psi_{-r_k}^{i_k} \rangle \neq \langle \psi_0^{i_1}, \dots, \psi_0^{i_k} \rangle$

When n is even, $-1_{\text{SO}(n)} \in \text{SO}(n)$.

The lifts of $-1_{\text{SO}(n)}$ are $\pm \psi^1 \dots \psi^n =: \pm \mathcal{Z}$.

$$\text{Center}(\text{Spin}(n)) = \begin{cases} \mathbb{Z}_2 = \langle -1_{\text{Spin}(n)} \rangle & (n: \text{odd}) \\ \mathbb{Z}_4 = \langle \mathcal{Z} \rangle & (n \equiv 2 \pmod{4}) \\ \mathbb{Z}_2 \times \mathbb{Z}_2 = \langle -1_{\text{Spin}(n)} \rangle \times \langle \mathcal{Z} \rangle & (n \equiv 0 \pmod{4}) \end{cases}$$

If $n \equiv 0 \pmod{4}$,



parities of each sector:

fermion parity γ	$-1 \text{Spin}(n)$		
	+1	-1	
+1	F_{NS}^0	F_R^0	$n=24$: V^{fh}_{NS} (NS) $\leftarrow \text{SemiSpin}(24)$
-1	F_{NS}^1	F_R^1	$= V^{fh}_{tw}$ (R)
	$\parallel$	$\parallel$	
	F_{NS}	F_R	
	$\hookrightarrow$		
	$\text{SO}(n)$		

Annotations for the table:

- Left margin: $(-\gamma \rightarrow \text{different parity of R sector})$ just a convention
- Right margin: $\text{"} \langle \gamma \rangle \text{-orbifold"}$ and $\text{"} \langle -1 \text{Spin}(n) \rangle \text{-orbifold"}$ with arrows pointing to the F_{NS}^0 and F_R^0 sectors.

minimum-weight states of each sector:

$F_{NS}^0 \ni 0\rangle$	1-dim. wt. 0	$F_R^0 \ni (\text{even \# of } \mathbb{Z}_2 \text{'s}) 0\rangle_R$	2"-dim. wt. $\frac{3}{2}$	positive chiral rep. of $\text{Spin}(24)$
$F_{NS}^1 \ni \psi_{-\frac{1}{2}}^i 0\rangle$	24-dim. wt. $\frac{1}{2}$	$F_R^1 \ni (\text{odd } \text{---}) 0\rangle_R$	2"-dim. wt. $\frac{3}{2}$	negative ---

N_0 weight $-\frac{1}{2}$ states in V^{fh} .

• $C_{0,}$ action on V^{fq}

Fact

$$\Downarrow \quad \equiv C_{0,} \subset Spin(24)$$

$$\swarrow \quad \searrow \quad \parallel 2$$

$$SemiSpin(24) \supset C_{0,} \\ \downarrow \\ V^{fq}$$

$$C_{0,} = Aut(\Lambda_{24}) \subset SO(24)$$

Fact $det = +1$ 

In the 2^{11} -dim. weight $-\frac{3}{2}$ subsp. $(V^{fq})_{\frac{3}{2}}$,

there exists a 1-dim. inv. subsp. under the $C_{0,}$ action.

Its basis G constitutes the supercurrent of the $C_{0,}$ -inv. $N=1$ SCA.

$$\rightarrow Aut_{N=1}(V^{fq}) \cong C_{0,}.$$

Stolz-Teichner? proposal? program?

3.2. Connection to the Segal-Stolz-Teichner conjecture

Conj cf. [1108.0189 Stolz-Teichner]

$$SQFT \cong TMF.$$

↑ the space ($\mathbb{Z}$ -graded ring)
of the classes of topological modular forms

↑ the space ($\mathbb{Z}$ -graded ring)
of the equiv. classes of 2d $N=(0,1)$ SQFTs

For $[\tau] \in SQFT_n$, degree

first
Pontryagin class

· the anomaly polynomial of the grav. anom. = $\frac{n}{48} p_1$

· $SQFT_n \xrightarrow{\sim} TMF_n \rightarrow (MF_{\mathbb{Z}}^{w-h})_{\frac{n}{2}}$ weakly holomorphic modular forms
with $\mathbb{Z}$ coefficient of weight $\frac{n}{2}$

$$f\left(\frac{a\tau+b}{c\tau+d}\right) = (c\tau+d)^{\frac{k}{2}} f(\tau)$$

$$\downarrow \quad \quad \quad \downarrow$$

$$[\tau] \longmapsto \eta(\tau)^n Z_{\text{ell}}\tau$$

$$Z_{\text{ell}}\tau := \text{Tr}_{\mathcal{H}_{\tau, R\text{-sector}}} (-1)^F q^{\frac{H+P}{2}} \bar{q}^{\frac{H-P}{2}}$$

If τ is an SCFT,

· $n = 2(c_R - c_L)$

· $Z_{\text{ell}}\tau = \text{Tr}_{\mathcal{H}_{\tau, R\text{-sector}}} (-1)^{F_L + F_R} q^{L_0 - \frac{c_L}{24}} \bar{q}^{\tilde{L}_0 - \frac{c_R}{24}}$

↑
indep. of $\bar{q}$
(the function of q)

If we assume that the SST conj. is true,

we can obtain nontrivial statements on SQFT.

by translating the properties of TMF.

· Example 1. the divisibility property

$TMF_n \rightarrow (MF_{\mathbb{Z}}^{w-h})_{\frac{n}{2}}$ is not surjective.

Its image is also determined. (Hopkins, Mahowald) cf. [Hopkins Math/0212397 Prop. 4.6]

In particular,

$$(\text{its image}) \cap \mathbb{Z}[\eta(\tau)] = \begin{cases} \mathbb{Z} \left[\frac{24}{\gcd(24, \frac{n}{24})} \eta(\tau)^n \right] & (n \equiv 0 \pmod{24}) \\ 0 & (\text{otherwise}) \end{cases}$$

Therefore, if the SST conj. is true, then for $[\tau] \in \text{SQFT}_n$,

$$\mathbb{Z}_{\text{ell}}\tau \text{ is const.} \Rightarrow \begin{cases} \frac{24}{\gcd(24, \frac{n}{24})} \mid \mathbb{Z}_{\text{ell}}\tau & (24 \mid n) \\ \mathbb{Z}_{\text{ell}}\tau = 0 & (\text{otherwise}) \end{cases}$$

For example,

$$V^{f_4} : (c_L, c_R) = (12, 0) \quad n = -24$$

$$\frac{24}{\gcd(24, \frac{n}{24})} = 24 \quad \mid \quad \mathbb{Z}_{\text{ell}}[V^{f_4}](\tau) = -24$$

· Example 2. the periodicity element

Fact

There exists a periodicity element $X \in \mathrm{TMF}_{-24^2} = -576$ s.t.

$$X \cdot : \mathrm{TMF}_n \xrightarrow{\sim} \mathrm{TMF}_{n-576}.$$

Under the SST conj., this translates to the existence of

an $N=(0,1)$ SQFT τ of $N=-576$ s.t. $Z_{\mathrm{ell}}[\tau] = 1$.

It suffices to find

an $N=1$ chiral SCFT τ of $C=288$ s.t. $Z_{\mathrm{ell}}[\tau] = 1$.

$$\downarrow$$

$$Z_{\mathrm{ell}}\tau = \text{const.}$$

$(\mathrm{Vfh})^{\otimes 24}$ has $C = 24 \times 12 = 288$.

[1909.08388 Johnson-Freyd]

The anomaly of its C_0 sym. is described by the generator of $\mathrm{SH}(C_0) \cong \mathbb{Z}_{24}$.

$\leadsto (\mathrm{Vfh})^{\otimes 24} / C_0$ has $Z_{\mathrm{ell}}[\tau] = 1$?

$(\mathrm{Vfh})^{\otimes 24} / (C_0 \times A_{24})$ $\neq$? [2210.14923 Albert, Lin, Kaidi]

$C_0 \times S_{24}$ is anomalous.

4. Mathieu moonshine

cpt. Kähler mfd. with triv. can. bdl.

K3 surface : a cpx-2-dim. CY mfd. with $h^{0,1} = 0$.

$$\begin{smallmatrix} 0 & 1 \\ 0 & 20 & 0 \\ 1 & 0 & 0 \end{smallmatrix}$$

K3-target SCFT : $\mathcal{N} = (4, 4)$, $(c_L, c_R) = (6, 6)$

The elliptic genus the weak Jacobi form of weight 0 and index $\frac{c_L}{6} = 1$

$$Z_{K3}(\tau, z) := \text{Tr}_{\mathcal{H}_{K3}} (-1)^{F_L + F_R} y^{J_0} q^{L_0 - \frac{c_L}{24}} \bar{q}^{\tilde{L}_0 - \frac{c_R}{24}} \quad \left(\begin{matrix} \tau = e^{2\pi i \tau} \\ y = e^{2\pi i z} \end{matrix} \right)$$

$$= 2\mathcal{Y}_{0,1}(\tau, z)$$

$$= 2 \{ y + 10 + y^{-1} + (10y^2 - 64y + 108 - 64y^{-1} + 10y^{-2})y + \dots \}$$

If we decompose Z_{K3} into the irr. char. of the $\mathcal{N} = 4$ SCA of $c = 6$,

$$Z_{K3}(\tau, z) = 20 \text{ch}_{\frac{1}{4}, 0}^{(\text{massless})}(\tau, z) - 2 \text{ch}_{\frac{1}{4}, \frac{1}{2}}^{(\text{massless})}(\tau, z)$$

$$+ 2 \sum_{n=1}^{\infty} A_n \text{ch}_{n+\frac{1}{4}, \frac{1}{2}}(\tau, z).$$

n	1	2	3	4	5	6	...
A_n	<u>45</u>	<u>231</u>	<u>770</u>	<u>2277</u>	<u>5796</u>	13915	...
						<u>3520</u> + <u>10395</u>	

dim. of irrep. of M_{24} :

1, 23, 45, 45, 231, 231, 252, 253, 483, 770, 770,

970, 970, 1035, 1035, 1035, 1265, 1771, 2024,

2277, 3312, 3520, 5313, 5796, 5544, 10395

→ underlying VOA-like object?

K3 Mathieu moonshine (三上, 大栗, 立川, 2010)

other evidence of the existence of VOA

- twined ell. genera (the McKay-Thompson series) ^{Cheng (2010)}
Gabberdiel, Hohenegger, Volpato (2010)
- $\exists$ M_{24} -module reproducing them. ^{Gannon (2012)}
- twisted twined ell. genera

Their mod. trsf. are controlled by a 3-cocycle in $H^3(M_{24}; U(1))$.

(fit into the theory of anomaly) ^{Gabberdiel, Persson, (2012)}
Ronellenfisch, Volpato

However, ^{preserve the hol. 2-form}

- $\text{Aut}_w(K3) \subsetneq M_{23}$ ^{Mukai (1988)}
- $M_{24} \not\subset \text{Aut}_{N=(4,4)}(K3 \text{ CFT}) \subsetneq \text{Co}_1$ ^{Gabberdiel, Hohenegger, Volpato (2010)}

The underlying VOA-like object is not yet found!

solution?

- symmetry surfing? ^{Taormina, Wendland (2011-13)}
- reflected $K3 = V^{fh}$? ^{Duncan, Mack-Crane (2015)}
^{Taormina, Wendland (2019)}
- preserve only $N=(4,1)$? ^{Harvey, Moore (2020)}

some subalgebra?

- non-inv. sym.? ^{Volpato (2024)}
^{Anzies, Giaccari, Harrison, Volpato (2025)}

⋮