

Elementary Weak Solutions of the Navier-Stokes Equations by a New Fixed Point Theorem

Masatoshi Ohnri

August 23, 2026

Abstract

We solve the problem in mathematics. The problems are not in physics, so we do not use any physics or assumptions-falsified mathematics.

We use mathematics only. We can solve the problem by using an exactly and completely FALSIFIED resolution, where large initial values destroy the earth, because uniqueness does NOT hold, or SMALL initial values love your cup of coffee.

There are no long or complicated calculations, or no theory of evolution equations.

1 Introduction

The existence of solutions is actually known. For example, Fujita-Kato Theory, Shibata Theory: Kato [4], Zhang [12], Charve-Danchin [2], Shibata-Murata [6]. The semi-group theory or a priori estimates are in these theories, but these are not elementary. Semi-group theory or a priori estimates for contraction mappings are not used in the proof of the existence of Leray-Hopf's weak solutions (for example, Wasao Sibagaki and Hisako Rikimaru [7]), but they are not elementary either. We define new weak solutions with uniqueness and smoothness, without semi-group theory or a priori estimates. We apply the local solvability of the partial differential operators with constant coefficients. The policy is to let L be the heat operator $\partial_t - \Delta$ in the initial value problem of the Navier-Stokes equations in

$$(N-S) \quad \partial_t u - \Delta u = f - \nabla p - (u \cdot \nabla)u$$

$$(\sigma) \quad \operatorname{div}(u) = 0,$$

$$(A) \quad u(0, x) = a(x), (0, x) \in \Omega$$

$$(S) \quad \lim_{t, x \rightarrow \infty} \partial^\alpha u(t, x) = 0$$

$$(C) \quad f \mapsto u, a \mapsto u \text{ are continuous.}$$

The uniqueness follows without boundary conditions. Our solutions are more rapidly decreasing than [4], satisfy the energy inequity, and analytic. There are no long or complicated calculations; semi-groups, a priori estimates, or boundary conditions on $\Omega' \subset R^3$ are not used at all. We apply the local solvability of linear partial differential operators with constant coefficients ([8], [3]), erase the pressure p by the Helmholtz projection P , and solve

$$\begin{aligned}\frac{\partial u}{\partial t} - \Delta u &= Pf - P((u_n \cdot \nabla)u_n), \\ \frac{\partial u}{\partial t} - \Delta u &= Pf - P((u \cdot \nabla)u),\end{aligned}$$

in $X = \bigcap_{m \geq 5, p=1,2} W^{m,p}(\Omega)$, $\Omega \subset \mathbb{R} \times \mathbb{R}^3$ is a bounded domain with Lipschitz boundary such that $(0,0) \in \Omega$.

For example, cylinder sets $\Omega = I \times \Omega' = (-T, T) \times B_3(0, R)$. We handle the variables t, x simultaneously. Cylinder sets are Lipschitz domains, so we can use the Sobolev embedding theorem in Ω and Helmholtz decomposition in Ω' .

2 Definition

For convenience, we write the index of the component of the vectors in the upper right corner.

"Function space" and "space" are abbreviations for "linear topological space" (of functions or distributions), other than pressure $\mathbf{p}$, which are $\mathbb{R}$ -valued. The absolute value of the functions in the norm of the normal function space is interpreted as the length of the number vector (the absolute value of $\mathbb{R}^3$ in the norm of the space of the $\mathbb{R}$ -valued functions). We write the space of the real numeric functions and the space of the $\mathbb{R}^3$ -valued functions using the same symbol to keep the notation simple. Let $|\Omega|$ be the Lebesgue measure of Ω . Let χ_Ω be the characteristic function on Ω . For any natural number $m > \max\{0 + 4/1, 0 + 4/2\} = 4$, $p = 1, 2$, We define the function spaces on $\mathbb{R} \times \mathbb{R}^3$:

$$\begin{aligned}V^{m,p}(\Omega) &= \left\{ u = (u^1, u^2, u^3) : u^i \in C^\infty(\Omega), \|u^i\|_{W^{m,p}(\Omega)} < \infty \right\}, \\ V_\sigma^{m,p}(\Omega) &= \{ u \in V^{m,p}(\Omega) : \operatorname{div}(u) = 0 \}.\end{aligned}$$

Let $W^{m,p}(\Omega)$ and $W_\sigma^{m,p}(\Omega)$ be the Sobolev spaces defined by the completions of $V^{m,p}(\Omega)$ and $V_\sigma^{m,p}(\Omega)$ (cf. [11]).

Let $\mathcal{D}(\Omega)$ be the space of the test functions ($C_0^\infty(\Omega)$ as a set); let $\mathcal{D}_\sigma(\Omega)$ be the space of the test functions for which the divergence is 0 with respect to spatial variables. Let $P: L^2(\Omega) \rightarrow L_\sigma^2(\Omega) = \overline{\mathcal{D}_\sigma(\Omega)}^{\|\cdot\|_{L^2(\Omega)}}$ be the projection. Let $C^{k,\varepsilon}(\Omega)$ be the Hölder space. Let

$$\begin{aligned}\langle w, \varphi \rangle &:= (w, \varphi)_{L^2(\Omega)} \\ &= \int_\Omega w(t, x) \cdot \varphi(t, x) \, dt \, dx \\ &= \int_\Omega \sum_{i=1}^3 w^i(t, x) \varphi^i(t, x) \, dt \, dx\end{aligned}$$

for $w = (w^1, w^2, w^3)$, $\varphi = (\varphi^1, \varphi^2, \varphi^3)$.

In general, if for two Banach spaces X, Y , there exists a linear Hausdorff space Z such that $X, Y \subset Z$, then $X \cap Y$ is a Banach space with norms given by $\|u\|_X + \|u\|_Y$ or $\max\{\|u\|_X, \|u\|_Y\}$.

$$\max\{\|u\|_X, \|u\|_Y\} \leq \|u\|_X + \|u\|_Y \leq 2 \max\{\|u\|_X, \|u\|_Y\}$$

so these are equivalent. We put

$$\begin{aligned}\mathcal{X} &= \bigcap_{m \geq 5} W_\sigma^{m,1}(\Omega) \cap W_\sigma^{m,2}(\Omega) \\ \mathcal{X}' &= \bigcap_{m \geq 5} W^{m,1}(\Omega) \cap W^{m,2}(\Omega)\end{aligned}$$

We define for any $u \in \mathcal{X}$,

$$\|u\|_X = \sum_{m=5}^{\infty} \sum_{k=0}^{\infty} \frac{1}{m!^4 (c'_{k,m})^2} \|u\|_{W^{m,1}(\Omega) \cap W^{m,2}(\Omega)},$$

and for any $u \in \mathcal{X}'$,

$$\|u\|_{X'} = \sum_{m=5}^{\infty} \sum_{k=0}^{\infty} \frac{1}{m!^4 (c'_{k,m})^2} \|u\|_{W^{m,1}(\Omega) \cap W^{m,2}(\Omega)}.$$

We put $X = \{u \in \mathcal{X} : \|u\|_X < \infty\}$, $X' = \{u \in \mathcal{X}' : \|u\|_{X'} < \infty\}$. For constants $M, C > 0$, let S be a subset of X :

$$S = \{u \in X : \|u\|_X \leq M, \|\partial_{x^j} u\|_X \leq C \|u\|_X\}.$$

Let E be the fundamental solution of $\partial_t - \Delta$. That is, in the sense of that $\mathbb{R}^3$ -valued distribution,

$$(\partial_t - \Delta)E(t, x) = \delta(t, x) = \delta(t) \otimes \delta(x).$$

Here,

$$E(t, x) = \begin{cases} \frac{1}{\sqrt{4\pi t}^3} \exp\left(-\frac{|x|^2}{4t}\right) & (t > 0) \\ 0 & (t \leq 0) \end{cases}.$$

(cf. [8], [3]).

3 Existence of elementary weak solutions

Assumption. Let $\Omega \subset \mathbb{R} \times \mathbb{R}^3$ be a bounded domain with Lipschitz boundary such that $(0, 0) \in \Omega$. Let $f : \mathbb{R} \times \mathbb{R}^3 \rightarrow \mathbb{R}^3$ be an external force satisfying

$Pf \in S$. When $f \neq 0$, we further assume that there exists $(0, x) \in \Omega$ such that

$$\int_{\mathbb{R} \times \mathbb{R}^3} E(s, y)(\chi_\Omega Pf)(0 - s, x - y) ds dy \neq 0.$$

Let A denote the set of initial values given by

$$A = \left\{ u(0, \cdot) : u \in X, u(t, x) = \int_{\mathbb{R} \times \mathbb{R}^3} E(s, y)(\chi_\Omega P(f - (u \cdot \nabla)u))(t - s, x - y) ds dy \right\}.$$

Proposition 1 (Local solvability of linear partial differential operator with constant coefficients). Let L be a linear partial differential operator with constant coefficients in $\mathbb{R}^N$, and let $E \in \mathcal{D}'(\mathbb{R}^N)$ be a fundamental solution of L , i.e., $LE = \delta$. For $f \in L^1_{\text{loc}}(\mathbb{R}^N)$ and a bounded domain $\Omega \subset \mathbb{R}^N$, a weak solution $u \in \mathcal{D}'(\Omega)$ to the equation $Lu = f$ in Ω is given by $u = E * \chi_\Omega f \in \mathcal{D}'(\Omega)$.

Proof. For any $\varphi \in \mathcal{D}(\Omega)$,

$$\begin{aligned} \langle L(E * \chi_\Omega f), \varphi \rangle &= \pm \langle E * \chi_\Omega f, L\varphi \rangle \\ &:= \pm \langle E(x), \langle \chi_\Omega(y)f(y), L\varphi(x + y) \rangle \rangle \\ &= \pm \langle \chi_\Omega(x)f(x), \langle E(y), L\varphi(x + y) \rangle \rangle \\ &= \langle \chi_\Omega(x)f(x), \langle LE(y), \varphi(x + y) \rangle \rangle \\ &= \langle LE(x), \langle \chi_\Omega(y)f(y), \varphi(x + y) \rangle \rangle \\ &= \langle \chi_\Omega(y)f(y), \varphi(y) \rangle \\ &= \langle f, \varphi \rangle \end{aligned}$$

□

Theorem 1 (Main theorem). $A \neq \emptyset$. For $a \in A$, there are elementary weak solutions $u, \mathbf{p}$ of the initial value problem in Ω :

$$\begin{aligned} \frac{\partial u}{\partial t} - \Delta u &= f - \nabla \mathbf{p} - (u \cdot \nabla)u, \\ \operatorname{div}(u) &= 0, \\ u(0, x) &= a(x). \end{aligned}$$

in the sense that $u \in X, \mathbf{p} \in L^2_{\text{loc}}(\Omega)/(p \sim q \iff \nabla(p - q) = 0)$ and $u, \mathbf{p}$ satisfy for any $\varphi \in \mathcal{D}_\sigma(\Omega)$,

$$\langle \partial_t u + (u \cdot \nabla)u - \Delta u + \nabla \mathbf{p} - f, \varphi \rangle = 0,$$

for any $\varphi \in \mathcal{D}(\Omega)$,

$$\langle \operatorname{div}(u), \varphi \rangle = - \sum_{j=1}^3 \langle u^j, \partial_{x^j} \varphi \rangle = 0.$$

For any multi-index $\alpha \in \mathbb{Z}_{\geq 0}^4$,

$$\lim_{t, |x| \rightarrow \infty} \partial^\alpha u(t, x) = 0.$$

If solutions $u, \mathbf{p}$ and $v, \mathbf{q}$ satisfy $u, v \in S, u - v \in S$, then $u = v, \mathbf{p} = \mathbf{q}$ on Ω . The map $f \mapsto u$ is continuous.

We prove Theorem 1 later. The method is similar to the proof of Banach fixed-point theorem. f defines A and the proof of existence does not depend on A .

Proposition 2 (Smoothness of elementary weak solutions). The solutions $(u, \mathbf{p})$ are C^∞ -functions.

Proof. $m \in \mathbb{N}$ can be taken arbitrarily large, so the embedding theorem into Hölder space ([1]),

$$\text{If } \mathbb{N} \ni m - 4/p > 0, \text{ then } W^{m,p}(\Omega) \subset C^{(m-4/p)-1,\varepsilon}(\Omega) \text{ for all } \varepsilon \in (0, 1).$$

in the sense of existence of suitable representative elements, u is a C^∞ -function.

f is smooth and $\partial_t u + (u \cdot \nabla)u - \Delta u - f = -\nabla \mathbf{p}$. Because $-\nabla \mathbf{p}$ is smooth, so $\mathbf{p}$ is also smooth. $\square$

Lemma 1. The normed spaces X and X' are non-trivial.

Proof. It suffices to exhibit non-zero elements in each space. We note that χ_Ω and $\text{curl}(\exp(x^1), \exp(x^3), \exp(x^2)) \in X$, hence $X \neq 0$. Similarly, $\chi_\Omega \in X'$ and $\exp(x^1), \sin(x^2) \in X'$, so $X' \neq 0$. $\square$

Lemma 2 (Completeness). X, X' are Banach spaces.

Proof. Let $\{u_n\}$ be a Cauchy sequence in X . $\{u_n\}$ is a Cauchy sequence of $W_\sigma^{m,1}(\Omega) \cap W_\sigma^{m,2}(\Omega)$. Since $W^{m,1}(\Omega) \cap W^{m,2}(\Omega)$ are Banach spaces, $\{u_n\}$ converges in $W^{m,1}(\Omega) \cap W^{m,2}(\Omega)$. Let u be the limit of $\{u_n\}$ in $W^{m,1}(\Omega) \cap W^{m,2}(\Omega)$. For any $\varepsilon > 0$, there exists $N \in \mathbb{N}$ such that if $l, n \geq N$, then $\|u_l - u_n\|_X < \varepsilon$. Using Fatou's lemma for counting measures,

$$\begin{aligned} \|u - u_n\|_X &= \sum_{m=5}^{\infty} \sum_{k=0}^{\infty} \frac{1}{m!^4 (c'_{k,m})^2} \|u - u_n\|_{W^{m,1}(\Omega) \cap W^{m,2}(\Omega)} \\ &= \sum_{m=5}^{\infty} \sum_{k=0}^{\infty} \frac{1}{m!^4 (c'_{k,m})^2} \liminf_{l \rightarrow \infty} \|u_l - u_n\|_{W^{m,1}(\Omega) \cap W^{m,2}(\Omega)} \\ &\leq \liminf_{l \rightarrow \infty} \sum_{m=5}^{\infty} \sum_{k=0}^{\infty} \frac{1}{m!^4 (c'_{k,m})^2} \|u_l - u_n\|_{W^{m,1}(\Omega) \cap W^{m,2}(\Omega)} \\ &\leq \varepsilon. \end{aligned}$$

Therefore,

$$\|u\|_X \leq \|u - u_N\|_X + \|u_N\|_X \leq \varepsilon + \|u_N\|_X < \infty,$$

hence $u \in X$. $\square$

Lemma 3 (Separation of product). There exists a constant $C_1 > 0$ such that

$$\|u^i v^i\|_{X'} \leq C_1 \|u^i\|_{X'} \|v^i\|_{X'}$$

holds for all $u, v \in X'$.

Proof. For multi-indices $\alpha, \beta \in \mathbb{Z}_{\geq 0}^4$, let $c_{\alpha, \beta}$ denote the binomial coefficients. Define

$$c_\alpha = \sum_{\beta \leq \alpha} c_{\alpha, \beta}.$$

There is a continuous embedding $X' \subset C^{k, \varepsilon}(\overline{\Omega})$ for any $k \in \mathbb{N}$ because

$$\begin{aligned} \|u_n - u\|_X \rightarrow 0 &\implies \|u_n - u\|_{W^{m,1}(\Omega) \cap W^{m,2}(\Omega)} \rightarrow 0 \\ &\implies \|u_n - u\|_{C^{k, \varepsilon}(\overline{\Omega})} \rightarrow 0. \end{aligned}$$

([1]), so there exists a constant $c'_{k,m} > 0$ such that

$$\|u\|_{C^{k, \varepsilon}(\overline{\Omega})} \leq c'_{k,m} \|u\|_{X'}.$$

By Leibniz formula,

$$\begin{aligned} \|\partial^\alpha(uv)\|_{L^p(\Omega)} &\leq c_\alpha \|u\|_{C^{k, \varepsilon}(\overline{\Omega})} \|v\|_{C^{k, \varepsilon}(\overline{\Omega})} |\Omega|^{1/p} \\ &\leq c_\alpha c'_{k,m} |\Omega|^{1/p} \|u\|_{X'} c'_{k,m} \|v\|_{X'} \\ &\leq c_\alpha (c'_{k,m})^2 |\Omega|^{1/p} \|u\|_{X'} \|v\|_{X'}, \end{aligned}$$

for $|\alpha| \leq k$. Therefore, there exists a constant $C'_1 > 0$ such that

$$\|uv\|_{X'_{\text{new}}} \leq C'_1 \|u\|_{X'_{\text{old}}} \|v\|_{X'_{\text{old}}}.$$

For any old $c'_{k,m}, c'_{k+1,m+1} \geq c'_{k,m} > 0$, there exists a new $c''_{k,m} := c'_{k,m}$. We correct : $c''_{k,m} \geq c'_{k,m} \geq \sqrt{\alpha_{0,5}} c''_{k,m}$.

We can prove Lemma 3 exactly.

If there exists $0 < \alpha = \alpha_{c', c''} < 1$ such that

$$\alpha \|uv\|_{X'_{\text{old}}} \leq \|uv\|_{X'_{\text{new}}} \leq C'_1 \|u\|_{X'_{\text{old}}} \|v\|_{X'_{\text{old}}}.$$

Hence, it is sufficient to take $C_1 := C'_1 / \alpha$ and

$$\|uv\|_{X'_{\text{old}}} \leq C_1 \|u\|_{X'_{\text{old}}} \|v\|_{X'_{\text{old}}}.$$

$$\begin{aligned}
& c'_{k,m} \geq c''_{k,m} \\
& \|u\|'' \leq \|u\|' \\
& \left\{ \begin{aligned} & \|u_m - u_n\|' \rightarrow 0 \\ & \|u_m - u_n\|'' \rightarrow 0 \end{aligned} \right\} \wedge \left\{ \begin{aligned} & \|u_m - u_n\|' \rightarrow 0 \\ & \|u_m - u_n\|'' \rightarrow 0 \end{aligned} \right\} \\
& 0 \leq \|u' - u''\|'' \\
& \leq \|u' - u_n\|'' + \|u_n - u''\|'' \\
& \leq \|u_n - u'\|' + \|u_n - u''\|'' \rightarrow 0 \\
& \therefore u' = u'' \therefore u_m - u_n \rightarrow 0 \text{ in } X_{c',c''} \\
& \Rightarrow \exists u \in X_{c',c''}, u_n \rightarrow u \text{ in } X_{c',c''}
\end{aligned}$$

We have $\|u\|_{C^{0,\varepsilon}(\bar{\Omega})} \leq \|u\|_{C^{k,\varepsilon}(\bar{\Omega})} \leq c''_{k,m} \|u\|_X$,
 $\|u\|_{C^{0,\varepsilon}(\bar{\Omega})} \leq c''_{0,5} \|u\|_X$. Therefore we can take

$$c''_{k,m}$$

as a constant. (Reference: Haim Brezis, *Functional Analysis, Sobolev Spaces and Partial Differential Equations*, Springer, 2010.) To justify Lemma 3, take

$$c'_{k,m} \geq 1.$$

(Hisashi Okamoto, *Principal of the Navier-Stokes Equations*, University of Tokyo Press, 2023, ISBN: 978-4-13-061315-6;

Yoshihiro Shibata, *Basic of Mathematical Fluids 1, 2*, Iwanami books, 2022, ISBN: 9784000298582, 9784000298599.)

From

$$\frac{1}{(c'_{k,m})^2} \leq \frac{1}{(c''_{k,m})^2} \frac{(c''_{k,m})^2 m!^4}{(c'_{k,m})^2},$$

if


$$m'' \geq m, \quad k'' \geq k,$$

then

$$\begin{aligned}
& \frac{1}{m''!^4 (c''_{k'',m''})^2} \frac{1}{m!^4 (c'_{k,m})^2} \|u_{m'} - u_{n'}\|_{W^m} \\
& \leq \frac{1}{m!^4 (c''_{k,m})^2} \|u_{m'} - u_{n'}\|_{W^m} \frac{1}{(c'_{k,m})^2} \\
& \leq \frac{1}{m!^4 (c''_{k,m})^2} \|u_{m'} - u_{n'}\|_{W^m}.
\end{aligned}$$

Therefore, there exists constant $C' > 0$ such that

$$C' \|u_{m'} - u_{n'}\|' \leq \|u_{m'} - u_{n'}\|''.$$



To obtain α , choose $0 < \alpha_{k,m} \leq 1$ so that

$$\alpha_{0,5} < \alpha_{k+1,m+1} \leq \alpha_{k,m},$$

and

$$0 < \frac{\alpha_{k,m}}{(c'_{k,m})^2} \leq \frac{1}{(c''_{k,m})^2}.$$

Using

$$\frac{\alpha}{(c'_{k,m})^2} \leq \frac{\alpha_{k,m}}{(c'_{k,m})^2} \leq \frac{1}{(c''_{k,m})^2}$$

and

$$\exp\left(1 + \frac{1}{\alpha_{1,6}}\right) \geq \exp\left(1 + \frac{1}{\alpha_{k,m}}\right) \geq \frac{1}{\alpha_{k,m}} \geq 1,$$

for

$$\alpha = \frac{1}{\inf_{k,m} \exp\left(1 + \frac{1}{\alpha_{k,m}}\right)}.$$

We can also choose $c_{k,m}'$ and justify the existence of α by using [10, Problem 9.6]: “If the linear space

$$X_{c',c''} := \left\{ u \in \bigcap_{m \geq 5} \bigcap_{p=1}^2 W^{m,p}(\Omega) : \|u\|' < \infty, \|u\|'' < \infty \right\}$$

is complete with respect to the norms $\|\cdot\|'$ and $\|\cdot\|''$, and if $\|\cdot\|'' \leq \|\cdot\|'$, then these norms are equivalent.”

□

Lemma 4 (Absorption of differential). There exists a constant $C_2 > 0$ such that



$$\|\partial_{x^j} u\|_X \leq C_2 \|u\|_X$$

for any $u \in S$.

Lemma 5

$$\left\| \int_{\mathbb{R} \times \mathbb{R}^3} E(s, y) \chi_\Omega(t-s, x-y) u(t-s, x-y) ds dy \right\|_X \leq C_3 \|u\|_X,$$

$$\left\| \int_{\mathbb{R} \times \mathbb{R}^3} E(s, y) \partial_{x^j} \chi_\Omega(t-s, x-y) u(t-s, x-y) ds dy \right\|_X \leq C_3 C_2 \|u\|_X$$



Let $I = (0, 1)$. Since

$$t^{-3/2}t^2 = \sqrt{t} > t^2 \quad (t \in I),$$

therefore $I = (0, 1)$,

$$u(t, x) = \frac{1}{\sqrt{4\pi}} t^2 x_1 x_2 x_3$$

are the counterexamples.

We correct Lemma 5:

$$\exists C_3 > 0, \quad \forall u \in S \setminus \{v \in S : \exists \alpha, \partial^\alpha v = 0\},$$

$$E * \chi_\Omega u \in X \setminus \{v \in S : \exists \alpha, \partial^\alpha v = 0\},$$

and

$$\|E * \chi_\Omega u\|_X \leq C_3 \|u\|_X,$$

$$\|E * \partial_{x^j}(\chi_\Omega u)\|_X \leq C_3 C_2 \|u\|_X.$$

To prove Lemma 5, we don't derivative χ_Ω , derivative

$$\chi_\Omega u = u,$$

consider the common derivative almost everywhere on Ω , and correct the value of

$$E(s, y)U(t-s, x-y)$$

for fixed a.e. $(t, x) \in \Omega$:

$$E(s, y)U(t-s, x-y), \quad (s, y) \in -\Omega + (t, x);$$

$$E(s, y)U(t-s, x-y), \quad (s, y) \in \Omega.$$

By second inequality of Lemma 5, and Lemma 7, we can construct the elementary weak solutions even if

$$\Phi[S] \not\subset S.$$

$$\begin{aligned} & \int_\Omega \partial^\alpha (E * \chi_\Omega U) \varphi \\ &= (-1)^{|\alpha|} \int_\Omega (E * \chi_\Omega U) \partial^\alpha \varphi \\ &= (-1)^{|\alpha|} \int_\Omega \int_{-\Omega + (t, x)} E(s, y) U(t-s, x-y) ds dy \partial^\alpha \varphi(t, x) dt dx \\ &= (-1)^{|\alpha|} \int_\Omega \int_\Omega E(s, y) U(t-s, x-y) ds dy \partial^\alpha \varphi(t, x) dt dx \\ &= \int_\Omega \int_\Omega E(s, y) \partial^\alpha U(t-s, x-y) ds dy \varphi(t, x) dt dx. \end{aligned}$$

Proof. As a function of (s, y) , for almost every $(t, x) \in \Omega$,

$$\begin{aligned} \text{supp}(E(s, y)(\chi_\Omega u)(t - s, x - y)) &\subseteq -\bar{\Omega} + (t, x) \\ &= \{(s, y) \in \mathbb{R} \times \mathbb{R}^3 : (t - s, x - y) \in \Omega\} \end{aligned}$$

is the translation of the reverse of Ω , so it is compact, and

$$\begin{aligned} |\partial_{t,x}^\alpha (E(s, y)(\chi_\Omega u)(t - s, x - y))| &\leq E(s, y) \sup\{|\partial_{t,x}^\alpha u(t - s, x - y)| : (t, x) \in \Omega\} \\ &\leq C_\alpha E(s, y) \in L_{s,y}^1(\Omega). \end{aligned}$$

Combining the theorem of differentiation under the integral sign, Hölder's inequality ([11]), and continuous embedding $S \subset L^\infty(\Omega)$, we have

$$\begin{aligned} \|\partial^\alpha (E * (\chi_\Omega u))\|_{L^p(\Omega)} &\leq \|E * (\partial^\alpha (\chi_\Omega u))\|_{L^p(\Omega)} \\ &\leq \left\| \|E\|_{L^1(-\Omega+(t,x))} \|\partial^\alpha u(t - s, x - y)\|_{L^\infty(-\Omega+(t,x))} \right\|_{L^p(\Omega)} \\ &\leq \sup\left\{ \|E\|_{L^1(-\Omega+(t,x))} : (t, x) \in \Omega \right\} |\Omega|^{1/p} \|\partial^\alpha u\|_{L^\infty(\Omega)} \\ &\leq \sup\left\{ \|E\|_{L_{s,y}^1(-\Omega+(t,x))} : (t, x) \in \Omega \right\} c'' C_2^{|\alpha|} |\Omega|^{1/p} \|u\|_X \\ &< \infty. \end{aligned}$$

Therefore, we obtain

$$\|E * (\chi_\Omega u)\|_X \leq C_3 \|u\|_X.$$

□

We take $C = \max\{C_1, C_2, C_3\}$. Note that Lemma 3, Lemma 4, and Lemma 5 hold for $C = O(|\Omega|)$. We take $\Omega, f, M > 0$ satisfying

$$C\|f\|_{X'} + 3C^3 M^2 \leq M.$$

We solve

$$(N-S)' \quad \partial_t u - \Delta u = f - (u \cdot \nabla)u,$$

that is, for any $a \in A$, there exist $u \in S$ and $\mathbf{p} \in L^2(\Omega)$ such that

$$\begin{cases} \langle \partial_t u + (u \cdot \nabla)u - \Delta u + \nabla \mathbf{p} - f, \varphi \rangle = 0 & (\varphi \in \mathcal{D}_\sigma(\Omega)), \\ \langle \text{div}(u), \varphi \rangle = -\sum_{j=1}^3 \langle u^j, \partial_{x^j} \varphi \rangle = 0 & (\varphi \in \mathcal{D}(\Omega)), \\ u(0, x) = a(x). \end{cases}$$

Define $\Phi: S \rightarrow X$ as follows:

$$\Phi[u](t, x) = \int_{\mathbb{R} \times \mathbb{R}^3} E(s, y) (\chi_\Omega P(f - (u \cdot \nabla)u))(t - s, x - y) ds dy.$$

We take a sequence $\{u_n\} \subset X$ as $u_0 \in S, u_1 \in S, u_1 - u_0 \in S$; if $n \geq 1$ then

$$\begin{aligned} u_{n+1}(t, x) &= \Phi[u_n](t, x) \\ &= \int_{\mathbb{R} \times \mathbb{R}^3} E(s, y) (\chi_\Omega P(f - (u_n \cdot \nabla)u_n))(t - s, x - y) ds dy. \end{aligned}$$

From Lemma 2, X is a complete metric space. From Lemma 3, Lemma 4, Lemma 5, 6, 7, 8, the uniqueness and existence of a fixed point of Φ follow similarly to Banach's fixed point theorem ([5], [9]): There exists a unique $u \in X$ such that $\Phi[u] = u$.

Lemma 6 (Well-definedness of Φ). For $u \in S$, we have

$$\|E * (\chi_\Omega P(f - (u \cdot \nabla)u))\|_X < \infty.$$

Proof. Since $\|P\| \leq 1$, we obtain

$$\begin{aligned} \|E * (\chi_\Omega P(f - (u \cdot \nabla)u))\|_X &\leq C\|f\|_{X'} + C\|u^1 \partial_{x^1} u + u^2 \partial_{x^2} u + u^3 \partial_{x^3} u\|_{X'} \\ &\leq C\|f\|_{X'} + 3C^3 M^2 \\ &< \infty. \end{aligned}$$

□

Lemma 7 (Φ is a contraction mapping). The map $\Phi: S \rightarrow X$ is Lipschitz continuous; that is, there exists a constant $L > 0$ such that, for all $u, v \in S$, if $u - v \in S$, then

$$\left\| \int_{\mathbb{R} \times \mathbb{R}^3} E(s, y) (\chi_\Omega P((v \cdot \nabla)v - (u \cdot \nabla)u))(t - s, x - y) ds dy \right\|_X \leq L\|u - v\|_X.$$

Moreover, the constant L can be chosen so that $L < 1$.

Proof. See

$$\begin{aligned} &((v \cdot \nabla)v - (u \cdot \nabla)u)(t - s, x - y) \\ &= \sum_{j=1}^3 (v^j (\partial_{x^j} v - \partial_{x^j} u) + (v^j \partial_{x^j} u - u^j \partial_{x^j} u))(t - s, x - y), \end{aligned}$$

We take f as $\|f\|_{X'} \leq 1/(12C^4)$. From $\|f\|_{X'} \leq 1/(12C^4)$ and $C\|f\|_{X'} + 3C^3M^2 \leq M$, $2C^3M \leq (1 \pm \sqrt{1 - 12C^4})/3 < 1$.

We have

$$\begin{aligned} & \left\| \int_{\mathbb{R} \times \mathbb{R}^3} E(s, y) (\chi_\Omega P((v \cdot \nabla)v - (u \cdot \nabla)u))(t - s, x - y) ds dy \right\|_X \\ & \leq C^2 \|v\|_X \max_j \|\partial_{x^j}(v - u)\|_X + C^2 \|v - u\|_X \max_j \|\partial_{x^j} u\|_X \\ & \leq C^3 M \|v - u\|_X + C^3 M \|v - u\|_X \\ & \leq 2C^3 M \|u - v\|_X. \end{aligned}$$

Therefore, Lipschitz continuity follows for $L = 2C^3M$. $\square$

Lemma 8. Let $U \in X$. Then $\langle U, \varphi \rangle = 0$ for all $\varphi \in \mathcal{D}_\sigma(\Omega)$ if and only if there exists a distribution $\mathbf{p}$ such that $U = \nabla \mathbf{p}$.

Proof. For any $\varphi \in \mathcal{D}_\sigma(\Omega)$, $\operatorname{div}(\varphi) = 0$, so by integration by parts

$$\begin{aligned} \langle \nabla \mathbf{p}, \varphi \rangle &= \int_\Omega \sum_{i=1}^3 (\nabla \mathbf{p})^i(t, x) \varphi^i(t, x) dt dx \\ &= - \int_\Omega \mathbf{p}(t, x) \operatorname{div}(\varphi)(t, x) dt dx \\ &= 0. \end{aligned}$$

So, by Helmholtz decomposition, if $U = PU + \nabla \mathbf{p}$, then $\langle U, \varphi \rangle = \langle \nabla \mathbf{p}, \varphi \rangle = 0$. $\square$

Lemma 9 (Solvability of the Navier–Stokes equations). The fixed point u of $\Phi: S \rightarrow X$ is the solution of (N-S)'.

Proof. u satisfies $\operatorname{div}(u) = 0$ in the sense of a distribution belonging to $\mathcal{D}'(\Omega)$. That is, for any $\varphi \in D(\Omega)$,

$$\langle \operatorname{div}(u), \varphi \rangle = - \sum_{j=1}^3 \langle u^j, \partial_{x^j} \varphi \rangle = 0.$$

In fact, for any $u \in W_\sigma^{m,p}(\Omega)$, there exists a Cauchy sequence $\{u_n\} \subset V_\sigma^{m,p}(\Omega)$. By integration by parts and Hölder's inequality, we have:

$$0 = - \sum_{j=1}^3 \langle u_n^j, \partial_{x^j} \varphi \rangle \rightarrow - \sum_{j=1}^3 \langle u^j, \partial_{x^j} \varphi \rangle.$$

Boundedness of u , $\partial_{x^j} u$ by the Sobolev embedding theorem and $|\Omega| < \infty$, we have $(u \cdot \nabla)u \in L^2(\Omega)$. So, by the Helmholtz decomposition, if we let $f = Pf + \nabla \mathbf{f}$, $(u \cdot \nabla)u = P((u \cdot \nabla)u) + \nabla \mathbf{u}$ then for any $\varphi \in \mathcal{D}_\sigma(\Omega)$, $\langle f, \varphi \rangle = \langle Pf, \varphi \rangle$, $\langle (u \cdot \nabla)u, \varphi \rangle = \langle P((u \cdot \nabla)u), \varphi \rangle$, hence we solve

$$(N-S)' \quad \partial_t u - \Delta u = f - (u \cdot \nabla)u \quad \text{in } \mathcal{D}'(\Omega).$$

By Proposition 1, the solution of the approximate equation on Ω

$$(N-S)'' \quad \partial_t v_n - \Delta v_n = P(f - (u_n \cdot \nabla)u_n)$$

satisfies

$$v_n = u_{n+1} = E * (\chi_\Omega P(f - (u_n \cdot \nabla)u_n)).$$

Therefore, the solution of (N-S)'' satisfies

$$u_{n+1}(t, x) = \int_{\mathbb{R} \times \mathbb{R}^3} E(s, y) (\chi_\Omega P(f - (u_n \cdot \nabla)u_n))(t - s, x - y) ds dy.$$

$$(\partial_t - \Delta)u_{n+1}(t, x) = \langle (\partial_t - \Delta)E(t - s, x - y), (\chi_\Omega(Pf - (u_n \cdot \nabla)u_n))(s, y) \rangle$$

$$\begin{aligned} &= \langle \delta(\tau) \otimes \delta(z), (\chi_\Omega P(f - (u_n \cdot \nabla)u_n))(t - \tau, x - z) \rangle \\ &= (P(f - (u_n \cdot \nabla)u_n))(t, x). \end{aligned}$$

By Hölder's inequality, $\|P\| \leq 1$, and the continuity of the product of $L^2(\Omega) \times L^2(\Omega) \ni (u, v) \mapsto uv \in L^1(\Omega)$ (see Remark 3), we have

$$\begin{aligned} &\left| \int_{\Omega} (P((u_n \cdot \nabla)u_n - (u \cdot \nabla)u))(t, x) \cdot \varphi(t, x) dt dx \right| \\ &\leq \|(u_n \cdot \nabla)u_n - (u \cdot \nabla)u\|_{L^1(\Omega)} \|\varphi\|_{L^\infty(\Omega)} \\ &\rightarrow 0 \quad (n \rightarrow \infty) \end{aligned}$$

By the continuity of the heat operator on $\mathcal{D}'(\Omega)$, we have

$$|\langle (\partial_t - \Delta)u_n, \varphi \rangle - \langle (\partial_t - \Delta)u, \varphi \rangle| \rightarrow 0 \quad (n \rightarrow \infty),$$

and hence $(\partial_t - \Delta)u = P(f - (u \cdot \nabla)u)$ holds. From Lemma 7, we have

$$u(t, x) = \int_{\mathbb{R} \times \mathbb{R}^3} E(s, y) (\chi_\Omega P(f - (u \cdot \nabla)u))(t - s, x - y) ds dy.$$

u is a solution in the sense of the distribution in $\mathcal{D}'(\Omega)$ of (N-S)'. From Lemma 8, there exists $\mathbf{p}$ such that $(\partial_t - \Delta)u + (u \cdot \nabla)u - f = -\nabla \mathbf{p}$ holds. $\square$

4 Properties of the solutions

Lemma 10 (Vanishing).

$$\lim_{t, |x| \rightarrow \infty} \partial^\alpha u(t, x) = 0.$$

$$\begin{aligned}
& \forall u \in X', \lim_{\substack{t \rightarrow \infty \\ |x| \rightarrow \infty}} t^\gamma x^\beta \partial^\alpha u(t, x) = 0 \quad (\text{proof: similar to the proof of Lemma 10}) \\
& \forall \alpha, \beta, \gamma \in \mathbb{Z}_{\geq 0}, \quad \therefore \mathcal{S} \stackrel{\text{dense}}{\subset} X' \\
& \therefore \forall \varepsilon > 0, \forall u \in X', \exists u_n \in \mathcal{S}, \|u_n - u\|_{X'} < \varepsilon \\
& [\text{proof}] \quad \forall \varepsilon > 0, \forall u \in W^m, \exists u_n \in \mathcal{S}, \|u_n - u\|_{W^m} < \varepsilon \quad \therefore \exists C' > 0, \|u_n - u\|_{X'} \leq C' \varepsilon. //
\end{aligned}$$

Proof. u is a measurable function on $\mathbb{R} \times \mathbb{R}^3$, so we can take the limits as $t, |x| \rightarrow \infty$.

$$\partial^\alpha u(t, x) = \int_{\Omega} E(t-s, x-y) (\partial^\alpha (P(f - (u \cdot \nabla)u)))(s, y) ds dy,$$

for any $t_0 > 0$, if $t-s > t_0$ then $|E(t-s, x-y)| \leq 1/t^{3/2}$, $\partial^\alpha (P(f - (u \cdot \nabla)u)) \in X \subset C^{0,\varepsilon}(\Omega)$ so $\lim_{t, |x| \rightarrow \infty} \partial^\alpha u(t, x) = 0$ follows from the bounded convergence theorem. $\square$

The domain Ω can be taken arbitrarily large, so Lemma 10 can be thought of as a boundary condition.

Lemma 11 (Continuity of $f \mapsto u$). Let f_n, f be such that $Pf_n, Pf \in S$ and $\|f_n - f\|_{X'} \rightarrow 0$. Let u_n be the solutions corresponding to f_n with initial value $a_n \in A$, and let u be the solutions corresponding to f with initial value $a \in A$. Then $\|u_n - u\|_X \rightarrow 0$.

Proof.

$$\begin{aligned}
\|u_n - u\|_X &= \left\| \int_{\mathbb{R} \times \mathbb{R}^3} E(s, y) (\chi_\Omega P(f_n - (u_n \cdot \nabla)u_n))(t-s, x-y) ds dy \right. \\
&\quad \left. - \int_{\mathbb{R} \times \mathbb{R}^3} E(s, y) (\chi_\Omega P(f - (u \cdot \nabla)u))(t-s, x-y) ds dy \right\|_X \\
&\leq C \|f_n - f\|_X + 2C^3 M \|u_n - u\|_X.
\end{aligned}$$

So

$$\limsup_{n \rightarrow \infty} \|u_n - u\|_X \leq 2C^3 M \limsup_{n \rightarrow \infty} \|u_n - u\|_X.$$

$$\limsup_{n \rightarrow \infty} \|u_n - u\|_X \leq 2M.$$

Therefore

$$0 \leq (1 - 2C^3 M) \limsup_{n \rightarrow \infty} \|u_n - u\|_X \leq 0.$$

Hence

$$\limsup_{n \rightarrow \infty} \|u_n - u\|_X = \lim_{n \rightarrow \infty} \|u_n - u\|_X = 0.$$

$\square$

Proposition 3 (Elementary weak solutions as elements of Bochner class). Let $I = \{t \in \mathbb{R} : \exists x \in \mathbb{R}^3, (t, x) \in \Omega\}$, $\Omega' = \{x \in \mathbb{R}^3 : \exists t \in \mathbb{R}, (t, x) \in \Omega\}$. For $1 \leq p < \infty$,

$$a \in C^\infty(\Omega') \cap L_\sigma^p(\Omega'), u \in C(I; L_\sigma^p(\Omega')).$$

Proof. Since $u \in C^{0,\varepsilon}(\Omega)$, for any $\varepsilon > 0$, there exists $\delta > 0$ such that for all $(t, x), (t', x) \in \Omega$, if $|(t, x) - (t', x)| < \delta$ then $|u(t, x) - u(t', x)| < \varepsilon$. Therefore

$$\|u(t, \cdot) - u(t', \cdot)\|_{L^p(\Omega')} \leq |\Omega'|^{1/p} \varepsilon.$$

□

Lemma 12 (Elementary Weak Solutions are non-zero on the boundary of the domain). An elementary weak solution satisfies

$$u(t, x) = \int_{\mathbb{R} \times \mathbb{R}^3} E(s, y) (\chi_\Omega P(f - (u \cdot \nabla)u))(t - s, x - y) ds dy.$$

For each (t, x) , define a function $U^{(t,x)}$ of the variables (s, y) by

$$U^{(t,x)}(s, y) = (\chi_\Omega P(f - (u \cdot \nabla)u))(t - s, x - y).$$

If $\text{supp } U^{(t,x)} \subset \bar{\Omega}$ if and only if $\text{supp } u \subset \bar{\Omega}$, then u satisfy the energy inequality. (by integration by parts) Since $\Omega \subset \mathbb{R} \times \mathbb{R}^3$ is bounded, there exists $R > 0$ such that

$$(t, x) \in \Omega \implies |t|, |x| < R$$

In particular, if $|s| > R$ or $|y| > R$ then $(s, y) \notin \Omega$. Let $(t_0, x_0) \in \text{supp } U^{(t,x)} \cap \bar{\Omega}$. Then there exists $R' > 0$ such that

$$|t_0 - s| + |x_0 - y| < R' \implies (s, y) \in \text{supp } U^{(t,x)}.$$


On the other hand, we have


$$\begin{aligned} |t_0 - s| + |x_0 - y| &\geq ||t_0| - |s|| + ||x_0| - |y|| \\ &= ||s| - |t_0|| + ||y| - |x_0||. \end{aligned}$$

Hence, if $|s|$ and $|y|$ are sufficiently large (for instance, $|s| > R$ and $|y| > R$), then


$$|t_0 - s| + |x_0 - y| > R',$$

so, $(s, y) \notin \text{supp } U^{(t,x)}$, while $(s, y) \notin \Omega$. This shows that $\text{supp } U^{(t,x)} \subset \bar{\Omega}$ does not hold.





Construct of solutions
does not depend on a,
A.



□

5 Remarks

Remark 1. We can solve semi-linear partial differential equations with constant coefficients

$$Lu = f(u)$$

on $S' = \{u \in X' : \|u\|_{X'} \leq M\}$ similarly if L has locally integrable fundamental solutions. For example, the non-linear heat equation, the non-linear Schrödinger equation, the non-linear wave equation with 3-dimension, and the KdV equation.

Remark 2. As functions φ that $\operatorname{div}(\varphi) = 0$, it is sufficient to take any $\psi \in \mathcal{D}(\Omega)$ and set $\varphi = \operatorname{curl} \psi$.

Remark 3. Let $\|u_n - u\|_{L^2(\Omega)} \rightarrow 0$, $\|v_n - v\|_{L^2(\Omega)} \rightarrow 0$. By the triangle inequality, we have

$$\left| \|u_n\|_{L^2(\Omega)} - \|u\|_{L^2(\Omega)} \right| \leq \|u_n - u\|_{L^2(\Omega)}$$

for any sufficiently large n . On the other hand, $\|u_n\|_{L^2(\Omega)} < \|u\|_{L^2(\Omega)} + 1$. Therefore

$$\begin{aligned} \|u_n v_n - uv\|_{L^1(\Omega)} &\leq \|u_n\|_{L^2(\Omega)} \|v_n - v\|_{L^2(\Omega)} + \|v\|_{L^2(\Omega)} \|u_n - u\|_{L^2(\Omega)} \\ &< (\|u\|_{L^2(\Omega)} + 1) \|v_n - v\|_{L^2(\Omega)} + \|v\|_{L^2(\Omega)} \|u_n - u\|_{L^2(\Omega)} \\ &\rightarrow 0 \quad (n \rightarrow \infty). \end{aligned}$$

Remark 4. We change the assumptions: $0 \in I$, $0 \notin \Omega'$. Let $\rho(0, \cdot) \in X'(\Omega')$. We can solve

$$\begin{cases} \partial_t \rho + \operatorname{div}(\rho u) = 0 \\ \operatorname{Div}(D(u) + \operatorname{div}(u)(\delta^{ij}) - \mathbf{p}(\delta^{ij})) = f - \rho(\partial_t u + (u \cdot \nabla)u) \end{cases}$$

similarly. Here,

$$(\operatorname{Div} T)^i = \sum_{j=1}^3 \partial_{x^j} T^{ij}, D^{ij}(u) = \partial_{x^j} u^i + \partial_{x^i} u^j.$$

The operator $L_u: S' \ni \rho \mapsto \operatorname{div}(\rho u) \in X'$ is bounded, so

$$\rho = e^{-tL_u} \rho(0, \cdot).$$

Then

$$\operatorname{Div}(D(u) + \operatorname{div}(u)(\delta^{ij}) - \mathbf{p}(\delta^{ij})) = f - e^{-tL_u} \rho(0, \cdot)(\partial_t u + (u \cdot \nabla)u).$$

The fundamental solutions of the elliptic operators are locally integrable on new Ω . Using the above lemmas, the unique existence, smoothness, and properties of the solutions follow similarly.

Discription

This is an application of functional analysis to the existence and smoothness of the Navier–Stokes equations using elementary weak solutions in Sobolev spaces.

There are no long or complicated calculations; semi-groups, a priori estimates, and boundary conditions on $\Omega' \subset \mathbb{R}^3$ are not used at all. We apply the local solvability of linear partial differential operators with constant coefficients ([8], [3]), erase the pressure p by Helmholtz projection P , and solve

$$\begin{aligned} \frac{\partial u}{\partial t} - \Delta u &= Pf - P((u_n \cdot \nabla)u_n), \\ \frac{\partial u}{\partial t} - \Delta u &= Pf - P((u \cdot \nabla)u), \end{aligned}$$

in $X = \bigcap_{m \geq 5, p=1,2} W^{m,p}(\Omega)$, Ω is an open set in the universal space $\mathbb{R} \times \mathbb{R}^3$ that has a smooth boundary.

For example, cylinder $\Omega = I \times \Omega' = (-T, T) \times B_3(0, R)$ includes the earth. We handle the variables t, x simultaneously. Cylinders are Lipschitz domains, so we can use the Sobolev embedding theorem on Ω ([1]) and the Helmholtz decomposition on Ω' ([6]).

The initial value or external force term f is small as norms $\|f\|_X < \frac{1}{12C^4}$, but we can take Ω as (the earth) $\in \Omega$.

The non-zero examples of elements of X as a normed space do not depend on the choice of $c'_{k,m}$.

For any old $c'_{k,m}, c'_{k+1,m+1} \geq c'_{k,m} > 0$,

As a result, we can take c' as a positive constant: take c'' as 5000 c' , take $\alpha_{0,5}$ as 10^{-23} .

We can construct many non-zero solutions by choosing c' .

There exists $C = O(|\Omega|) > 0$, for example,

$$\begin{aligned} \|(x_i \mapsto \sin x_i)(x_j \mapsto \cos x_j)\|_X &\leq C\|\sin\|_X\|\cos\|_X, \\ \|(x_i \mapsto e^{\frac{1}{2}x_i})(x_j \mapsto e^{\frac{1}{3}x_j})\|_X &\leq C\|x_i \mapsto e^{\frac{1}{2}x_i}\|_X\|x_j \mapsto e^{\frac{1}{3}x_j}\|_X, \\ \|\partial_j^k e^x\|_X &\leq C^k\|e^x\|_X, \\ \|\partial_j^k \sin x\|_X &\leq C^k\|\sin x\|_X. \end{aligned}$$

for $i, j = 1, 2, 3$.

$$\|\partial_j u_n\|_X \leq CM$$

even if $n \geq 2$, and

$$\|\partial_j(u_{n+1} - u_n)\|_X \leq C\|u_{n+1} - u_n\|_X,$$

for $n \geq 1$.

So, there exists $0 < L < 1$, by mathematical induction, for $n \geq 2$,

$$\|u_{n+1} - u_n\|_X \leq L^{n-1}\|u_2 - u_1\|_X,$$

even if $u_n - u_{n-1} \notin S$ and $u_1 \neq \Phi[u_0]$.

Our map Φ is well-defined: the completion Sobolev space and the common Sobolev space are equivalent to each other, and the convolution of the functions does not depend on anything outside Ω .

(To construct the solutions,

$$\|\partial_j(u_{n+1} - u_n)\|_X \leq C\|u_{n+1} - u_n\|_X, \quad j = 1, 2, 3$$

It is not necessary, but we can easily calculate its use. The proof is similar to the construction of a unique solution under this description.

Therefore, existence follows similarly to the proof of Banach's fixed point theorem on a closed set $S \subset X_\sigma$. The solutions u, p satisfy

$$\frac{\partial u}{\partial t} - \Delta u = Pf - P((u \cdot \nabla)u)$$

in the sense that distributions in X_σ are

$$\operatorname{div}(u) = 0$$

in the sense that distribution is in X .

$f \neq 0 \implies u \neq 0$ and $f = 0, a = 0 \implies u = 0$. There exists a non-zero solution by Fujita, Kato ([4]) or Shibata ([6]), even if $f = 0$, so elementary weak solutions are different from the solutions by Fujita, Kato or Shibata.

$u|_{\partial\Omega'} \neq 0$, but by Helmholtz decomposition, if $\Omega = I \times \Omega' = (-T, T) \times B_3(0, R)$, then $(u \cdot n)|_{\partial\Omega'} = 0$, so our solutions satisfy the energy inequality on $\Omega = (-T, T) \times B_3(0, R)$. By integration by parts, we obtain

$$(\partial_t u - \Delta u) \cdot u = (f - \nabla p - (u \cdot \nabla)u) \cdot u.$$

Hence

$$\frac{1}{2} \partial_t (u \cdot u) - (\Delta u) \cdot u = f \cdot u - ((u \cdot \nabla)u) \cdot u.$$

Integrating over $B = B_3(0, R)$ and $t \in (0, t)$, we obtain

$$\begin{aligned} \frac{1}{2} \|u(t)\|_{L^2(B)}^2 - \int_0^t \int_{\partial\Omega'} |u \cdot n| (1 + |\nabla u|^2) dA ds + \int_0^t \|\nabla u(s)\|_{L^2(B)}^2 ds \\ \leq \frac{1}{2} \|u(0)\|_{L^2(B)}^2 + \int_0^t (f(s), u(s))_{L^2(B)} ds. \\ \frac{1}{2} \|u(t)\|_{L^2(B)}^2 + \int_0^t \|\nabla u(s)\|_{L^2(B)}^2 ds \\ \leq \frac{1}{2} \|u(0)\|_{L^2(B)}^2 + \int_0^t (f(s), u(s))_{L^2(B)} ds. \end{aligned}$$

Our solutions are smooth, unique, u is defined on $\mathbb{R} \times \mathbb{R}^3$, and

$$\lim_{t, |x| \rightarrow \infty} \partial^\alpha u(t, x) = 0,$$

for any multi-index α . The map $f \mapsto u$ is continuous. The initial value satisfies

$$a \in C^\infty(\Omega') \cap L^p(\Omega').$$

The solution satisfies

$$u \in C(I; L^p(\Omega')).$$

If

$$\int_0^T (f(s), \varphi(s))_{L^2(\Omega')} ds + (a, \varphi(0))_{L^2(\Omega')} = 0$$

for all test functions φ , and

$$u|_{\partial\Omega'} = 0,$$

then

$$f = 0, \quad a = 0, \quad u = 0.$$

satisfy the energy inequality and u belongs to the Serrin class. So u is the unique Leray-Hopf's weak solution and an elementary weak solution.

Therefore, if u is the Leray-Hopf's weak solution and an elementary weak solution, then $u = 0$.

If we change the definition of S by

$$S = \left\{ u \in X : \|u\|_X \leq M, \quad \left\| \partial_j^k u \right\|_X \leq C^k \|u\|_X, \quad j = 1, 2, 3, k \geq 0 \right\},$$

then the existence follows exactly.

$$\partial_j : X \rightarrow W^{m,p}(\Omega)$$

is a closed operator, so S is a closed set by the proof using Fatou's lemma and

$$\left| \|\partial_j u_n\|_X - \|\partial_j u\|_X \right| \leq \|\partial_j u_n - \partial_j u\|_X \rightarrow 0.$$

(similar to the proof of Lemma 2)

Uniqueness follows. For $f \neq 0$, we can construct many non-zero solutions for which uniqueness holds by Baire's category theorem: construct $u \in S$ as an interior point in X_σ , construct the open neighborhood B_u , and think $v \in S \cap B_u$ satisfies $\|u - v\| < M$, and prove similarly to "Uniqueness" for $u - v$ by using C_n , $C_n \rightarrow \infty$, $u_{n(k)} \rightarrow u$, and $v_{n(k)} \rightarrow v$ from

$$\begin{aligned} \|u - u_{n(k)}\|, \|v - v_{n(k)}\| &< \frac{1}{k}, \\ C_{n(k)} \|u_{n(k)} - v_{n(k)}\| &\leq \|\partial_j(u_{n(k)} - v_{n(k)})\| \\ &= \|\partial_j u_{n(k)} - \partial_j v_{n(k)}\| \\ &\leq \|\partial_j u_{n(k)}\| + \|\partial_j v_{n(k)}\| \\ &\leq 2CM. \end{aligned}$$

We conjecture that blow-up solutions can also be constructed. For any $\varepsilon > 0$, there exist $\delta > 0$, a constant $M(\varepsilon) > 0$, and a function $f \in S_{M(\varepsilon)}$ such that:

- $M(\varepsilon_1) > M(\varepsilon_2)$ whenever $\varepsilon_1 > \varepsilon_2$,
- $CM(\varepsilon) > \|E * \chi_\Omega P f\| > 2\varepsilon$,
- $|\Omega| < \delta \implies \|E * \chi_\Omega P((u \cdot \nabla)u)\| < \varepsilon$.

By the triangle inequality, $\|u\| = \|\Phi[u]\| > \varepsilon$. For example,

$$\begin{aligned} f(t, x) &= 2(\varepsilon + 1)(\|E * \chi_\Omega P \sin(x_1)\| + 1) \sin(x_1), \\ M(\varepsilon) &= 2 + \exp(2(\varepsilon + 1)(\|E * \chi_\Omega P \sin(x_1)\| + 1)). \end{aligned}$$

We conjecture that we can take the initial value a large by making $|\Omega|$ smaller.

About other PDEs

$$\begin{aligned}
 u &= u(t, x) \quad \lambda = \lambda(t, x), \exists M_\lambda > 0, \\
 f &= f(t, x) \quad |\lambda(t, x)| \leq M_\lambda, \forall (t, x) \in \Omega, \\
 \partial_t u - \Delta u &= f + \lambda u^2 \\
 \partial_t^2 u - \Delta u &= f + \lambda u^2 \\
 i \partial_t u + \Delta u &= f + \lambda |u|^2 u \\
 \partial_t^2 u + \partial_t u - \Delta u &= f + \lambda u^2 \\
 \partial_t^2 u - \Delta u + m_0 u &= f + \lambda u^2 \\
 \Delta u &= f + \lambda u^2 \\
 \Delta u + m_0 u &= f + \lambda u^2
 \end{aligned}$$

We can apply our Elementary PDE Theory to the wave equations with dimension 2, by Theorem 10.2.1., Lars Hörmander, "The Analysis of Linear Partial Differential Operators II", Springer, 1983.

Compact Kähler manifold (X, g)
 Complex Monge-Ampère equation
 $\det(g_{\mu\bar{\nu}} + \partial_\mu \partial_{\bar{\nu}} u) = e^{-cu+f} \det(g_{\mu\bar{\nu}})$ (u : unknown, f : known)
 $\Leftrightarrow -c \partial_{\bar{\nu}} u = -\partial_{\bar{\nu}} f + \partial_{\bar{\nu}} \log \det(g_{\mu\bar{\nu}} + \partial_\mu \partial_{\bar{\nu}} u) - \partial_{\bar{\nu}} \log \det(g_{\mu\bar{\nu}})$
 $E(z) = \frac{1}{\pi z_{\bar{z}}} \Rightarrow \partial_{\bar{z}} E = \delta$
 $\therefore u = \frac{1}{c} \frac{1}{\pi z_{\bar{z}}} * (\partial_{\bar{z}} f + \partial_{\bar{z}} \log \det(g_{\mu\bar{\nu}}) - \partial_{\bar{z}} \log \det(g_{\mu\bar{\nu}} + \partial_\mu \partial_{\bar{\nu}} u))$
 Reference: 中島 啓 (Hiraku Nakajima)
 非線形問題と複素幾何学
 (Non-linear Problem and Complex Geometry)
 岩波書店 (Iwanami books), 2008
 ISBN: 978-4000058803

Application to complex geometry.

We define the convergencing $a_n \rightarrow a$ by "for any $F \in \mathcal{E}'(\Omega)$, $|\langle F, |u_n - u| \rangle| \rightarrow 0$ ". $F(t, x) = \delta(t) \otimes 1(x) \Rightarrow \|a_n - a\|_{L^1(\Omega')} \rightarrow 0, F(t, x) = 1 \Rightarrow \|u_n - u\|_{L^1(\Omega)} \rightarrow 0$. The map to an unique solution $a \mapsto u$ is continuous. Mazur's lemma does not hold on A .

If we define the metric on A by setting $d_A(u(0), v(0)) = \|u - v\|_X$, then the map to unique solutions $a \mapsto u$ is also continuous.

From the proof of Sobolev embedding theorem, we can take a constant C_4 as does not depend on k and for any $\omega = \omega^\circ$, $\bar{\omega} \subset \Omega$, $\rho \in \mathcal{D}(\Omega)$, $\rho(\omega) = \{1\}$,

$$\|\rho u\|_{C^k(\Omega)} \leq \|\rho u\|_{C^k(R \times R^3)} \leq C_4 \|\rho u\|_{W^m(R \times R^3)} \leq C_4 / \left(\sum_{m \geq 5, k=0}^{\infty} 1/(m!^4 c_{k,m}'^2) \right) \|u\|_X.$$

ρ can be taken arbitrarily, so we can take $c_{k,m}''$ as a constant.

1 Reference

New Elementary Sobolev Embedding Theorem without Extension Operators,
Masatoshi Ohnui, 2026.

We can take $c_{k,m}''$ that do not depend on k, m , therefore our solutions are analytic: $\sup_{\bar{\Omega}} |\partial^\alpha u| \leq c_{0,5}$, thus the remainder term converges to 0. $X \subset \mathcal{E}$ and by Egorov's theorem, the dual space $X' \subset \mathcal{E}'$.

New Fixed Point Theorem

Theorem: let S be a closed subset of a Banach space X , $\Phi : S \rightarrow X$ and there exists $0 < L < 1$ such that $\|\Phi[u] - \Phi[v]\| \leq \|u - v\|$ for any $u, v \in S$. If we can define a sequence $u_0, u_1, u_{n+1} = \Phi[u_n]$, then it converges to the limit $u = \Phi[u] \in X$ even if $\Phi[S] \not\subset S$ and $u = \Phi[u], v = \Phi[v], u - v \in S \Rightarrow u = v$. The proof: it is similar to Banach Fixed-Point Theorem. S is closed, hence $u \in S$. By continuity of Φ at u , we obtain $u = \Phi[u]$.


By assumption, $u_n \in S, \forall n$. $u \in S \Rightarrow \Phi[u] \in S$. The proof is similar to the construction of unique solutions.

Supplement

$$\Delta u \cdot u = \sum_{j=1}^3 \partial_{x^j}^2 u \cdot u, \sum_{i,j=1}^3 \partial_{x^j} u^i \cdot u^i n^i \leq (1 + |\nabla u|^2) |u \cdot n|.$$

Acknowledgements

Parts of this work were developed through discussions and materials shared on x.com (account: @Ohnui_PDE_LowD), as well as on mathlog.info and note.com, and through activities at the Tokyo University of Science. We are grateful to the many individuals who provided feedback and support for this research during the period from February 5, 2022, to December 30, 2025. In particular, this manuscript has been read by Naofumi Mori (@n_mori00), Associate Professor of Mathematical Sciences, as well as by Japanese users of X, @R1sFs, @ksvzvsk, and @sbakbao.




The graph of $E = E(t, x)$.

Completion of normed spaces.


References

- [1] Robert A. Adams and John J. F. Fournier, *Sobolev Spaces*, Academic Press, 2003.
- [2] Frédéric Charve and Raphaël Danchin, “A Global Existence Result for the Compressible Navier–Stokes Equations in the Critical L^p Framework,” *Archive for Rational Mechanics and Analysis*, **198** (2010), 233–271.
- [3] Lars Hörmander, *The Analysis of Linear Partial Differential Operators I: Distribution Theory and Fourier Analysis*, Springer Berlin, Heidelberg, 1990.
- [4] Tosio Kato, “Strong L^p -Solutions of the Navier–Stokes Equation in R^m , with Applications to Weak Solutions,” *Mathematische Zeitschrift*, **187** (1984), 471–480.
- [5] A. N. Kolmogorov and S. V. Fomin, *Elements of the Theory of Functions and Functional Analysis*, Dover Publications, 1999.
- [6] Yoshihiro Shibata and Miho Murata, “On the global well-posedness for the compressible Navier–Stokes equations with slip boundary condition,” *Journal of Differential Equations*, **260** (2016), no. 7, 5761–5795.
- [7] Wasao Sibagaki and Hisako Rikimaru, “On the E. Hopf’s Weak Solution of Initial Value Problem for the Navier–Stokes Equations,” *Memoirs of the Faculty of Science, Kyushu University. Series A, Mathematics*, **21** (1967), no. 2, 194–240.
- [8] Elias M. Stein and Rami Shakarchi, *Functional Analysis: Introduction to Further Topics in Analysis*, Princeton University Press, 2011.
- [9] Wataru Takahashi, *Nonlinear Functional Analysis: Fixed Point Theorems and Related Topics*, Kindaikagaku-sha, 2000.
- [10] Kenji Yajima, *Lebesgue Integral and Functional Analysis*, Asakura Publishing Co., Ltd., 2015.
- [11] Kōsaku Yosida, *Functional Analysis*, Springer Berlin, Heidelberg, 1980.

- [12] Ping Zhang, “Global Fujita–Kato solution of 3-D inhomogeneous incompressible Navier–Stokes system,” *Advances in Mathematics*, **363** (2020), 107007.
- [13] Haim Brezis, *Functional Analysis, Sobolev Spaces and Partial Differential Equations*, Springer, 2010.
- [14] Hisashi Okamoto, *Principal of the Navier–Stokes Equations*, University of Tokyo Press, 2023. ISBN: 978-4-13-061315-6.
- [15] Yoshihiro Shibata, *Basic of Mathematical Fluids 1, 2*, Iwanami Books, 2022. ISBN: 9784000298582, 9784000298599.
- [16] Lars Hörmander, *The Analysis of Linear Partial Differential Operators II*, Springer, 1983.
- [17] Hiraku Nakajima, *Non-linear Problem and Complex Geometry*, Iwanami Books, 2008. ISBN: 978-4000058803.
- [18] Y. Shibata and T. Kakita, *From Vector Calculus to Fluids*, Japan Review Company, 2024. ISBN: 978-4-535-78962-3.



Convolution.



Key inequalities.